

Expression of Osteopontin in Synovial Tissue of Rheumatoid Arthritis: Its Role in Joint Destruction

Thesis

Submitted for Fulfillment of the Requirements for the MD Degree in Physical Medicine and Rehabilitation

By

Nermin Farouk Labib

M.B., B. Ch., M. Sc.

Under supervision of

Professor Doctor/ Nahed Mounir Sherif

*Professor of Physical Medicine and Rehabilitation
Faculty of Medicine, Ain Shams University*

Professor Doctor/ Hala Abdul-Hady Saleh

*Professor of Physical Medicine and Rehabilitation
Faculty of Medicine, Ain Shams University*

Professor Doctor/ Omar Hussein Omar

*Professor of Radiodiagnosis
Faculty of Medicine, Ain Shams University*

Professor Doctor/ Nedal Ahmed Hegazy

*Professor of Pathology
Faculty of Medicine, Ain Shams University*

Doctor/ Iman Mahmoud Ghanima

*Assistant professor of Physical Medicine and Rehabilitation
Faculty of Medicine, Ain Shams University*

**Faculty of Medicine,
Ain Shams University**

٢٠٠٨

Acknowledgement

First and foremost, thanks to **ALLAH**, the most beneficent and most merciful.

I find no words by which I can express my sincere thanks, supreme gratitude to **Prof. Dr. Nahed Mounir Sherif**, Professor of Physical Medicine, Rheumatology and Rehabilitation, Faculty of Medicine, Ain Shams University. It was an honor to me to carry out this work under her continuous guidance, patience, encouragement and expert supervision.

I would like also to express my gratitude to **Prof. Dr. Hala Abdul Hady Saleh**, Professor of Physical Medicine, Rheumatology and Rehabilitation, Faculty of Medicine, Ain Shams University. For her unlimited help, precise instructions, kind encouragement and directions throughout this work.

I would like also to express my gratitude to **Prof. Dr. Omar Hussein Omar**, Professor of Radiodiagnosis, Faculty of Medicine, Ain Shams University, for his instructions, unlimited help throughout this work.

I am deeply grateful to **Prof. Dr. Nedal Ahmed Hegazy**, Professor of Pathology, Faculty of Medicine, Ain Shams University, who devoted her time, effort and experience most generously throughout this work.

I am greatly indebted to **Dr. Iman Mahmoud Ghanima**, Assistant Professor of Physical Medicine, Rheumatology and Rehabilitation, Faculty of Medicine, Ain Shams University, for her great help and guidance throughout this work.

I am grateful to the orthopedic surgery professors for their help which was essential for accomplishment of this work.

I would like to thank all our patients who were involved in this study for their cooperation and understanding of the importance of our research work and for their friendly attitude.

Also, I would like to thank all my Professors and colleagues in Physical Medicine, Rheumatology and Rehabilitation, Faculty of Medicine, Ain Shams University, for their help and support.

Finally, I wish to thank everyone who offered me help during the achievement of this research work especially my parents and husband whom with their continuous enthusiastic encouragement and everlasting help made the completion of this work possible.

Nermin Farouk Labib

List of Contents

	<i>Page</i>
• List of abbreviations	ii
• List of Figures	vi
• List of Tables	x
• Introduction & aim of the work	1
• Review of Literature:	
➤ Rheumatoid arthritis	4
➤ Osteopontin	97
• Patients and methods	129
• Results	142
• Discussion	183
• Summary and conclusion	209
• Recommendations	213
• References	214
• Arabic summary	

List of Abbreviations

ACR	American college of rheumatology
AFA	antifilaggrin antibodies
AIMs	Arthritis impact measurements Scales
AKA	Antikeratin antibody
ALP	Alkaline phosphatase
ANA	Antinuclear antibodies
ANCA	Antineutrophil cytoplasmic antibodies
Ang-λ	Angiopoietin- λ
Anti-CII ...	Anticollagen type II
APF	Antiperinuclear factor
Bcl-γ	B cell lymphoma- γ
BiP	Binding protein
BMD	Bone mineral density
Cλ-γ	Cervical λ - γ
CAM	Chorioallantoic membrane
CCPs	Cyclic citrullinated peptides
CCR	chemokine receptor (γ adjacent cysteines)
CD	Cluster of differentiation
cDNA	Complementary deoxy ribonucleic acid
CFD	Color flow doppler
COX	Cyclooxygenase
CPPD	Calcium pyrophosphate dehydrate
CR	Conventional radiography
CRP	C-reactive protein
CT	Computed tomography
CTLA-ξ ...	Cytotoxic T-lymphocyte associated antigen ξ
CXC	Chemokine (γ cysteines separated by X amino acid)
DAB	Diaminobenzidine
DAS	Disease activity score
DEXA	Dual Energy X ray Absorptiometry
DIP	Distal interphalangeal
DMARDs ..	Disease modified antirheumatic drugs
ds-DNA ...	Double stranded deoxyribonucleic acid
DTH	Delayed type hypersensitivity
EAE	Experimental autoimmune encephalomyelitis
EBV	Epstein bar virus
ECM	Extracellular matrix

EDTA	Ethylenediaminetetraacetic acid
ELISA	Enzyme linked immunosorbent assay
ELR	Glycine-leucine-arginine amino acid sequences
ENA	Epithelial neutrophil activating protein
ERM	Ezrin, radixin, moesin proteins
ESR	Erythrocyte sedimentation rate
Eta-λ	Early T-cell activation gene λ
Fc	Fragment-crystalline
FGF	Fibroblast growth factor
FLS	Fibroblast like synoviocytes
fMLP	Formyl methionyl-leucyl-phenylalanine
Gly	Glycolysation
GM-CSF ..	Granulocyte macrophage colony stimulating factor
GPI	Glucose- γ -phosphate isomerase
GRO	Growth related oncogene
HAQ	Heath assessment questionnaire
Hb	Hemoglobin
HIF	Hypoxia inducible transcription factor
HLA	Human leukocytic antigen
HMG	High mobility group protein
HPF	High power field
HRUS	High resolution ultrasound
IAP	Inhibitor of apoptosis
ICAM	Intercellular adhesion molecule
IFN-γ	Interferon γ
Ig	Immunoglobulin
IL	Interleukin
iNOS	Inducible nitric oxide synthetase
IP	Inducible protein
I-TAC	Inducible T cell alpha chemoattractant
JAK-STAT	Janus kinases, signal transducers and activators of transcription
JIA	Juvenile idiopathic arthritis
kDa	Kilodalton
KO mice ..	Knock out mice
LPS	Lipopolysaccharides
Lt	Left
MACTAR	McMaster Toronto arthritis
MAdCAM	Mucosal addressin cell adhesion molecule

Max	Maximum
MCP	Metacarpophalangeal
MCP-1	Monocyte chemoattractant protein-1
MDHAQ ..	Multidimensional health assessment questionnaire
Mg	Milligram
Mg²⁺	Magnesium
MHAQ	Modified health assessment questionnaire
MHC	Major histocompatibility complex
MHz	Megahertz
Mig	Monokine induced by interferon gamma
Min	Minimum
MLS	Macrophage like synoviocyte
Mm	Millimeter
MMP	Matrix metalloproteinase
Mn²⁺	Manganese
MRI	Magnetic resonance imaging
mRNA	Messenger ribonucleic acid
MSUS	Musculoskeletal ultrasound
MTP	Metatarsophalangeal
MTX	Methotrexate
NF-κB	Nuclear factor kappa beta
NHP	Nottingham health profile
NO	Nitric oxide
NS	Non significant
NSAIDs ...	Nonsteroidal anti-inflammatory drugs
OA	Osteoarthritis
OPN	Osteopontin
PBMC	Peripheral blood mononuclear cell
PBS	Phosphate buffered saline
PDGF	Platelet derived growth factor
PDUS	Power doppler ultrasonography
PECAM ...	Platelet-endothelial cellular adhesion molecule
PF	Platelet factor
PGA	Patient's global assessment
PGE²	Prostaglandin E ²
PIP	Proximal interphalangeal
Plt	Platelets
PMNLs ...	Polymorphnuclear leukocytes
PVP	Polyvinyl pyrrolidone
PXR	Plain X ray

r	Correlation coefficient
RA	Rheumatoid arthritis
RANA	Rheumatoid arthritis associated nuclear antigen
RANK	Receptor of activation of nuclear factor
RANTES .	Regulated on activation normally T cell expressed and secreted
RF	Rheumatoid factor
RGD	Arginine-glycine-aspartic amino acid sequences
rOPN	Recombinant OPN
Rt	Right
RT-PCR ..	Reverse transcriptase polymerase chain reaction
S	Significant
SD	Standard deviation
SDF-1	Stromal cell derived factor-1
SE	Shared epitope
SFMC	Synovial fluid mononuclear cell
SFs	Synovial fibroblasts
Sig.	Significance
SLAYGLR	Ser-Leu-Ala-Tyr-Gly-Leu-Arg amino acid sequences
SLE	Systemic lupus erythematosus
SLEDAI ..	Systemic lupus erythematosus disease activity index
SSI	Spread Severity Index
SVVYGLR	Ser-Val-Val-Tyr-Gly-Leu-Arg amino acid sequences
TGF-B	Transforming growth factor beta
Th	T helper
Tie	Receptor tyrosine kinase
TIMPs	Tissue inhibitor matrix metalloproteinase
TMJ	Tempromandibular joint
TNF	Tumor necrosis factor
TUNEL	Terminal deoxynucleotidyl transferase dUTP nick end labeling
UC	Ulcerative colitis
US	Ultrasound
VAS	Visual analogue scale for pain
VCAM	Vascular cell adhesion molecule
VEGF	Vascular endothelial growth factor
VLA	Very late antigen

List of Figures

Figure NO.	Figure title	Page NO.
Figure (١):	Outline of the management of rheumatoid arthritis	٨٤
Figure (٢):	Some of the features of the OPN molecule	٩٨
Figure (٣):	Pro- and anti-inflammatory actions of OPN in cell injury	١١٣
Figure (٤):	Shows HRUS apparatus	١٣٥
Figure (٥):	Shows suprapatellar longitudinal application in moderate knee flexion	١٣٧
Figure (٦):	Shows sex distribution among RA patients....	١٤٣
Figure (٧):	Shows modified disease activity score (DAS) among RA patients	١٤٤
Figure (٨):	Shows the distribution of different extra-articular manifestations of RA patients	١٤٦
Figure (٩):	A comparison between patient and control groups regarding the laboratory data	١٤٨
Figure (١٠):	Shows the frequency of the radiological grades among patients under study	١٤٩
Figure (١١):	Longitudinal sonographic scan of the right knee joint in a RA patient showing, thickened irregular synovium situated between the femur (F) and the quadriceps tendon (T) and the thickness of the synovium was determined	١٥١
Figure (١٢):	Longitudinal sonographic scan of the right knee joint in a RA patient showing decreased in cartilage thickness	١٥١

Figure (١٣): Longitudinal sonographic scan of the right knee joint in a RA patient showing, bone erosions	١٥٢
Figure (١٤): Power Doppler sonographic image of the left knee joint in a RA patient showing no increase in synovial blood flow (power Doppler grade ٠)	١٥٢
Figure (١٥): Longitudinal sonographic scan of the right knee joint in a RA patient showing increased synovial blood flow (power Doppler grade ١).	١٥٣
Figure (١٦): Longitudinal sonographic scan of the left knee joint in a RA patient showing increased synovial blood flow (power Dopple grade ٢).	١٥٣
Figure (١٧): Power Doppler sonographic scan of the right knee joint in a RA patient showing, hypertrophic synovium and its vascularization. The flow pattern of the synovial blood flow was determined and the resistive index (RI) was calculated automatically	١٥٤
Figure (١٨): Shows comparison between RA patients group and controls group regarding synovial blood flow grades	١٥٥
Figure (١٩): Shows comparison of high resolution ultrasound data between RA patients group and control group under study	١٥٦
Figure (٢٠): Control synovial tissue showing weak minimal immunoreactivity for OPN in synovial lining and weak scattered	

	perivascular staining in the sublining layer (X ¹ ··)	161
Figure (21):	Control synovial tissue showing weak minimal immunostaining for OPN in synovial lining cells (X ² ··)	162
Figure (22):	Showing expression of OPN in RA synovium: OPN staining is located predominantly in synovial lining rather than the sublining (X ¹ ··)	162
Figure (23):	Showing expression of OPN in RA synovium: synovial lining cells stain strongly for OPN while no staining is detected in the lymphoid aggregates (X ² ··)	163
Figure (24):	RA synovial sublining tissue showing less remarkable OPN expression, mainly located in the perivascular extracellular tissue and some scattered inflammatory cells. It is absent in the lymphoid aggregates (X ² ··)	163
Figure (25):	RA synovial tissue showing strong cytoplasmic staining for OPN (+++) in lining synoviocytes (X ² ··)	164
Figure (26):	RA synovial tissue showing strong cytoplasmic staining for OPN (+++) in lining synoviocytes more than the sublining layer (X ² ··)	164
Figure (27):	RA synovial tissue showing intermediate cytoplasmic staining for OPN (++) in lining synoviocytes (X ² ··)	165

Figure (٣٨): The correlation between synovial thickness and bone erosion	١٦٩
Figure (٣٩): The correlation between synovial thickness and cartilage thickness	١٦٩
Figure (٣٠): The correlation between synovial thickness and spread severity index	١٧٠
Figure (٣١): The correlation between synovial thickness and resistive index (RI)	١٧٠
Figure (٣٢): The correlation between resistive index and disease activity score	١٧٤
Figure (٣٣): The correlation between resistive index and spread severity index	١٧٤
Figure (٣٤): The correlation between OPN positive cells and DAS	١٧٦
Figure (٣٥): The correlation between OPN positive cells and spread severity index	١٧٦
Figure (٣٦): The correlation between OPN positive cells and synovial thickness	١٧٧
Figure (٣٧): The correlation between OPN positive cells and bone erosion	١٧٧
Figure (٣٨): The correlation between OPN positive cells and cartilage thickness	١٧٨
Figure (٣٩): The correlation between OPN positive cells and RI	١٧٨

List of Tables

Table No.	Table Title	Page No.
Table (1):	Shows major adhesion molecule interactions with rheumatoid arthritis synovial endothelium	27
Table (2):	Shows RA autoantigens with relatively high specificity	53
Table (3):	Shows the benefits of MSUS for rheumatologists	60
Table (4):	Shows the demographic data of both groups under study	142
Table (5):	Shows the clinical data of RA patients	143
Table (6):	Shows the classification of RA patients according to their disease activity	144
Table (7):	Shows modified index of synovitis activity of the knee joint among RA patients under study	145
Table (8):	Shows the frequency of different extra-articular manifestations of RA patients	146
Table (9):	Shows laboratory data among patient group	147
Table (10):	Shows laboratory data among control group	147
Table (11):	Shows comparison of laboratory data between RA patients group and control group under study	148
Table (12):	Shows the radiological data among patients under study	149

Table (١٣): Shows comparison between RA patients group and control group regarding synovial blood flow grades	١٥٥
Table (١٤): Shows comparison between RA patients group and control group regarding other HRUS findings	١٥٦
Table (١٥): Shows the comparison between RA patients with positive RF and those with negative RF as regards HRUS findings	١٥٧
Table (١٦): Shows the comparison between RA patients with moderate modified DAS and those with severe modified DAS as regards HRUS findings	١٥٨
Table (١٧): Shows grades of modified index of knee synovitis activity in relation to HRUS findings of RA patients group	١٥٩
Table (١٨): Shows the different radiological grades according to Larsen's score in relation to HRUS results of RA patients group	١٦٠
Table (١٩): Shows a comparison between patients and controls regarding the immuno-histochemical findings	١٦٥
Table (٢٠): Shows the comparison between RA patients with positive rheumatoid factor (RF) and those with negative RF as regards the immunohistochemical data	١٦٦
Table (٢١): Shows a comparison between grades of modified DAS and the immuno-histochemical findings	١٦٦
Table (٢٢): Shows the modified index of synovitis activity grades of the knee joint in relation	

to the immunohistochemical data among RA patient group	167
Table (23): Shows the different radiological grades according to Larsen's score in relation to immunohistochemical data among RA patient group	167
Table (24): Shows correlation between synovial thickness and various clinical, laboratory, radiological and immunohistochemical parameter of RA patients group	168
Table (25): Shows correlation between cartilage thickness and various clinical, laboratory, radiological and immunohistochemical parameter of RA patients group	171
Table (26): Shows correlation between bone erosion and various clinical, laboratory, radiological and immunohistochemical parameter of RA patients group	172
Table (27): Shows correlation between resistive index (RI) and various clinical, laboratory, radiological and immunohistochemical parameter of RA patients group	173
Table (28): Shows correlation between percentage of OPN positive cells and various clinical, laboratory and radiological parameters of RA patients group	175
Table (29): Shows synovial blood flow grades in relation to various clinical data of RA patient	179