

**ROLE OF DIALYZER REPROCESSING IN
TRANSMISSION OF HCV INFECTION**

Thesis
Submitted for Partial Fulfillment of The Master Degree
In
Internal Medicine

Presented By
Khalid Mahmoud Makboul
M.B., B.Ch.

Under Supervision of
Prof. Dr. Badawy Labib Mahmoud
Professor of Internal Medicine and Nephrology
Faculty of Medicine
Ain Shams University

Prof. Dr. Ali Khalifa Ali
Professor of Biochemistry
Faculty of Medicine
Ain Shams University

Dr. Hesham Mohamed El-Sayed
Lecturer of Internal Medicine and Nephrology
Faculty of Medicine
Ain Shams University

Faculty of Medicine
Ain Shams University
1998



بِسْمِ اللَّهِ الرَّحْمَنِ الرَّحِيمِ

قَالُوا سُبْحَانَكَ لَا عِلْمَ لَنَا إِلَّا
مَا عَلَّمْتَنَا إِنَّكَ أَنْتَ الْعَلِيمُ
الْحَكِيمُ

صَدَقَ اللَّهُ الْعَظِيمُ

سورة البقرة الآية ٢٢

Acknowledgment

Before all, I would like to express my deep thanks to **ALLAH** without his great blessings, I would never accomplish this work.

I wish to express my sincere and deep appreciations to Prof.Dr. Badawy Labib, Professor of Internal Medicine and Nephrology, Ain Shams University, for his continuous guidance, paternal support and constructive criticism through the work.

Also, my profound gratitude to Prof.Dr. Aly Khalifa Aly, Professor of Biochemistry, Ain Shams University, for his kind supervision and support.

I am also indebted to Dr. Hesham Mohamed El-Sayed, Lecturer of Internal Medicine and Nephrology, Ain Shams University, without his sincere help this work could not be accomplished.

Many thanks to Prof. Dr. Mahmoud Ismael, Professor of Biochemistry, Ain Shams University for his great help

Special thanks are dedicated to Dr. Abdul Basset El-Shaarawy, Lecturer of

Internal Medicine and Nephrology, Ain Shams University, and to my colleague Dr. Saeed Abdul Wahab, Assistant Lecturer of Internal Medicine and Nephrology, Ain Shams University, for their kind help.

TABLE OF CONTENTS

| | |
|---|-------------------|
| <i>List of tables</i> | <i>i</i> |
| <i>List of figures</i> | <i>ii</i> |
| <i>List of abbreviations</i> | <i>iii</i> |
| <i>INTRODUCTION AND AIM OF THE WORK</i> | <i>1</i> |
| <i>REVIEW OF LITERATURE</i> | <i>3</i> |
| <i>Viral Hepatitis</i> | <i>3</i> |
| □ <i>Hepatitis A Virus</i> | <i>4</i> |
| □ <i>Hepatitis B Virus</i> | <i>5</i> |
| □ <i>Hepatitis D Virus</i> | <i>8</i> |
| □ <i>Hepatitis E Virus</i> | <i>9</i> |
| □ <i>Hepatitis G Virus</i> | <i>10</i> |
| □ <i>Hepatitis C Virus</i> | <i>11</i> |
| <i>Hepatitis C Virus And Haemodialysis Patients.</i> | <i>29</i> |
| <i>Dialyzer Reuse</i> | <i>39</i> |
| □ <i>HCV And Reprocessing Of Dialyzers.</i> | <i>56</i> |
| <i>PATIENTS AND METHODS</i> | <i>58</i> |
| <i>RESULTS</i> | <i>67</i> |
| <i>DISCUSSION</i> | <i>76</i> |
| <i>CONCLUSION AND RECOMMENDATIONS</i> | <i>80</i> |
| <i>SUMMARY</i> | <i>81</i> |
| <i>REFERENCES</i> | <i>83</i> |
| <i>ARABIC SUMMARY</i> | |

LIST OF TABLES

| <i>Tables of review</i> | | |
|--------------------------|---|----|
| <i>Table 1:</i> | <i>Proposed theories to explain low aminotransferases in patients with renal failure.</i> | 33 |
| <i>Table 2:</i> | <i>Comparison between formaldehyde and peracetic acid.</i> | 47 |
| <i>Table 3:</i> | <i>OSHA environmental exposure limits.</i> | 56 |
| <i>Tables of results</i> | | |
| <i>Table 1:</i> | <i>Prevalence of HCV RNA positivity among anti-HCV positive HD patients.</i> | 69 |
| <i>Table 2:</i> | <i>Mean values of age, duration on HD and serum creatinine for HC viremic group.</i> | 69 |
| <i>Table 3:</i> | <i>Mean values of serum ALT levels and mean number of blood transfusion among HC viremic group.</i> | 69 |
| <i>Table 4:</i> | <i>Descriptive statistics of various parameters among HC viremic females.</i> | 70 |
| <i>Table 5:</i> | <i>Descriptive statistics of various parameters among HC viremic males.</i> | 70 |
| <i>Table 6:</i> | <i>Correlation between different data among HC viremic HD patients.</i> | 71 |
| <i>Table 7:</i> | <i>Comparison between results of PCR for HCV RNA of the fifteen HD patients and the results of PCR test for HCV RNA in their dialyzers' wash after reprocessing.</i> | 72 |
| <i>Table 8:</i> | <i>Comparison between results of PCR test for HCV RNA in the blood samples obtained from the patients during the HD sessions (after passing through the dialyzers) and PCR results for HCV RNA in dialyzers' wash of these patients after reprocessing.</i> | 72 |
| <i>Table 9:</i> | <i>Crude data of patients</i> | 73 |

LIST OF FIGURES

| | | |
|------------------|---|-----------|
| <i>Figure 1:</i> | <i>Comparison between results of PCR test for HCV RNA of the fifteen HD patients and the PCR test for HCV RNA in their dialyzers' wash after reprocessing.</i> | <i>74</i> |
| <i>Figure 2:</i> | <i>Comparison between results of PCR test for HCV RNA in the blood samples obtained during the HD sessions and PCR results for HCV RNA in dialyzers' wash after reprocessing.</i> | <i>75</i> |

LIST OF ABBREVIATIONS

| | |
|---------------------|-----------------------------------|
| <i>ALT</i> | Alanine aminotransferase |
| <i>ELISA</i> | Enzyme-linked immunosorbant assay |
| <i>ETO</i> | Ethylene oxide |
| <i>KUF</i> | Ultrafiltration coefficient |
| <i>n-PCR</i> | Nested-polymerase chain reaction. |
| <i>PCR</i> | Polymerase chain reaction. |
| <i>RIBA</i> | Recombinant immunoblot assay. |

Errata

| <u>Page</u> | <u>Row</u> | <u>Mistake</u> | <u>Correct</u> |
|-------------|--------------|---|---|
| 58 | /13 | Chronic glomerulonephritis in 3 patients | Chronicglomerulone phritis in 2 patients and SLE in one patient |
| 70 | Table (4) | Number = 15 | Number = 9 |
| 70 | Table (5) | Number = 15 | Number = 6 |
| 111 | /1 | Cuypers H.T Reesink H.W | Cuypers H.T Reesink H.W |

