



**Isolation and characterization of glucose-6-phosphate dehydrogenase enzyme from liver of some mammals for laboratory diagnosis**

**Thesis Submitted**

**By**

***Mohamed Mosaad Abdel-Monseif Soliman***

M. Sc. in Biochemistry, Biochemistry Department  
Faculty of Science, Ain Shams University, 2010  
Assistant Researcher, Molecular Biology Department  
Genetic Engineering and Biotechnology Division  
National Research Centre

**Ph. D. Degree in Biochemistry**

**Supervised by**

***Prof. Dr.***

***Ahmed Mohamed Hassan Salem***

Prof. of Biochemistry  
Biochemistry Department  
Faculty of Science  
Ain Shams University

***Prof. Dr.***

***Mahmoud Abdel-Aziz Ibrahim***

Prof. of Biochemistry  
Molecular Biology Department  
Head of Genetic Engineering and  
Biotechnology Division  
National Research Centre

***Dr.***

***Mohamed Abdel-Hady Mahmoud Ghazy***

Assistant Professor of Biochemistry  
Biochemistry Department  
Faculty of Science  
Ain Shams University

**Biochemistry Department  
Faculty of Science  
Ain Shams University  
(2014)**

**Isolation and characterization of glucose-6-phosphate dehydrogenase enzyme from liver of some mammals for laboratory diagnosis**

**Board of Scientific Supervision**

**Prof. Dr. Ahmed Mohamed Hassan Salem**  
**Emeritus Professor of Biochemistry**  
**Biochemistry Department**  
**Faculty of Science**  
**Ain Shams University**  
**Cairo - Egypt**

**Prof. Dr. Mahmoud Abdel-Aziz Ibrahim**  
**Professor of Biochemistry**  
**Molecular Biology Department**  
**Head of Genetic Engineering and Biotechnology Division**  
**National Research Centre**  
**Cairo - Egypt**

**Dr. Mohamed Abdel-Hady Mahmoud Ghazy**  
**Assistant Professor of Biochemistry**  
**Biochemistry Department**  
**Faculty of Science**  
**Ain Shams University**  
**Cairo - Egypt**



## **ACKNOWLEDGEMENT**

*I would like to express my deep appreciations to **Prof. Dr. Ahmed Mohamed Hassan Salam**, emeritus Professor of Biochemistry, Department of Biochemistry, Faculty of Science, Ain Shams University for his valuable supervision, guidance and kind encouragement.*

*I would like to express my sincere and great appreciation to **Prof. Dr. Mahmoud Abdel-Aziz Ibrahim**, Professor of Biochemistry, Molecular Biology Department, National Research Center, for suggesting the research point, his close supervision and giving the full facilities for the proper research work,*

*My deep appreciations are due to **Prof. Dr. Abdel-Hady Mahmoud Ghazy**, emeritus Professor of Biochemistry, Molecular Biology Department, National Research Center, for his generous help, kindness and encouragement throughout the course of this work,*

*I would like to express my sincere gratitude to **Dr. Mohamed Abdel-Hady Mahmoud Ghazy**, Assistant Professor of Biochemistry, Biochemistry Department, Faculty of Science, Ain Shams University for his valuable supervision and kind support.*

*The financial support offered by National Research Centre and facilities offered by Molecular Biology Department, that enabled me to carry out this work, are greatly appreciated.*

*Deepest gratitude is indebted to my colleagues in Molecular Biology Department, National Research Centre.*

*Deepest gratitude is indebted to my family; my father, my mother, my brothers, my sisters and my wife for their help and continuous encouragement. I wouldn't be here if I didn't have them beside me.*

---

---

## TABLE OF CONTENTS

	Page
List of Figures-----	VII
List of Tables-----	XIV
List of Abbreviations-----	XV
<b>Abstract</b> -----	XVII
<b>Aim of work</b> -----	1
<b>Introduction</b> -----	3
I- Importance of glucose-6-phosphate dehydrogenase -----	6
II- Distribution of glucose-6-phosphate dehydrogenase -----	7
III- Molecular properties of glucose-6-phosphate dehydrogenase -----	8
1- Genetics of glucose-6-phosphate dehydrogenase-----	8
2- Structure and function of glucose-6-phosphate dehydrogenase -----	9
3- Molecular weight and subunits-----	10
4- Substrate and coenzyme specificity-----	11
5- Optimum pH-----	12
6- Optimum temperature -----	12
7- Inhibition -----	13
(A) Product inhibition-----	13
(B) Inhibition by various inhibitors-----	13
8- Regulation of glucose-6-phosphate dehydrogenase-----	14
(A) Regulation of G6PD by nutritional and hormonal factors -----	14
(B) Regulation of glucose-6-phosphate dehydrogenase by oxidative stress-----	15
IV- Purification of glucose-6-phosphate dehydrogenase -----	16
V- The hexokinase/glucose-6-phosphate dehydrogenase (HK/G6PD) diagnostic kit -----	17
<b>Materials and Methods</b> -----	20
A- MATERIALS-----	20
1- Liver material-----	20
2- Chemicals-----	20
3- Buffers-----	21

---

---

- II -

---

---

B- METHODS-----	21
I- Enzyme assay-----	21
Assay of glucose-6-phosphate dehydrogenase activity-	21
II- Protein determination-----	21
1- Principle-----	21
2- Preparation of the protein assay reagent-----	22
3- Procedure-----	22
III- Preparation of columns for chromatography-----	24
1- Preparation of DEAE-cellulose column-----	24
2- Preparation of sephacryl S-300 column-----	25
3- Preparation of 2', 5' ADP Sepharose 4B affinity column -----	25
IV- Purification of the liver glucose-6-phosphate dehydrogenase -----	26
1- Preparation of crude extract -----	26
2- Preparation of Ammonium sulfate precipitation ---	27
3- DEAE-cellulose column chromatography-----	27
4- Sephacryl S-300 column chromatography-----	28
5- 2', 5' ADP Sepharose 4B affinity column Chromatography-----	28
V- Molecular weight determination by gel filtration-----	28
VI- Electrophoretic analyses-----	29
A- Native polyacrylamide gel electrophoresis (PAGE) ----	29
1- Principle-----	29
2- Reagents-----	31
3- Procedure-----	32
B- Sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE) -----	33
1 – Principle-----	33
2 – Reagents-----	33
3 – Procedure-----	35
4 – Molecular weight determination by SDS-PAGE----	37
C- Isoelectric focusing (IEF) -----	39
1- Principle-----	39
2- Reagents-----	39

---

---

- III -

---

3- Procedure-----	41
4- Determination of isoelectric point <i>pI</i> value-----	43
D- Staining of the glucose-6-phosphate dehydrogenase enzyme activity-----	43
VII- Statistical analysis-----	45
VIII- Preparation of glucose hexokinase diagnostic kit by using the purified G6PD -----	45
1- Reagents Preparation -----	45
2- System parameters -----	47
3- Assay Procedure -----	48
4- Calculation-----	48
5- Reference Range -----	48
<b>Results</b> -----	49
Comparison of the liver glucose-6-phosphate dehydrogenase activity: -----	49
1- Based on the specific activity level: -----	49
2- Based on the anion exchange chromatographic pattern: --	51
3- Based on electrophoretic pattern: -----	55
<b>I- Buffalo liver glucose-6-phosphate dehydrogenases: -----</b>	<b>58</b>
-----	58
A- Purification of glucose-6-phosphate dehydrogenase from buffalo liver: -----	58
1- Preparation of crude extract: -----	58
2- Ammonium sulfate precipitation: -----	58
3- Chromatography on the DEAE-cellulose column: --	60
4- Chromatography on sephacryl S-300 column: -----	62
5- Affinity chromatography on 2', 5' ADP Sepharose 4B column: -----	65
B- Characterization of the buffalo liver glucose-6-phosphate dehydrogenases BLG6PD1 and BLG6PD2:-----	68
1- Molecular weight determination of the buffalo liver glucose-6-phosphate dehydrogenase by gel filtration: - -----	68
2- Electrophoretic analyses of buffalo liver glucose-6-phosphate dehydrogenase: -----	68
A- Homogeneity by native 7 % PAGE: -----	68

---

---

---

B- Subunit molecular weight by SDS-PAGE: -----	69
C- Isoelectric point determination: -----	69
3- Determination of optimum pH: -----	77
4- Effect of divalent cations: -----	77
5- Michaelis-Menten constant ( <i>K<sub>m</sub></i> ) value: -----	80
6- Effect of various inhibitors: -----	82
7- Kinetics of BLG6PD1 and BLG6PD2 inhibition by NADPH: -----	87
A- Effect of NADPH concentration on BLG6PD1 and BLG6PD2 activity: -----	88
B- Number of binding site: -----	88
C- Type of inhibition: -----	88
D- Determination of the inhibition constant ( <i>K<sub>i</sub></i> ): -----	92
<b>II- Camel liver glucose-6-phosphate dehydrogenase: ----</b>	<b>92</b>
A- Purification of glucose-6-phosphate dehydrogenase from camel liver: -----	92
1- Preparation of crude extract: -----	92
2- Ammonium sulfate precipitation: -----	94
3- Chromatography on the DEAE-cellulose column: -	96
4- Chromatography on Sephacryl S-300 column: -----	96
5- Affinity chromatography on 2', 5' ADP Sepharose 4B column: -----	99
B- Characterization of the camel liver glucose-6-phosphate dehydrogenase CLG6PD:-----	101
1- Molecular weight determination of the camel liver glucose-6-phosphate dehydrogenase by gel filtration: - -----	101
2- Electrophoretic analyses of camel liver glucose-6-phosphate dehydrogenase: -----	101
A- Homogeneity by native 7 % PAGE: -----	101
B- Subunit molecular weight by SDS-PAGE: -----	104
C- Isoelectric point determination: -----	104
3- Determination of optimum pH: -----	104
4- Effect of divalent cations: -----	108
5- Michaelis-Menten constant ( <i>K<sub>m</sub></i> ) value: -----	108

---

---

---

6- Effect of various inhibitors: -----	111
7- Kinetics of CLG6PD inhibition by NADPH: -----	116
A- Effect of NADPH concentration on CLG6PD activity: -----	116
B- Number of binding site: -----	116
C- Type of inhibition: -----	119
D- Determination of the inhibition constant ( $K_i$ ): -----	119
<b>III- Sheep liver glucose-6-phosphate dehydrogenases: --</b>	<b>122</b>
A- Purification of glucose-6-phosphate dehydrogenase from sheep liver: -----	122
1- Preparation of crude extract: -----	122
2- Ammonium sulfate precipitation: -----	122
3- Chromatography on the DEAE-cellulose column: --	124
4- Chromatography on sephacryl S-300 column: -----	124
5- Affinity chromatography on 2', 5' ADP Sepharose 4B column: -----	126
B- Characterization of the sheep liver glucose-6-phosphate dehydrogenase SLG6PD1 and SLG6PD2:-----	131
1- Molecular weight determination of the sheep liver glucose-6-phosphate dehydrogenase by gel filtration: ----	131
2- Electrophoretic analyses of sheep liver glucose-6-phosphate dehydrogenase: -----	131
A- Homogeneity by native 7 % PAGE: -----	131
B- Subunit molecular weight by SDS-PAGE: -----	136
C- Isoelectric point determination: -----	136
3- Determination of optimum pH: -----	140
4- Effect of divalent cations: -----	140
5- Michaelis-Menten constant ( $K_m$ ) value: -----	143
6- Effect of various inhibitors: -----	145
7- Kinetics of SLG6PD1 and SLG6PD2 inhibition by NADPH: -----	150
A- Effect of NADPH concentration on SLG6PD1 and SLG6PD2 activity: -----	150
B- Number of binding site: -----	152
C- Type of inhibition: -----	152

---



---

---

D- Determination of the inhibition constant ( $K_i$ ): -----	155
<b>IV- Comparison of the constructed kit with the commercially available kits -----</b>	<b>155</b>
<b>Discussion-----</b>	<b>160</b>
<b>References-----</b>	<b>178</b>
<b>Summary-----</b>	<b>192</b>
<b>Arabic Summary</b>	
<b>Arabic Abstract</b>	

---

---

---

---

**LIST OF FIGURES**

<b>Figure No.</b>	<b>Page</b>
(1): Standard curve of the protein concentration using bovine serum albumin as a standard protein -----	23
(2): A Calibration curve for the molecular weight determination of glucose-6-phosphate dehydrogenase by gel filtration on the Sephacryl S-300 column-----	30
(3): A Calibration curve for the molecular weight determination by 12 % SDS-PAGE-----	38
(4): A calibration curve for isoelectric point ( <i>pI</i> ) value determination by isoelectric focusing (IEF) -----	44
(5): Standard curve of the prepared glucose hexokinase diagnostic kit using glucose as a standard (100 mg/dl) -----	46
(6): A typical elution profile for the chromatography of the buffalo liver crude extract on DEAE-cellulose column-----	52
(7): A typical elution profile for the chromatography of the camel liver crude extract on DEAE-cellulose column -----	53
(8): A typical elution profile for the chromatography of the sheep liver crude extract on DEAE-cellulose column-----	54
(9): Electrophoretic analysis of liver crude extract of the isoenzyme pattern of three mammals on 7 % native polyacrylamide gel -----	56
(10): Electrophoretic analysis of liver crude extract of the protein pattern of three mammals on 7 % native polyacrylamide gel -----	57
(11): A typical elution profile for the chromatography of the buffalo liver ammonium sulfate fraction on DEAE-cellulose column -----	61
(12): A typical elution profile for the chromatography of the concentrated pooled DEAE-cellulose fractions containing the buffalo liver glucose-6-phosphate dehydrogenase activity (BLG6PD1) on Sephacryl S-300 column-----	63

---

---

- VIII -

- 
- 
- (13): A typical elution profile for the chromatography of the concentrated pooled DEAE-cellulose fractions containing the buffalo liver glucose-6-phosphate dehydrogenase activity (BLG6PD2) on Sephacryl S-300 column -----64
- (14): A typical elution profile for the chromatography of the concentrated pooled Sephacryl S-300 column fractions containing the buffalo liver glucose-6-phosphate dehydrogenase activity (BLG6PD1) on 2', 5' ADP sepharose 4B column ----- 66
- (15): A typical elution profile for the chromatography of the concentrated pooled Sephacryl S-300 column fractions containing the buffalo liver glucose-6-phosphate dehydrogenase activity (BLG6PD2) on 2', 5' ADP sepharose 4B column -----67
- (16): Electrophoretic analysis of buffalo liver glucose-6-phosphate dehydrogenase (BLG6PD1) of the different purification steps on 7 % native polyacrylamide gel, protein pattern ----- 70
- (17): Electrophoretic analysis of buffalo liver glucose-6-phosphate dehydrogenase (BLG6PD1) of the different purification steps on 7 % native polyacrylamide gel, isoenzyme pattern -----71
- (18): Electrophoretic analysis of buffalo liver glucose-6-phosphate dehydrogenase (BLG6PD2) of the different purification steps on 7 % native polyacrylamide gel, protein pattern:----- 72
- (19): Electrophoretic analysis of buffalo liver glucose-6-phosphate dehydrogenase (BLG6PD2) of the different purification steps on 7 % native polyacrylamide gel, isoenzyme pattern -----73
- (20): Subunit molecular weight determination by electrophoretic analysis of buffalo liver glucose-6-phosphate dehydrogenase isoenzyme (BLG6PD1) on 12 % SDS-polyacrylamide gel ----- 74
- (21): Subunit molecular weight determination by electrophoretic analysis of buffalo liver glucose-6-
- 
-

- IX -

---

---

phosphate dehydrogenase isoenzyme (BLG6PD2) on 12 % SDS-polyacrylamide gel -----	75
(22): Isoelectrofocusing; (1) Isoelectric point ( <i>pI</i> ) marker proteins, (2) the purified buffalo liver glucose-6-phosphate dehydrogenase BLG6PD1 and (3) the purified buffalo liver glucose-6-phosphate dehydrogenase BLG6PD2-----	76
(23): Effect of pH on the purified buffalo liver glucose-6-phosphate dehydrogenase (BLG6PD1 and BLG6PD2) using 0.02 M Tris / HCl buffer of various pH values -	93
(24): Effect of divalent cations on BLG6PD1 and BLG6PD2--- -----	79.
(25): Effect of the substrate $\beta$ -Nicotinamide adenine dinucleotide phosphate ( $\text{NADP}^+$ ) concentration in mM on the reaction velocity of the purified buffalo liver BLG6PD1 and BLG6PD2-----	81
(26): Lineweaver-Burk plot relating the reciprocal of the reaction velocity of the purified buffalo liver BLG6PD1 and BLG6PD2 to $\beta$ -Nicotinamide adenine dinucleotide phosphate ( $\text{NADP}^+$ ) concentration in mM -----	83
(27): Effect of inhibitors on BLG6PD1 and BLG6PD2-----	84
(28): Inhibition of the purified buffalo liver BLG6PD1 and BLG6PD2 by varying concentrations of NADPH ----	89
(29): Hill plot for inhibition of the purified liver BLG6PD1 and BLG6PD2 by varying concentrations of NADPH -----	90
(30): Lineweaver-Burk plots showing the type of inhibition of the purified buffalo liver BLG6PD1 and BLG6PD2 by NADPH. -----	91
(31): Determination of the inhibition constant ( <i>K<sub>i</sub></i> ) value for the inhibition of the purified buffalo liver BLG6PD1 and BLG6PD2 by NADPH -----	93
(32): A typical elution profile for the chromatography of the camel liver ammonium sulfate fraction on DEAE-cellulose column -----	97

---

---

- 
- 
- (33): A typical elution profile for the chromatography of the concentrated pooled DEAE-cellulose fractions containing the camel liver glucose-6-phosphate dehydrogenase activity (CLG6PD) on Sephacryl S-300 column----- 98
- (34): A typical elution profile for the chromatography of the concentrated pooled Sephacryl S-300 column fractions containing the camel liver glucose-6-phosphate dehydrogenase activity (CLG6PD) on 2', 5' ADP sepharose 4B column ----- 100
- (35): Electrophoretic analysis of camel liver glucose-6-phosphate dehydrogenase (CLG6PD) of the different purification steps on 7 % native polyacrylamide gel, protein pattern ----- 102
- (36): Electrophoretic analysis of camel liver glucose-6-phosphate dehydrogenase (CLG6PD) of the different purification steps on 7 % native polyacrylamide gel, isoenzyme pattern ----- 103
- (37): Subunit molecular weight determination by electrophoretic analysis of camel liver glucose-6-phosphate dehydrogenase CLG6PD on 12 % SDS-polyacryl-amide gel ----- 105
- (38): Isoelectrofocusing; (1) Isoelectric point ( $pI$ ) marker proteins, (2) the purified camel liver glucose-6-phosphate dehydrogenase CLG6PD ----- 106
- (39): Effect of pH on the purified camel liver glucose-6-phosphate dehydrogenase CLG6PD using 0.02 M Tris / HCl buffer of various pH values ----- 107
- (40): Effect of divalent cations on the purified camel liver glucose-6-phosphate dehydrogenase CLG6PD ----- 109
- (41): Effect of the substrate  $\beta$ -Nicotinamide adenine dinucleotide phosphate ( $NADP^+$ ) concentration in mM on the reaction velocity of the purified camel liver CLG6PD ----- 110
- (42): Lineweaver-Burk plot relating the reciprocal of the reaction velocity of the purified camel liver CLG6PD to
- 
-

---

---

β-Nicotinamide adenine dinucleotide phosphate (NADP <sup>+</sup> ) concentration in mM -----	112
(43): Effect of inhibitors on the purified camel liver glucose-6- phosphate dehydrogenase CLG6PD -----	113
(44): Inhibition of the purified camel liver CLG6PD by varying concentrations of NADPH -----	117
(45): Hill plot for inhibition of the purified camel liver CLG6PD by varying concentrations of NADPH -----	118
(46): Lineweaver-Burk plots showing the type of inhibition of the purified camel liver CLG6PD by NADPH. The activity of CLG6PD was measured with varying concentrations of the substrate β-Nicotinamide adenine dinucleotide phosphate (NADP <sup>+</sup> ) in absence and presence of three various concentrations of NADPH -----	120
(47): Determination of the inhibition constant ( <i>K<sub>i</sub></i> ) value for the inhibition of the purified camel liver CLG6PD by NADPH-----	121
(48): A typical elution profile for the chromatography of the sheep liver ammonium sulfate fraction on DEAE- cellulose column -----	125
(49): A typical elution profile for the chromatography of the concentrated pooled DEAE-cellulose fractions containing the sheep liver glucose-6-phosphate dehydrogenase activity (SLG6PD1 and SLG6PD1) on Sephacryl S-300 column -----	127
(50): A typical elution profile for the chromatography of the concentrated pooled Sephacryl S-300 column fractions containing the sheep liver glucose-6-phosphate dehydrogenase activity (SLG6PD1) on 2', 5' ADP sepharose 4B column -----	129
(51): A typical elution profile for the chromatography of the concentrated pooled Sephacryl S-300 column fractions containing the sheep liver glucose-6-phosphate dehydrogenase activity (SLG6PD2) on 2', 5' ADP sepharose 4B column -----	130

---

---

- 
- 
- (52): Electrophoretic analysis of sheep liver glucose-6-phosphate dehydrogenase (SLG6PD1) of the different purification steps on 7 % native polyacrylamide gel, protein pattern -----132
- (53): Electrophoretic analysis of sheep liver glucose-6-phosphate dehydrogenase (SLG6PD1) of the different purification steps on 7 % native polyacrylamide gel, isoenzyme pattern ----- 133
- (54): Electrophoretic analysis of sheep liver glucose-6-phosphate dehydrogenase (SLG6PD2) of the different purification steps on 7 % native polyacrylamide gel, protein pattern ----- 134
- (55): Electrophoretic analysis of sheep liver glucose-6-phosphate dehydrogenase (SLG6PD2) of the different purification steps on 7 % native polyacrylamide gel, isoenzyme pattern ----- 135
- (56): Subunit molecular weight determination by electrophoretic analysis of sheep liver glucose-6-phosphate dehydrogenase isoenzyme (SLG6PD1) on 12 % SDS-polyacrylamide gel ----- 137
- (57): Subunit molecular weight determination by electrophoretic analysis of sheep liver glucose-6-phosphate dehydrogenase isoenzyme (SLG6PD2) on 12 % SDS-polyacrylamide gel: ----- 138
- (58): Isoelectrofocusing; (1) Isoelectric point ( $pI$ ) marker proteins, (2) the purified sheep liver glucose-6-phosphate dehydrogenase SLG6PD1-----139
- (59): Effect of pH on the purified sheep liver glucose-6-phosphate dehydrogenase (SLG6PD1 and SLG6PD2) using 0.02 M Tris / HCl buffer of various pH values - 141
- (60): Effect of divalent cations on SLG6PD1 and SLG6PD2---  
----- 142
- (61): Effect of the substrate  $\beta$ -Nicotinamide adenine dinucleotide phosphate ( $NADP^+$ ) concentration in mM on the reaction velocity of the purified sheep liver SLG6PD1 and SLG6PD2----- 144
- 
-