

سامية محمد مصطفى



شبكة المعلومات الجامعية

بِسْمِ اللَّهِ الرَّحْمَنِ الرَّحِيمِ




سامية محمد مصطفى



شبكة المعلومات الجامعية



شبكة المعلومات الجامعية التوثيق الالكتروني والميكروفيلم



سامية محمد مصطفى



شبكة المعلومات الجامعية

جامعة عين شمس

التوثيق الإلكتروني والميكروفيلم


قسم

نقسم بالله العظيم أن المادة التي تم توثيقها وتسجيلها
علي هذه الأقراص المدمجة قد أعدت دون أية تغييرات



يجب أن

تحفظ هذه الأقراص المدمجة بعيدا عن الغبار




سامية محمد مصطفى



شبكة المعلومات الجامعية



بعض الوثائق الأصلية تالفة



سامية محمد مصطفى




شبكة المعلومات الجامعية



بالرسالة صفحات

لم ترد بالأصل



**EVALUATION OF RESULTS OF OPERATIVE
TREATMENT OF FRACTURE NECK FEMUR
IN YOUNG ADULTS**

THIS IS

*SUBMITTED IN PARTIAL FULFILMENT FOR THE
DEGREE OF M.D. ORTHOPAEDIC SURGERY*

BY

MOHAMAD AHMAD MAHMAUD A. ALAM EL-DIN
*ASSISTANT LECTURER
DEPARTMENT OF ORTHOPAEDIC SURGERY
SOHAG FACULTY OF MEDICINE*

UNDER THE SUPERVISION OF

PROF. DR. ABDEL-KHALIK H. IBRAHIM
*PROFESSOR OF ORTHOPAEDIC SURGERY
FACULTY OF MEDICINE
ASSIUT UNIVERSITY*

PROF. DR. ESSAM k. EL-SHERIF
*PROFESSOR OF ORTHOPAEDIC SURGERY
FACULTY OF MEDICINE
ASSIUT UNIVERSITY*


DR. KAMAL a. M. EL-GAAFARY
*ASSISTANT PROFESSOR OF ORTHOPAEDIC SURGERY
FACULTY OF MEDICINE
ASSIUT UNIVERSITY*

**FACULTY OF MEDICINE
ASSIUT UNIVERSITY**


1995

B

17.70



بِسْمِ اللَّهِ الرَّحْمَنِ الرَّحِيمِ



DEDICATION

To : *The memory of my parents*

To : *My uncles who have added a very
special dimension to my life.*

To: *My dear wife.*



ACKNOWLEDGEMENT

ACKNOWLEDGMENT

All gratitude and thanks are due to Allah, the Beneficial, the Merciful whose great help I always sought not only in this work but in every affair throughout my life.

I would like to express my great appreciation and much gratitude to Prof Dr. ABDEL KHALIK H. IBRAHIM, Professor of Orthopaedic Surgery and Traumatology, Faculty of Medicine, Assiut University, for his excellent supervision, generous discussions, clarifying suggestions and constant encouragement. It was he who suggested the topic of this thesis to me.

I offer my warm thanks to Prof. Dr. ESSAM K. EL-SHERIF, Professor of Orthopaedic Surgery, for constant support and continuous friendly encouragement. I am greatly indebted to him for his stimulating and ever enthusiastic support during a great number of years.

I would like also to thank Dr. KAMAL EL-GAAFARY, Assistant Professor of Orthopaedic Surgery, for his kind assistance and help throughout the preparation of this work.

I wish to express my deep gratitude to Prof. Dr. GALAL Z. SAID, Head of the Department of Orthopaedics and Traumatology, Faculty of Medicine, Assiut University, my teacher in orthopaedic surgery, for his advice and support during this study.

All thanks to all the staff members of the Orthopaedic Surgery and Traumatology department and to my colleagues for their co-operation and help.

Mohammed Alam El-Din


1995



CONTENTS

CONTENTS

	<i>Page</i>
INTRODUCTION AND AIM OF THE WORK	1
REVIEW OF THE LITERATURE	3
* <i>General Consideration</i>	3
* <i>Historical Review</i>	36
* <i>Mechanism of Femoral Neck Fractures</i>	42
* <i>Classification of Femoral Neck Fractures</i>	45
* <i>Healing of Femoral neck Fractures</i>	74
* <i>Osteoporosis and Fracture of Femoral Neck</i>	104
MATERIALS AND METHODS	108
RESULTS	134
DISCUSSION	178
SUMMARY AND CONCLUSIONS	207
REFERENCES	212
ARABIC SUMMARY	



INTRODUCTION
AND
AIM OF THE WORK

INTRODUCTION AND AIM OF THE WORK

The history of treatment of femoral neck fractures in young adults is one of a persistent search for adequate reduction and fixation in an effort to minimize the incidence of non-union and avascular necrosis⁽⁵⁴⁾.

This fracture continues to be considered "unsolved" until the incidence of these two complications can be remarkably diminished or abolished⁽⁶⁰⁾. According to Wilson⁽⁴⁰⁾, united femoral neck fracture with an intact femoral head is better than any prosthesis, and if there is any reasonable chance for accurate reduction and good internal fixation, nailing should be the operation of choice. The materials of this work included 47 cases of recent displaced femoral neck fractures admitted to the department of Orthopaedic Surgery and Traumatology, Assiut University. Patients had been observed for an average of 31 months following surgery. These cases had been assessed clinically by the grading method described by D.Aubigne⁽²²⁾. In this method three items had been taken into consideration namely pain, mobility of the hip and the ability to walk. Each of the three items was given a degree from 0 to 6. Also, radiological assessment had been done by plain x-rays taken every three months to detect the state of union of the fracture and the early signs denoting the non viability of the femoral head.

The aim of this work is to study the results of treatment of recent displaced fractures of the femoral neck in adults treated by 3 cancellous lag screws fixation and to compare our results with those of other authors.