



Value of 18-FDG PET/CT in assessment of solitary pulmonary nodule

Essay

Submitted for partial fulfillment of Master Degree in
Radiodiagnosis

By

Mohammed Ashraf Mohammed Keshk

M.B., B.Ch.

Ain Shams University

Under The Supervision Of

Prof. Dr. Mervat Mohammed Elgohary

Professor of Radio-diagnosis

Faculty of Medicine - Ain Shams University

Dr. Mary Yaftah Tadros

Lecturer of Radio-diagnosis

Faculty of Medicine - Ain Shams University

Faculty of Medicine

Ain-shams University

2015



First and foremost, I thank **Allah** for endowing me with health, patience, and knowledge to complete this work, till it reached its end.

I acknowledge, with deep gratitude and appreciation, encouragement, valuable time and guidance that was given to me by **Prof. Dr. Mervat Mohammed Elgohary**, Professor of Radiodiagnosis, Ain Shams University, who I am deeply thankful for her valuable supervision.

Thereafter, I am deeply grateful to **Dr. Mary Yafrah Tadros**, Lecturer of Radiodiagnosis, Ain Shams University for her extensive guidance, continuous support, and personal involvement in all phases of this work.

Thanks are due to my senior colleagues at the hospital for their Friendship and great help in all times.

Finally, I would like to express my deepest gratitude to my mother, father, my wife for their emotional and moral support throughout this work and also for their love, patience, encouragement and prayers.



LIST OF CONTENTS

Introduction and Aim of the work.

Chapter I: - Lung anatomy.

Chapter II: - Pathology of Solitary pulmonary nodule.

Chapter III: - Technology & Principles of PET/CT.

Chapter IV: - PET/CT findings in solitary pulmonary nodule.

Summary & Conclusion.

References.

Arabic summary.

List of Figures

Figure	Title	Page
1	The position of lungs within rib cage	4
2	Lung surfaces	5
3	Pleural cavities	6
4	Diagram illustrating lung fissures	7
5	Bronchopulmonary segments A. Right lung, B. Left lung	8
6	Roots and hila of lungs	9
7	Diagram illustrating lung vasculature	10
8	Lymph nodes related to trachea and main bronchi	12
9	Bronchopulmonary segments	14
10	Bronchial veins	15
11	Normal CT appearance of chest, Both Lung and mediastinal window	16
12	Normal major fissures. High-resolution CT scans show the major fissures as lines, arrows	18
13	CT Chest image using lung window at level of apex of lung	20
14	CT Chest image using lung window at level of apex of lung, lower level	21

List of Figures

15	CT Chest image using lung window at level of the arch of aorta	22
16	CT Chest image using lung window at level of Carina	23
17	CT Chest image using lung window at the level of the left atrium	24
18	Normal distribution of FDG. Coronal CT (a), PET (b), and PET-CT fusion (c) images demonstrate the physiologic accumulation of FDG in the cerebral-cerebellar cortex at the base of the skull and in the myocardium, liver, kidneys, renal pelvis, bone marrow, and urinary bladder	25
19	MIP FDG PET image (a) and transverse FDG PET (b), CT (c), and fused PET/CT (d) images show mildly increased uptake in the arytenoid muscles, arrows	26
20	A. Squamous cell carcinoma usually begins as a central (hilar) mass and growing into the peripheral parenchyma. B. Histology showing keratinization and pearls	32
21	Adenocarcinoma in situ, mucinous subtype, with characteristic growth along preexisting alveolar septa, without invasion	33
22	Small cell carcinoma with small deeply basophilic cells and areas of necrosis	34
23	Bronchial carcinoid. A. Carcinoid growing as a spherical, pale mass (arrow) protruding into the lumen of the bronchus. B. Histologic appearance demonstrating small, rounded, uniform nuclei and moderate cytoplasm	35
24	Histological appearance of hamartoma containing benign	38

List of Figures

	cartilage, adipocytes and respiratory epithelium	
25	Hemorrhagic, roughly wedge-shaped pulmonary infarct	41
26	Photomicrograph in Bronchogenic cyst in a 49-year-old man with a 2-week history of fever and cough , showing a portion of the cyst wall with respiratory epithelium (arrow) that overlies fibrous tissue and cartilage	42
27	Cryptogenic organizing pneumonia. Some alveolar spaces are filled with fibroblasts (Masson bodies)	43
28	A. Vasculitis of a small artery with adjacent granulomatous inflammation including giant cells (arrows). B. Lung from a patient with Wegener granulomatosis, demonstrating large nodular cavitating lesions	44
29	The principles of PET imaging	47
30	Current commercial PET-CT scanners : (a) GE Discovery PET/CT 600 for the BGO version (GE Healthcare); (b) GE Discovery PET/CT 690 for the Lutetium version (GE Healthcare); (c) Philips Medical Systems ; (d) Siemens Medical Solutions; (e) Hitachi Medical Systems ; (f) Toshiba Medical Corporation	49
31	A schematic of a current PET-CT scanner design. The design incorporates a multidetector spiral CT scanner and a lutetium oxy-orthosilicate (LSO) PET scanner. The dimensions of the gantry are 228 cm wide, 200 cm high, and 168 cm deep	50
32	Illustration showing data acquisition with septa (left) and without septa (right). In 3D mode (without septa) sensitivity is increased compared with 2D mode (with	52

List of Figures

	septa) However, in 3D, scatter is more significant	
33	Glucose and Flourodeoxyglucose structure	54
34	Display screen of the syngo software platform shows fused PET-CT images in the sagittal, coronal, and axial planes of a patient with recurrent esophageal carcinoma. Placing the crosshairs on the tumor automatically coregisters it in all three orthogonal planes. By dragging the vertical bar (arrow) along the horizontal line in the lower right quadrant of the screen, the PET and CT weighting of the image can be altered	58
35	Whole-body PET-CT imaging protocol. (a) First, a topogram is acquired , second, a CT scan and an emission scan follow, covering the predefined imaging range, then Fused PET-CT images are available (b), Example of artifacts like implants and artificial agents, patient motion, as well as patient setup and positioning	60
36	Respiratory motion artifact , Coronal PET, CT and PET/CT images (a–d) show severe displacement and apparent detachment of a large portion of the liver (arrows) from using tidal respiration breathing protocol during the CT acquisition	62
37	Extensive patient motion ,in a patient with right liver lesion , between the CT and PET portion of a combined FDG-PET-CT scan was noticeable in this case through a misalignment of the anatomic and functional information in the head and neck region (a, arrow). Distortion of the metabolic activity pattern on the attenuation-corrected PET (b, left and middle). The uncorrected emission images (b, right) suggest further patient motion during the scan	63

List of Figures

38	Artifacts because of metallic dental fillings (arrowhead) and hip prosthetics (arrows). PET scan shows apparent increased activity in oral cavity and right hip joint	64
39	Patient setup with arms down may lead to a variety of image distortions in whole-PET-CT exams	66
40	Corona radiata sign in a malignant lesion with spiculation at the margin	69
41	Pulmonary carcinoma. A. Axial CT image. B. & C. The calculated doubling time (DT) is 154 days, suggesting a neoplasm. D. The fused PET/CT image locates an area of metabolic hyperactivity with $SUV > 2.5$ (E)	72
42	A 60-year-old man with a histologically proven adenocarcinoma in the left central lung. The images include a MIP scan (A), a trans axial PET section (B), a corresponding axial non-enhanced CT section (C), and a corresponding axial PET/CT section (D). Diagnosis: Central lung tumor in the left lung, with a lymph node metastasis in the aortopulmonary window	74
43	Pulmonary carcinoma. A. Example of the segmentation of A 73 years old man with Squamous cell carcinoma of lung. A. Lung window B. 18-f FDG PET , C. Integrated 18-f FDG PET/CT show a mass around 30 mm (Arrow) with obstructive pneumonia (arrowhead) with increased FDG uptake in the right lower lobe	75
44	Bronchioalveolar carcinoma. (a) Axial chest PET/CT image shows a focal area of ground-glass attenuation in the right upper lobe (arrow) confirmed to be bronchioalveolar carcinoma at pathologic analysis. (b) Axial FDG PET scan shows only normal physiologic	76

List of Figures

	cardiac uptake	
45	Small cell lung cancer. Chest CT scan (a) and corresponding PET/CT image (b) show a 28 mm nodule in the left lung (top arrow) with intense radiotracer uptake on the PET/CT image with confluent ipsilateral prevascular (bottom arrow) and left paratracheal lymphadenopathy (N2), shows intense uptake as well	77
46	Metastatic nodes harboring cancer cells in right hilar & subcarinal regions observed at PET/CT in a 67-year-old man with lung adenocarcinoma. A: Axial CT scan B: Enhanced CT scan obtained at distal main bronchial level demonstrates enlarged lymph nodes in subcarinal and right hilar regions. C: PET (left) and PET/CT (right) images show high glucose uptake at them	78
47	Superior Vena Cava (SVC) Syndrome in Small Cell Carcinoma. Contrast-enhanced CT at the level of the tracheal carina reveals a right paratracheal mass (Red arrow) that obliterates the SVC. There is associated mediastinal venous collaterals (red arrowheads) and dilated internal mammary veins (blue arrowheads)	79
48	Single Pulmonary metastases in a young man with rectal cancer , showing nodule in the left lung in axial CT (A) , high uptake of glucose (B) proved to be metastatic nodule	80
49	PET-negative neuroendocrine tumor in a 59-year-old woman. Unenhanced CT (a) and PET/CT (b) images show a nodule in the middle lobe (arrow) with no FDG uptake. Results from transthoracic needle biopsy revealed a well-differentiated neuroendocrine tumor (carcinoid)	81
50	Mixed cellularity type Hodgkin disease with lung and mediastinal involvement. Axial fused PET/CT image	82

List of Figures

	shows foci of abnormal FDG uptake in a right lung lesion along the major fissure (black arrow), the thymus (arrowhead), and enlarged right hilar lymph nodes (white arrow)	
51	Coronal maximum intensity projection (MIP) (A) and three-dimensional volume-rendered CT images (B , C) revealed a Pulmonary Arteriovenous Malformation PAVM with a feeding artery, an aneurysmal part, and a draining vein	83
52	Hamartoma containing calcifications. (A) On CT, popcorn pattern of calcifications with mild low attenuation areas possibly representing fat. A confirmatory PET/CT study demonstrates low pattern of metabolic activity on coronal PET/CT (B)	84
53	Focal organizing pneumonia. (a) Unenhanced CT image shows a subsolid nodule (arrow) in the right lower lobe with predominantly ground-glass attenuation. (b) PET/CT image shows low-grade FDG uptake in the nodule (arrow). Results of transthoracic needle aspiration biopsy were negative and revealed focal organizing pneumonia	85
54	Lipoid pneumonia. An elderly female demonstrating a spiculated nodule , which on 1.5-mm soft tissue section low attenuation consistent with macroscopic fat is identified in the nodule (arrow) and mass-like areas in the right lower lobe , confirming the diagnosis of lipoid pneumonia	86
55	(a) Anterior and (b) lateral MIP images of the positron emission tomography (PET)/CT scan. A, nasal cavity; B, right parotid; C, anterior mediastinal mass; D, multiple cavitating pulmonary nodules; and E, prostate gland	87

List of Figures

	involvement by Wegener's granulomatosis	
56	A. an example of a central tumor with distal collapse, the margins of which cannot be clearly separated on CT alone. B. the fused PET/CT image and it clearly shows the central active tumor and the distal collapsed lung allowing a limited radiation port and thus sparing normal tissues	90
57	Lung adenocarcinoma in a 45-year-old patient. (A) Axial contrast (MDCT) image shows nodule (arrow) in the right upper lobe (B) Axial image with mediastinal window showed no adenopathy, only an isolated hilar node axis (arrow). (C) & (D) PET/CT images allows better characterization of lymph nodes so, this tumor was classified as cT1aN1M0	92
58	Sarcoidosis in a 32-year-old male. A. Axial PET-CT scans show bilateral paratracheal and hilar lymph node enlargement with high FDG uptake. B. Coronal section of PET shows high FDG uptake (arrowheads) which is highly suggestive of Sarcoidosis. A mediastinoscopic biopsy confirms Sarcoidosis	93
59	Bronchioalveolar carcinoma in a 75-year-old male. A. Contrast-enhanced CT shows a cavitary lesion in the left lower lobe (arrow). B. Transverse FDG-PET image of the transverse scan shows subtle uptake (arrow) in the left lower lobe with a maximum SUV of 1.7	94
60	Axial CT scan shows a 2.8 cm pulmonary nodule in the right upper lobe with spiculated margins and emphysema (A). 18 FDG PET axial (B) , coronal (D) and 18 FDG PET/CT fusion images (C) reveal increased 18 FDG uptake in the nodule. The lesion corresponded to a	95

List of Figures

	squamous-cell carcinoma	
61	Post-chemotherapy related decreased lymph node uptake in a 57 year old male lung cancer patient. A. Pre-chemotherapy FDG-PET fusion CT shows increased FDG uptake in the subcarinal lymph node. B & C After two months of chemotherapy, the residual lesion in the subcarinal area shows no abnormal FDG uptake	96
62	Contiguous pleural involvement of primary lung cancer. Trans axial CT image (a) demonstrates thickened pleura adjacent to a known malignant right lower lobe pulmonary nodule. On the PET-CT fusion image (b) the thickened pleura clearly demonstrate abnormal FDG tracer uptake most consistent with pleural extension	97
63	(A) PET scan shows moderately intense uptake in the nodule (solid arrow), indicating metabolic activity and incidental uptake in metabolically active brown fat (dotted arrows). (B) CT scan showing lung nodule (arrow). (C) Fused PET/CT image showing alignment of metabolic activity with nodule seen on CT	98
64	(A) Axial CT scan at lung windows through the lower trachea shows an irregular right upper lobe nodule. (B) Fused image from PET-CT shows marked increase in metabolic activity in the nodule	99
65	Remarkable morphological and metabolic regression as regarding the prevascular hyper metabolic lymph node , with estimated SUV 13.7 (pre-chemotherapy) and then on follow up it decreased to 8.9 , indicating good therapeutic response	100
66	Axial Fused PET/CT image shows a 20 mm nodule in the right lower lobe, with SUV = 22.6, proven	101