Transcultural Prevalence of Depression in Type 2 Diabetics Attending Endocrine Clinics in Both Egypt and Yemen

Submitted by

Ahmed Salem Saeed Basabbea
M.B., B.Ch
For Partial Fulfillment of Master Degree in Endocrinology

Under Supervision of

Prof. Dr. Mohammed R. Halawa

Professor of Internal Medicine and Endocrinology Faculty of Medicine - Ain Shams University

Dr. Ghada EL-Khouly

Consultant of Neuropsychiatry
Faculty of Medicine - Ain Shams University

Dr. Abdullah Saleh Bin-Nabhan

Assistant Professor of Internal Medicine Faculty of Medicine_Hadhramout University (Yemen)

Faculty of Medicine
Ain Shams University
2012

Acknowledgement

First and foremost, I am thankful to Allah, for without His grace, this work would have never been accomplished as we feel his great care, support and guidance in every step in our life.

I am genuinely and deeply indebted to **Dr. Mohmmed R. Halawa** Professor of Internal Medicine and Endocrinology, Faculty of Medicine, Ain shams University for his constant guidance, gentle encouragement, foresight and faith in me. Thank you for your help and unfailing support pushing me forward all the time, to overcome every obstacle in the way.

I would like to express my deepest appreciations to **Dr. GhadhaEl-kholy** consultant of neuropsychiatry, faculty of medidcine, Ain Shams University for her keen supervision, and valuable advice that guided me all the way through the various phases of this work, and for teaching me through her tremendous experience. Really, I owe to her much more than can be expressed. God blessing all of her faithful efforts.

My special thanks and cardinal appreciations to **Dr. abdellah S. Binnabhan** head of Internal Medicine depratment, Faculty of medicine, Hadhramout University for his continuous interest, precious suggestions and inexhaustible patience.

I would like to express my sincere appreciation to **Dr. Manal M. Abushady** Assistant Professor of Internal
Medicine and Endocrinology Faculty of Medicine - Ain
Shams University for her support at the beginning of this
work and final revision of it.

Lastly, but not least, I would like to express my thanks to all patients who participated in this study

Contents

Subjects	Page
List of Abbreviations	I
• List of Tables	II
• List of Figures	VI
• Introduction	1
• Aim of the Work	3
• Review of literature:	
- Chapter (1): Depression	4
- Chapter (2): Type 2 diabetes mellitus	20
- Chapter (3): Depression comorbid with typ	e 2 diabetes
mellitus	46
Subjects and Methods	83
• Results	91
• Discussion	129
• Conclusions	142
Recommendations	143
• Summary	145
• References	149
• Appendix	
Arabic Summary	

🕏 List of Figures 🗷

E List of Abbreviations &

List of Abbreviations

A1c	Glycosylated hemoglobin
ACEI	Angiotensin converting enzyme inhibitors
ADA	American diabetes association
ARB	Angiotensin receptor blockers
CHD	Coronary heart disease
CRH	Corticotropin releasing hormone
CVD	Cardiovascular disease
DKA	Diabetic ketoacidosis
DM	Diabetes mellitus
DSM-IV	Diagnostic and statistical manual of mental
	disorder-fourth edition
FPG	fasting plasma glucose
HHS	Hyperosmolar Hyperglycemic syndrome
HPA	Hypothalamic-pituitary adrenal axis
ICD-10	International classification of disease-10
IGT	Impaired glucose tolerance
OGTT	oral glucose tolerance test
SSRIs	selective serotonin reuptake inhibitors
T2DM	Type 2 diabetes mellitus
UKPDS	United kingdom prospective diabetes study

List of Tables

Table No.	Title	Pa
Tuble 110.	1100	ge
Table (1-1)	DSM_IV Diagnostic criteria of major	7
	depressive disorder	
Table (1-2)	ICD-10 diagnostic criteria of a depressive episode	8
Table (2-1)	Etiologic Classification of Diabetes Mellitus	24
Table (2-2)	Top 10 countries for estimated number of	26
	persons with diabetes in 2010 and 2030.	
Table (2-3)	Top 10 countries in prevalence of diabetes	27
	(20-79 age group)	
Table (2-4)	Causes of death in patients with type 1 and	29
	type 2diabetes.	
Table (2-5)	Insulin secretory defects in T2DM	32
Table (2-6)	Classical symptoms of diabetes	34
Table (2-7)	Criteria for the diagnosis of diabetes.	36
Table (2-8)	Advantages and disadvantages of Using of	36
	A 1C 6.5% (48 mmol/mol) as a cut - off	
	for making the diagnosis of diabetes	
Table (2-9)	clinical features of DKA	37
Table (2-10)	the clinical pictures of HHS	38
Table (2-11)	Neuropathic symptoms and signs in	42
	diabetes mellitus	
Table (2-12)	LEAP Program	45
Table (3-1)	Risk factors for depression in diabetes	58
Table (3-2)	Improving depression care in patients with	75
	diabetes	

Table No.	Title	Pa
Tuble 110.		ge
Table (1)	Comparing Egyptians and Yemeni groups	93
	regarding age, sex and religion.	
Table (2)	Comparing Egyptians and Yemeni groups	94
	regarding marital status, number of children and occupation	
Table (3)	Comparing Egyptians and Yemeni groups	97
	regarding social class, birth order, current	
	residence, smoking status and years of education	
Table (4)	Comparing Egyptians and Yemeni groups	99
	regarding complication, duration and	
	control of diabetes mellitus (DM).	100
Table (5)	Comparing Egyptians and Yemeni groups	100
	regarding to treatment of diabetes mellitus.	
Table (6)	Comparing Egyptians and Yemeni groups	101
	regarding to body mass index (BMI)	
Table (7)	Comparing Egyptians and Yemeni groups regarding comorbidities	102
Table (8)	Comparing Egyptians and Yemeni groups	103
	regarding drug for comorbidities	
Table (9)	Proportion of depression in Egyptians and Yemeni groups	104
Table (10)	Comparing non depressed and depressed	107
()	patients in Egyptians and Yemeni groups	
	regarding age, sex and religion.	
Table (11)	comparison between the non depressed	109
	and depressed groups as regard to marital	
	status, occupation, and number of children	

Table No.	Title	Pa
Table 10.	Title	ge
Table (12)	Comparing the non depressed and depressed patients regarding social class, birth order, current residence, smoking,	113
Table (13)	and number of cigarette per day Comparing non depressed and depressed participants in the two groups regarding diabetes mellitus duration in years and control	115
Table (14)	Comparing non depressed and depressed participants in both groups regarding diabetes related complications.	116
Table (15)	Comparing the non depressed and depressed participants in both groups regarding individual complications.	117
Table (16)	Comparing the non depressed and depressed participants in both groups regarding diabetes treatment.	118
Table (17)	Comparing the non depressed and depressed participants in both groups regarding medical co morbidities	119
Table (18)	the non depressed and depressed participants in both groups regarding drugs used for medical comorbidities	121
Table (19)	Comparing the non depressed and depressed participants in both groups regarding the number of drugs for medical comorbidities	122
Table (20)	Comparing the non depressed and depressed participants in both groups regarding the body mass index.	123

Table No.	Title	Pa
1401011101	Title	ge
Table (21 and	Comparing depressed Egyptians and	124
22)	Yemeni participants regarding	&
	sociodemographic characteristics.	125
Table (23)	Comparing depressed Egyptian and	126
	Yemeni participants regarding diabetes	
	related variables.	
Table (24)	Comparing depressed Egyptian and	126
	Yemeni participants regarding diabetes	
	related complications.	
Table (25)	Comparing depressed Egyptian and	127
	Yemeni participants regarding diabetes	
	medications.	
Table (26)	Predictors of depression in the both	128
	participating groups.	

🕏 List of Figures 🗷

List of Figures

Fig. No.	Title	Page
Fig. (1)	Comparing Egyptians and Yemeni groups regarding medical comorbidities.	102
Fig. (2)	Comparing the proportion of depression in both participating groups.	105

Scottish Intercollegiate Guidelines Network (2010), **Management of diabetes: A national clinical guideline,** available at www.sign.ac.uk

Introduction

is problem Diabetes mellitus a serious health worldwide. According to the international diabetes federation, 189million individuals have diabetes worldwide. The prevalence of diabetes is rapidly rising; worldwide the number of individuals with diabetes is expected to have doubled in 2025 (Scharm et al, 2009).

According to international diabetes federation, the prevalence of diabetes mellitus among age group between 20-79 years in Egypt is 11% in 2007 with a total number of 4.4 million persons affected (**I.D.F**, **2006**).

Depression is a serious medical condition that affects thoughts, feelings, and the ability to function in everyday life. Depression can occur at any age. National institute of mental health (NIMH)-sponsored studies estimate that 6 percent of 9- to 17-year-olds in the U.S. and almost 10 percent of American adults, or about 19 million people age 18 and older, experience some form of depression every year (**Braiana** *et al.* **2008**).

Depression may adversely impact outcomes of chronic illnesses, such as diabetes, in several ways. Depression has been shown in patients with diabetes to be associated with poor adherence to glucose monitoring diet exercise regimens and taking medication as prescribed (**Dunning and Ward**, 2008).

₹ Introduction 🗷

The high prevalence of depressive symptoms among patients with diabetes (between 18% and 35%) has been well documented in several studies. Patients with diabetes and co morbid depression display higher functional impairment and work loss and poorer self-management behavior, and they have more co morbidities (**Miranda** *et al*, 2009)

In an Egyptian study, 44.7% of diabetic children and adolescents were found to have psychiatric disorder, the most common being depression as 22.4% are showing depressive symptoms and 17.6% have major depression (**Shaker**, **2003**).

In a study of primary care diabetic patient in Al-Ain, Uuited Arab emirates, the author found that 33.8% of diabetic patients had psychiatric co morbidity. All psychiatric cases were identified as mild or moderate in severity. Depression was the most common disorder (16.9% of diabetic patients) (Sulaiman et al, 2010).

Because of lack of published studies on the prevalence of the depression in type 2 diabetic patients in Egypt and Yemen, we will carry on this study to evaluate the prevalence of depression among type 2 diabetics Egyptian and Yemeni adults, and correlate it to cultural differences.

₹Aim of the Study ≤

Aim of the Study

To study the prevalence of depression in a sample of type 2 diabetic patients taken from diabetic and endocrine outpatient clinics in both Egypt and Yemen and to correlate it to cultural differences.

Chapter (1) **Depression**

Depression

The word depression comes in Latin deprime, meaning "to press down". The term was first applied to mood at the seventeenth century. Depression was first named as a condition about 2,400 years ago by the famous ancient Greek doctor "Hippocrates", who called it melancholia (Gilbert, 2001).

Depression refers to a wide range of mental health problems characterized by the absence of a positive affect (a loss of interest and enjoyment in ordinary things and experiences), low mood and a range of associated emotional, cognitive, physical and behavioral symptoms (Lewinsohn et al., 2000).

physical Behavioral and symptoms typically include irritability, withdrawal. social tearfulness. an exacerbation of pre-existing pains, pains secondary to increased muscle tension (Gerber et al., 1992), a lack and diminished activity, of libido, fatigue although agitation is common and marked anxiety frequent. Typically there is reduced sleep and lowered appetite (sometimes leading to significant weight loss), but for some people it is recognized that sleep and appetite are increased. A loss of interest and enjoyment in everyday

Depression \mathcal{Z}

life, and feelings of guilt, worthlessness and that one deserves punishment, are common, as are lowered self-esteem, loss of confidence, feelings of helplessness, suicidal ideation and attempts at self-harm or suicide.

Cognitive changes include poor concentration and reduced attention, pessimistic and recurrently negative thoughts about oneself, one's past and the future, mental slowing and rumination (Cassano and Fava, 2002).

Major depression is generally diagnosed when a persistent low mood and an absence of positive affect are accompanied by a range of symptoms, the number and combination needed to make a diagnosis being operationally defined (APA, 1994).

People with severe depression may also develop (hallucinations psychotic symptoms and/or delusions). thematically commonly consistent with the most self-blaming cognitions negative. and low mood typically encountered in major depression, others may develop psychotic symptoms unrelated to mood (Anderson, 2010).

The average age of the first episode of major depression occurs in the mid-20s and, although the first episode may occur at any time from early childhood through to old age, a substantial proportion of people have their first depression in childhood or adolescence (Fava and Kendler, 2000).