Study of Sleep Disordered Breathing Among Obese Children

Thesis

Submitted for partial fulfillment of Master degree in pediatric

By

Mostafa Abd El-Latef Mostafa El-Mokadem M.B.,B.Ch.,(2003)

Under Supervision of Professor Doctor/ Karima Ahmed Abdel Khalik

Professor of Pediatrics Faculty of Medicine, Ain Shams University

Doctor/Terez Boshra Kamel

Lecturer of Pediatrics Faculty of Medicine, Ain Shams University

Doctor/ Omneya Ibrahim Youssef

Lecturer of Pediatrics Faculty of Medicine, Ain Shams University

> Faculty of Medicine Ain Shams University 2009



ACKNOWLEDGEMENT

Thanks to **ALLAH**, Most Gracious, Most Merciful, who gives everything we have and gave me the power and patience to finish this work.

I wish to express my thanks and deepest appreciation to **Prof. Dr. Karima Ahmed Abdel Khalik**, Professor of Pediatrics, Faculty of Medicine, Ain Shams University, for her keen guidance, kind supervision, valuable advice and continuous encouragement which made possible the completion of this work.

I would like to record my cardinal thanks to **Dr. Terez Boshra Kamel**, Lecturer of Pediatrics, Faculty of Medicine, Ain Shams University, for her great care, valuable instruction, constant help and helpful advice.

Also my great thanks to **Dr. Omneya Ibrahim Youssef**, Lecturer of Pediatrics, Faculty of Medicine, Ain Shams University, for her continuous supervision, stimulating support and valuable suggestions.


My deep thanks to all staff members of Pediatric Department for their help and kind cooperation.

I would like to express my hearty thanks to my father, my mother and all my family members for their support, understanding and tolerance till this work has been completed.

Last but not least my sincere thanks and appreciation to all patients participated in this study.

Mostafa El-mokadem

DEDICATION



CONTENTS

| | Page |
|------------------------|------|
| List of abbreviations | 1 |
| List of tables | Ш |
| List of figures | VII |
| Introduction | 1 |
| Aim of the work | 2 |
| Review of literature | 3 |
| Subjects and Methods | 69 |
| Results | 75 |
| Discussion | 98 |
| Summary and conclusion | 113 |
| Recommendations | 117 |
| References | 118 |
| Appendix | |
| Arabic summary | |

List of Abbreviations

ACTH : Adrenocorticotropic hormone

AgRP : Agouti-related Peptide
AHI : Apnea Hypoapnea Index

AI : Apnea Index

Alfa- MSH : Alfa Melanocyte stimulating hormone

ALT : Alanine transaminase

ARC : Arcuate Nucleus

AST : Aspartate aminotransferase

BMI : Body mass index
Blood pressure

CART : Cocaine-Amphetamine-Regulated

CCK : Cholecystokinin

Cm : Centimeter

CNS : Central nervous system

CO₂ : Carbon dioxide

CPAP : Continuous positive airway pressure

CRF : Corticotropin-releasing factor

CRP : C -reactive protein

CT : Computerized tomography

ECG : Electro-cardiography

EEG : Electro-encephalo-graphy

EMG : Electro-myography
EOG : Electro-oculography

GH : Growth hormone

HDL : High density lipoprotein

IGF-1 : Insulin-like growth factor 1

IL-6 : Interleukin-6

Kg : Kilogram

List of Abbreviations

LDL : Low density lipoprotein

M : Meter

MAOIs : Monoamine oxidase inhibitors

MC4R : Melanocortin receptor-4

MRI : Magnetic resonance imaging

NHANES : National Health and Nutrition Examination

Survey

NPY : Neuropeptide Y

NREM : Non rapid eye movement

 O_2 : Oxygen

OSA : Obstructive sleep apnea
PC-1 : Prohormone convertase-1
PCOS : Polycystic ovary syndrome

POMC : Pro-opiomelancortin

PSG : Polysomnography

PYY : Peptide YY

RDI : Respiratory Distress Index

REM : Rapid eye movement SES : Socioeconomic status

SBD : Sleep breathing disordered

SD : Standard deviation SFT : Skin fold thickness

TGs : Triglycerides

TLD : Traffic light diet

TNF-alpha : Tumor necrosis factor-alpha

Transcript Neurons

TSH : Thyroid stimulating hormone

W/H : Waist/Hip ratio

WC : Waist circumference

WHO : World Health Organization

| Table | Subject | Page |
|-------|--|------|
| (1) | BMI classification of children and adolescent | 14 |
| (2) | Cutoff values of Waist circumference in centimeters for 90th percentiles according to age and sex | 18 |
| (3) | Laboratory tests of obesity | 19 |
| (4) | Evaluating and Testing of comorbidities of obesity | 19 |
| (5) | Suggestion for prevention of childhood obesity | 28 |
| (6) | Factors predisposing to OSA and hypopnea | 45 |
| (7) | Common and different features in the clinical presentation of pediatric obstructive sleep apnea types I and II | 65 |
| (8) | Abramson and Abramson social status classification | 70 |
| (9) | Grades of tonsillar enlargement | 72 |
| (10) | Numbers and distribution percentages of subjected children according to sex, order of birth and social class | 75 |
| (11) | Anthropometric measurements of studied children | 75 |

| (12) | Mean age of presentation and duration of snoring and obesity | 76 |
|------|---|----|
| (13) | Numbers of children complaining of obstructed sleep apnea symptoms and their percentage | 76 |
| (14) | Distribution of blood pressure values in studied children | 77 |
| (15) | Throat examination finding of studied children | 78 |
| (16) | Apnea related data of studied children | 78 |
| (17) | Prevalence of obstructed sleep apnea in studied obese children | 79 |
| (18) | Prevalence of different degrees of OSA in studied children according to apnea hypopnea index (AHI) | 80 |
| (19) | Comparison between obese children with and without obstructed sleep apnea regarding gender, order of birth and social class | 81 |
| (20) | Comparison between obese children with and without obstructed sleep apnea according to anthropometric measures | 82 |
| (21) | Agreement of obstructed sleep apnea symptoms with apnea hypopnea index in diagnosis of OSA | 83 |
| (22) | Comparison between mean values of apnea hypopnea index in patients with and without OSA symptoms | 84 |

| (23) | Comparison between patients with different grades of OSA regarding OSA symptoms | 85 |
|------|---|----|
| (24) | Comparison between obese children with and without obstructed sleep apnea regarding blood pressure measurement | 86 |
| (25) | Comparison between children with different grades of apnea (according to apnea hypopnea index) as regards blood pressure measurements | 87 |
| (26) | Distribution of different grades of tonsillar enlargement in obese children with and without obstructed sleep apnea | 88 |
| (27) | Mean apnea hypopnea index values in different grades of tonsillar enlargement in studied children | 89 |
| (28) | Comparison between obese children with and without obstructed sleep apnea regarding the lowest O_2 saturation values | 90 |
| (29) | Comparison between obese children with and without obstructed sleep apnea according to mean values of lowest O_2 saturation and number of desaturation events /h as detected by apnea link device | 91 |
| (30) | Comparison between different grades of OSA (according to apnea hypopnea index) regarding the lowest O ₂ saturation values | 92 |
| (31) | Comparison between obese children with and | 92 |

| | without obstructed sleep apnea according to mean values of minimum pulse frequency and snoring events/h as detected by apnea link device | |
|------|--|----|
| (32) | Correlation between apnea index values and age of presentation and duration of obesity | 93 |
| (33) | Correlation between apnea hypopnea index values and age of presentation and duration of obesity | 95 |
| (34) | Correlation between apnea index values and age of presentation and duration of snoring | 96 |
| (35) | Correlation between apnea hypopnea index values and age of presentation and duration of snoring | 97 |

List of Figures

| Figure | Subject | Page |
|--------|--|------|
| (1) | Prevalence of obesity among children aged 2-19 years according to National Health and Nutrition Examination Surveys | 3 |
| (2) | Prevalence of obesity by Race/Ethnicity among children aged 12–19 years according to National Health and Nutrition Examination Surveys | 5 |
| (3) | Endogenous control of appetite | 6 |
| (4) | Diagrammatic representation of the effects of leptin on energy homeostasis. | 9 |
| (5) | The melanocortin pathway and its connection to leptin | 10 |
| (6) | BMI growth chart for age percentiles for boys and girls from 2 to 21 years | 15 |
| (7) | BMI growth chart for age percentiles for Egyptian boys and girls from 2 to 21 years | 16 |
| (8) | The percentiles for Triceps SFT | 17 |
| (9) | The percentiles for Subscapular SFT | 17 |
| (10) | Normal airway, Hypopnea and Obstructive sleep apnea | 37 |

List of Figures

| (11) | Anatomy of throat and anatomical position of Adenoids | 39 |
|------|--|----|
| (12) | Model of counterbalancing forces that influence airway patency | 43 |
| (13) | Tonsillar hypertrophy, a common cause of obstruction in the pediatric population | 46 |
| (14) | Plain x ray adenoid view showing adenoidal hypertrophy with narrowing of airway | 46 |
| (15) | Nasal continuous positive airway pressure usage in OSA | 62 |
| (16) | Apnea Link Device | 73 |
| (17) | Illustration of the Apnea Link Device usage steps | 73 |
| (18) | Percentage of children complaining of obstructed sleep apnea symptoms | 77 |
| (19) | Distribution of blood pressure values in studied children | 77 |
| (20) | Prevalence of obstructed sleep apnea | 79 |
| (21) | Prevalence of different degrees of OSA according to Apnea Hypopnea Index (AHI) | 80 |
| (22) | Comparison between obese children with and without Obstructed sleep apnea as regards body mass index and waist/hip ratio | 82 |

| (23) | Comparison between obese children with and without Obstructed sleep apnea regarding blood pressure measurements | 85 |
|------|---|----|
| (24) | Comparison between patients with different grades of apnea (according to apnea hypopnea index) as regards blood pressure measurements | 87 |
| (25) | Comparison between obese children with and without obstructed sleep apnea regarding distribution of different grades of tonsillar enlargement | 88 |
| (26) | Relationship between apnea hypopnea index and different grades of tonsillar enlargement in studied children | 89 |
| (27) | Comparison between obese children with and without obstructed sleep apnea regarding the lowest O_2 saturation | 90 |
| (28) | Comparison between obese children with and without obstructed sleep apnea according to mean values of lowest O ₂ saturation and number of desaturation events /h | 91 |
| (29) | Comparison between obese children with and without obstructed sleep apnea according to mean values of minimum pulse frequency and snoring events/h | 93 |

List of Figures

| (30) | Correlation between Apnea Index values and duration of obesity in studied patients | 94 |
|------|--|----|
| (31) | Correlation between Apnea Index values and age of presentation of obesity in studied patients | 94 |
| (32) | Correlation between Apnea Hypopnea Index values and duration of obesity in studied patients | 95 |
| (33) | Correlation between Apnea Hypopnea Index values and age of presentation of obesity in studied patients | 96 |
| (34) | Correlation between Apnea Hypopnea Index values and age of presentation of snoring in studied patients | 97 |