THE ROLE OF THE MAGIC FRUIT PLANT IN CONTROLLING THE SPREAS OF CANCERCELLS EX-VIVO

BY

Mofeed Mohamed Ahmed Selim

B. SC Faculty of science, Aim shams university 1981

Diploma in environmental sciences (department of Medical science Aim shams university 1994

A thesis submitted to in partial fulfillment of

The requirement for the Master degree
Environmental science
Department of Biology and Natural
Science

Institute of Environmental studies and Research
Aim shams university

APPROVAL SHEET THE ROLE OF THE MAGIC FRUIT PLANT IN CONTROLLING THE SPREAS OF CANCERCELLS EX- VIVO

BY

Mofeed Mohamed Ahmed Selim

B. SC Faculty of science, Aim shams university 1981

Diploma in environmental sciences (department of Medical science Aim shams university 1994

His thesis towards a Master Degree in Environmental science has been Approved by :

Signature

1- Prof. Dr. Rashwan Mohamed Farag

Prof of Biochemistry faculty of Medicine Ain Shams university

2- Prof. Dr . Fathy Mohamed Tash

Prof of Biochemistry faculty of Medicine Ain Shams university

3- Prof. Dr. Motawa a Eiss a El- Housseini

Prof of Biochemistry – cancer biology department – National cancer institute – Cairo university

4- Prof . Dr . Ibrahim Ali Elagoza

Prof of physiology – Faculty of science - – Cairo university

THE ROLE OF THE MAGIC FRUIT PLANT IN CONTROLLING THE SPREAS OF CANCERCELLS EX-VIVO

BY

Mofeed Mohamed Ahmed Selim

B. SC Faculty of science, Aim shams university 1981

Diploma in environmental sciences (department of Medical science Aim shams university 1994

A thesis submitted to in partial fulfillment

of

The requirement for the Master degree

in

Environmental science

Department of Biology and Natural Science
Institute of Environmental studies and Research
Ain shams university

Under the supervision of:

1- Prof. Dr. Rashwan Mohamed Farag

Prof of Biochemistry faculty of Medicine Ain Shams university

2-1-Prof. Dr. Shadia Abed Hamid Fathy

Prof of Biochemistry faculty of science Ain shams university

3- Prof. Dr. Motawa a Eiss a El- Housseini

Prof of Biochemistry – cancer biology department – National cancer institute – Cairo university

ACKNOWLEDGEMENTS

I Wish to express my deepest thanks and appreciation to **Dr. Rashwan Mohamed Farage**. Head of Biochemistry Department, Faculty of Medicine, Ain Shams University for his interest in the present work, kind supervision and continuous encouragement.

It is a pleasure to offer my sincere thanks and gratitude to **Dr. Shadia Abdel-Hamed Fathy**. Professor of Biochemistry, Faculty of Science, Ain Shams University for her kind supervision and useful advice.

I am much indebted to **Dr. Motawa Eissa El-Housseini**, Professor of Biochemistry, Cancer Biology Department, National Cancer Institute, Cairo University for his sincere cooperation and efforts throughout the work.

I wish express my deepest thanks to **Dr. Farid Abo-Bedair**, Assistant professor, Cancer Biology Department, National Cancer Institute, Cairo University and **Dr. Tarek Dardeer**, Assistant professor, Zoology Department Faculty of Science, Cairo University for their help during the preparation of the manuscript.

CONTENTS

INTRODUCTION1
AIM OF WORK
REVIEW OF THE
LITERATURE3
Cancer: origin and terminology3
Malignant transformation of cells5
Oncogenes and cancer induction7
Breast cancer8
Epidemiology of breast cancer
9
Risk factor for breast cancer11
1- Familial factors 11
2- gender 11
3- age
4- hormonal factors
5- dietary and lifestyle factors
13

6- radiation and age at exposure
13
Symptoms and screening for breast cancer14
Breast cancer diagnosis
14
Pathologic classification of tumor types
16
Invasive ductal carcinoma
16
Invasive lobular carcinoma
16
Staging of breast cancer
18
Clinical staging
18
Pathologic staging
18
Pathological grading of breast carcinomas
21
Prognostic and predictive factors in breast cancer22
Estrogen and progesterone receptor expression22
Her 2 (her2/neu , epidermal growth factor receptor -2,c-erbb2)
stauts
22
Lymphoyascular invasion
23
Angiogenesis
23

Proliferation rate24
Plasminogen activator system
Molecular and genomic factors
Tumor Infiltrating Lymphocytes26
Adaptive Immunotherapy 28
Lymphocyte tumor homing29
Lymphocyte markers and CD nomenclature system31
Activation of lymphocytes31
Synthesis of new proteins
Cellular proliferation
Differentiation into effectors cells32
Differentiation into memory cells35
Immunity to Tumors.,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,

Interleukin -2 immunotherapy37
The Magic fruit
Patients and Methods
Reagents and chemicals
Methods45
1- preparation of extract of magic fruit45
II – Enzymatic digestion Method for Tils separation form the Tumor Tissue.
III- Isolation of peripheral Mononuclear cells (PBMC)46
IV- Separation of Tils from fine needle47
Results
Clinic pathological data of the patients52
Liver and kidney function tests of the patients53
Effect of treatment by the magic fruit extract on the cell count54
Effect of treatment by the magic $$ extract on interferon gamma level (IFN- $\!\gamma$)
55

Correlations between the cell count and liver and kidney function
tests
56
Correlations between IFN-γ and liver and kidney function
tests59
Correlations between IFN-γ and liver and the cell count
61
Association between the clinic pathological and the cell count
62
Association between the clinic pathological date and the IFN-γ
level65
Discussion
71
Summary
78
References
81
Arabic summary
98

List of Tables:-

- 1-Table I: WHO Classification of Breast Carcinoma
- **2-Table II**: TNM stage definitions.
- 3 Table III: Nottingham combined histologic grades
- **4- Table (IV):** Food value per 100 g of edible portion of the magic fruit
- 5 Table 1 Clinicopathological data of the patients
- **6 Table 2.** Liver and kidney function tests of the patients
- **7- Table 3.** Effect of treatment of the cells by the magic fruit extract on the cell count.
- **8 Table 4.** Effect of treatment by the magic fruit extract on interferon gamma level (IFN- γ).
- **9 Table 5.** Correlations between the cell count and liver and kidney function examinations
- **10 Table 6**. Correlations between the IFN-γ level and liver and kidney function examinations
- 11 Table 7. Correlations between the IFN-γ level and the cell count before and after treatment by the magic fruit extract.
- **12 Table 8.** Association between the cloinicopathological and the cell count
- **13- Table 9.** Association between the cloinicopathological and the IFN-γ level.
- 14 Appendix Table. The raw data used in the present study

List of Figs

- Fig. 1. Effect of treatment of the cells by the magic fruit extract on the cell count. The count after treatment showed a highly significant increase than before treatment.
- Fig. 2. Effect of treatment of the cells by the magic fruit extract on IFN-γ level. The level after treatment showed a highly significant increase than before treatment.
- Fig. 3. Correlation between the cell count before treatment and ALP activity. The correlation is negative and significant (r = -0.497, p = 0.035).
- Fig. 4. Correlation between the cell count after treatment and ALP activity. The correlation is negative and significant (r = -0.497, p = 0.035).
- Fig. 5. Correlation between the cell count before treatment and urea. The correlation is positive and significant (r = 0.543, p = 0.019).
- Fig. 6. Correlation between the cell count after treatment and creatinin. The correlation is positive and significant (r = 0.488, p = 0.039).
- Fig. 7. Correlation between the IFN- γ before treatment and creatinin. The correlation is negative and significant (r = -0.480, p = 0.043).
- Fig. 8. Association between cell count and the tumor size.
- Fig. 9. Association between cell count and the tumor grades.
- Fig. 10. Association between cell count and the tumor stage.
- Fig. 11. Association between cell count and the LN status.
- Fig. 12. Association between cell count and the tumor size.
- Fig. 13. Association between cell count and the tumor grade.
- Fig. 14. Association between cell count and the tumor stage.
- Fig. 15. Association between cell count and the LN status.
- Fig.16. Tumor infiltrating lymphocytes in culture before treatment. (X 100). The cell number is very little if compared to the cell culture after treatment.

- Fig.17. Tumor infiltrating lymphocytes in culture after treatment. (X 100). The cell number is highly increased after treatment.
- Fig.18. Tumor infiltrating lymphocytes in culture before treatment. (X 400). The cell number is very little if compared to the cell culture after treatment.
- 11Fig.19. Tumor infiltrating lymphocytes in culture after treatment. (X 400). The cell number is highly increased after treatment.

Abbreviations

1- TilsTumor – Infiltrating lymphocytes
2- Il2Interleukin- 2
3- RNA Ribonucleic acid
4- DNADeoxy ribonucleic acid
5- ILCInvasive lobular carcinoma
6- TDCtype invasive ductal carcinoma
7- MFMagic fruit
8- SV 40 simian virus
9- BRCA1Breast cancer antigens
10- MTI Magnetic Resonance imaging
11- AJCCAmerican joint Committee on cancer
12- ER+ Estrogen Receptor positive
13- HER-2 Human Epidermal Receptor 2
14- EGFREpidermal growth factor Receptor
15- IHCImmunohistochemistry
16-CTLsCytolytic T cell
17- UPAUrokinase plasminogen activator
18-PAI-1Plasminogen activator inhibitor-1
19- CTLsCybolybic Tcell
20- APC8Antigen- presenting cells
21-IFN-8inberferons gamma
22- LAKLympakinge – activated killer
23-FC8Fetal calf serum
24- AlpAblalime phoshatars
25-L.NLemphnodes

The jujube is a member of Rhnaceae family. its botanical name is Ziziphus jujube or Chinese date. It is distributed throughout much of the Mediterranean region at least 3000 years ago (Farums medicine http:// www.Suudi medicine 2005.net) it helps in many of vital activities and functions such as weight gain , antiallergenic, improving stamina and strength, mildly sedating, improving of liver function , stimulating the immune system , and helping to efficiently get the energy form food and giving energy. It could be used as Mill Lam. In Sudan , it is fermented and prepared into a confection but in India, dry fruit powder is also used. It is 250 times sweeter than refined sugar and without calories (Clevallier, A.1996).

The major breakthrough is the treatment of terminals cancer with new approach of cell treatment. The idea is to exploit a subset of T-cells called tumor – infiltrating lymphocytes (TIIs) found deep inside cancerous tissue. These killer T-cells attack dividing cells and provide a natural protection against cancer (Ro senberg, 2001). The T-cell theraby might be more effective than other types of carcenoginic therapy.

Tumor – infiltrating lymphocytes (TILs) can be grown in vitro in a medium containing interleukin -2 in clinical trials (Bethesda 2005). patients with metastatic malignant melanomas were treated with intrleukin-2 plus the adoptive transfer of autologous (TILs) (Prevost-Blonde et al., 1998)

It can be used in the future in clinical trials post surgery in the Natural Cacer Institute. Lysis of autologous tumor cells by TIIs was significantly higher for responding than for non-responding patients.

AIM OF THE WORK

The aim of the present work is to study the possible effect (s) of the magic fruit extract on the viability, and count of TILs as well as on the level of interferon – gamma in the medium containing the cell culture. Tumor excision may not be a totally curative treatment of cancer, but the activation of TILs by using a natural product (plant) (Magic Fruit) may possibly inhibit the tumor cells propagated ex-vivo.