

Multislice CT angiography of peripheral pulmonary embolism

ESSAY

*SUBMITTED FOR FULFILLMENT
OF Msc degree IN RADIODIAGNOSIS*

BY

Ahmed Mohamed Salah
M.B.B.CH.

SUPERVISORS

Prof .Dr. Magdy Ibrahim Basiony
*Professor of Radiodiagnosis
Faculty of Medicine
Cairo University*

Dr. Noha Hosam El Din Behairy
*Lecturer of Radiodiagnosis
Faculty of Medicine
Cairo University*

Cairo University
Faculty of Medicine
2009

Abstract

The most important advantage of CT over other imaging modalities is that both mediastinal and parenchymal structures are evaluated, and thrombus is directly visualized. Studies have shown that up to two thirds of patients with an initial suspicion of PE receive another diagnosis, some with life threatening diseases, such as aortic dissection, pneumonia, lung cancer and pneumothorax. Most of these diagnoses are amenable to CT visualization, so that in many cases a specific etiology for the patient's symptoms and important additional diagnosis can be established.

Key Words :

Contrast Media - Inferior Vena Cava - Pulmonary Embolism .

Acknowledgment

First of all, I thank God for all his blessings and care, that helped me finish this work.

I would like to express my thanks and deep gratitude to Prof. Dr. Magdy Basiony, Professor of radiodiagnosis, Faculty of medicine, Cairo University, for his guidance, advise and thoughtful remarks and being there for me and my colleagues all the time.

Also I would like to express my thanks and appreciation to Dr. Noha Al Behairy, Lecturer of radiodiagnosis, Faculty of medicine, Cairo University, for her continuous encouragement, help and valuable comments.

Last but not least, I would like to thank my family for their help and encouragement and I would like to dedicate this work to them.

List of Figures

Figures	Content	Page
Fig. 1	Lung Anatomy (Quoted from Stranding et al., 2008).	4
Fig. 2	Pulmonary arteries and veins (Quoted from Stranding et al., 2008)	5
Fig. 3	The relations of the pulmonary arteries and primary bronchi seen from the front. (Quoted from Stranding et al., 2008).	8
Fig. 4	A, Anterior view showing main pulmonary trunk passing posteriorly. B, Posterior view with aorta and spine removed. (Quoted from Ulfacker et al, 2007).	9
Fig. 5	Distribution of the pulmonary artery in both lungs (Quoted from Ulfacker et al, 2007)	14
Fig. 6	Schematic drawing of the pulmonary veins and left atrium. (Quoted from Ulfacker et al, 2007)	16
Fig. 7	Chest CT Showing Pulmonary Artery Bifurcation (Quoted from Ryan et al., 2004)	17
Fig. 8	MHCT pulmonary angiography. Rotational (“paddlewheel”) 2D reconstruction image shows ability of	18

	MHCT to image subsegmental pulmonary arteries (arrow). (Quoted from Goo et al, 2003).	
Fig. 9	MDCT shows reconstructed coronal orientation of peripheral pulmonary vasculature. (Quoted from Goo et al, 2003).	19
Fig. 10	Bolus tracking. A,B,C,D: Coned-down views of axial images sequentially obtained at the same exact level following a bolus of intravenous contrast media, with a region-of-interest cursor placed over the main pulmonary artery in each image. E: A real-time graph demonstrating the level of contrast enhancement within the main pulmonary artery as a function of time. (Quoted from Naidich et al., 2007).	34
Fig. 11	Effects of image windowing. A,B: Identical contrast-enhanced axial sections through the right main pulmonary artery imaged with window widths of 400 HU (A) and 1,000 HU (B), respectively, with corresponding window widths of 40 HU (A) and 150 HU (B).. C,D: Identical coned-down sections through the inferior portion of the right hilum (Quoted from Naidich et al, 2007).	38

Fig. 12	Use of maximum intensity projections (MIPs). A: Axial 10-mm maximum MIP image demonstrates multiple left-sided PE. B: A 7-mm off-axis coronal MIP (Quoted from Naidich et al, 2007).	40
Fig. 13	Paddlewheel Reconstructions. A,C: Three 1-mm multiplanar reconstruction images selected from a set of images obtained using a 360-degree paddlewheel reconstruction algorithm. (Quoted from Naidich et al, 2007).	41
Fig. 14	Multiplanar reconstructions. A: 2-mm axial image demonstrates an completely occluded subsegmental pulmonary artery (arrow). B,C: Off-axis axial and coronal 5-mm maximum intensity projection reconstructions, respectively (Quoted from Naidich et al, 2007).	42
Fig. 15	Image depicting an experimental method for providing automated three-dimensional surface shaded renderings of the pulmonary vasculature. (Quoted from Naidich et al, 2007).	44
Fig. 16	Computer-assisted diagnosis. Axial (top) and coronal images (bottom), as viewed using an experimental CAD device. (Quoted from Naidich et al, 2007).	45

Fig. 17	Vessel bifurcation. (A) A small lucency in a subsegmental vessel in the right upper lobe. (B) Vessel bifurcation at this site.(Quoted from Phillip Boisselle and White, 2007)	48
Fig. 18	Mucous plug (Quoted from Phillip Boisselle and White, 2007)	50
Fig. 19	Transverse gadolinium-enhanced multi-detector row CT angiograms. Scans obtained at the levels of the (a) carina and (b) right bronchus intermedius. (Quoted from Martin Remy Jardin et al, 2006).	53
Fig. 20	Transverse gadolinium-enhanced multi detector row CT angiograms. Obtained at levels of (a) tracheal bifurcation, (b) right bronchus intermedius, and (c) lower lobes. (Quoted from Martin Remy Jardin et al, 2006).	54
Fig. 21	Transverse gadolinium-enhanced multi-detector row CT angiograms. Scans obtained at the levels of the (a) right bronchus intermedius, (b) right middle lobe bronchus, and (c) lower lung lobes. (Quoted from Martin Remy Jardin et al, 2006).	55
Fig. 22	Acute pulmonary embolus (PE). A partial-filling defect is seen in a right lower lobe (RLL) artery. On the left there are	62

	segmental pulmonary emboli in left lower lobe (LLL) (Quoted from Philip Boiselle and White, 2007).	
Fig. 23	Acute pulmonary embolus (PE). “Railway track sign” of clot floating in the anterior segmental left upper lobe (LUL) artery. A mural thrombus forming acute angles with the vessel wall is present in the RUL artery. (Quoted from Phillip Boiselle and White, 2007).	63
Fig. 24	CT scan (A) shows a pulmonary embolus within the postero-basal subsegmental of the right lower lobe artery (B) shows a wedge-shaped non enhancing soft tissue with small irregular air inclusions in the corresponding right lower lobe, “Hampton’s hump.” (Quoted from Phillip Boiselle and White, 2007).	65
Fig. 25	Post-processing techniques. Small subsegmental embolus that was primarily missed on axial section (A) but picked up on axial thin-slab maximum intensity projection (arrow) (B). (Quoted from Phillip Boiselle and White, 2007).	66
Fig. 26	Chronic PE CT scan (A) and axial oblique MIP image (B) (Quoted from Phillip Boiselle and White, 2007).	67
Fig. 27	Paddlewheel reformation method. (a)	69

	<p>Drawing shows how paddlewheel slabs pivot on a central horizontal axis between the lung hila. (b) Lateral scout image for paddlewheel reformations with central axis at main pulmonary artery bifurcation. (c) Paddlewheel image at level of main pulmonary artery bifurcation. (Quoted from Simon M et al, 2001).</p>	
Fig. 28	<p>Paddlewheel versus coronal and axial images show improved efficiency of review and enhanced assessment of embolic burden. (a) Paddlewheel image (b) Four coronal images are required for visualization of extent of embolism. (c) Multiple axial images are required for visualization of extent of embolism (arrow). (Quoted from Chiang et al, 2003).</p>	71
Fig. 29	<p>Chronic pulmonary embolus (PE). CT scan mediastinal window through mid-zone reveals marginated mural thrombus in the right inferior lobar pulmonary artery (arrow). Bilateral pleural effusions are present. (Quoted from Phillip Boiselle and White, 2007).</p>	73
Fig. 30	<p>Mosaic perfusion in a patient with chronic pulmonary embolus (PE). High-resolution CT through the upper lobes (A). Expiratory scan (B). (Quoted from Phillip</p>	75

	Boiselle and White, 2007).	
Fig. 31	Contrast-enhanced 16-detector row CT study obtained with 0.75-mm collimation. Consecutive transverse sections show isolated peripheral pulmonary embolus (arrows) in a subsegmental pulmonary artery in segment 9 of the left lung. (Quoted from Schoepf & Costello, 2004).	77
Fig. 32	Oblique sagittal multiplanar reformation shows an embolus (arrow). (Quoted from Schoepf & Costello, 2004).	78
Fig. 33	Transverse contrast-enhanced 16-detector row CT image. Isolated peripheral pulmonary embolus (arrow) in sixth-order pulmonary arterial branch in segment 8 of the right lung is shown. (Quoted from Schoepf & Costello, 2004).	78
Fig. 34	CT angiography show isolated pulmonary emboli (arrows) in segmental and subsegmental arteries in the right middle lobe of the lung and allow detailed visualization of course of obliquely oriented vessels and of isolated filling defects in segmental and subsegmental branches. (Quoted from Schoepf & Costello, 2004).	79
Fig. 35	Pulmonary infarct. Frontal chest	83

	radiograph reveals wedge-shaped area of air-space disease in left costophrenic sulcus. (Quoted from konstantinide et al, 2007).	
Fig. 36	Frontal chest radiograph in a patient with documented acute PE demonstrating enlargement of the right interlobar pulmonary artery-the “sausage” appearance. (Quoted from konstantinide et al, 2007).	83
Fig. 37	Frontal (A) and lateral (B) chest radiographs demonstrating lower lobe consolidation and moderate bilateral pleural effusions. The anteriorly located opacity in the right lung represents a pulmonary infarct. (Quoted from konstantinide et al, 2007).	84
Fig. 38	Acute pulmonary embolism. Perfusion scan reveals multiple wedge shaped perfusion defects. (Quoted from konstantinide et al, 2007).	85
Fig. 39	Acute pulmonary embolism. Selective digital subtraction angiography in left pulmonary artery. (Quoted from konstantinide et al, 2007).	87
Fig. 40	Pulmonary angiogram demonstrating an abrupt vascular cutoff representing acute	87

	pulmonary embolism. (Quoted from Coche et al., 2006).	
Fig. 41	A , Oblique multiplanar reformatted MR angiogram. Image quality suffers from motion artifacts and allows segmental analysis at best. B , Double oblique maximum intensity projection of CT angiogram shows peripheral saddle embolus in segment 9 of right lower lobe. (Quoted from Kluge et al, 2006).	90
Fig. 42	Acute pulmonary embolism: magnetic resonance angiography. A,C: Coronal images obtained in a normal subject demonstrating improved spatial resolution and visualization of peripheral pulmonary arteries with progressively thicker maximum intensity projections (MIPs) (A through C, respectively). (Quoted from Naidich et al., 2007).	91

List of tables

Table	Content	Page
Table no. 1	Computed Tomography Pulmonary Embolism Protocols	29
Table no. 2	Diagram illustrating scheme for pulmonary artery diagnosis	93

List of Abbreviations

CAD	Computer Aided Diagnosis
CM	Contrast Media
CNR	Contrast to Noise Ratio
CPTE	Chronic Pulmonary Thromboembolism
CT	Computed Tomography
CTA	Computed Tomography Angiography
CTP	Computed Tomography Perfusion
CTPA	Computed Tomography Pulmonary Angiography
CTV	Computed Tomography Venography
CWD	Continuous Wave Doppler
DLCO	Carbon Monoxide Diffusion Capacity
DSA	Digital Subtraction Angiography
DVT	Deep Venous Thrombosis
ECG	Electro Cardiography
ELIZA	Enzyme Linked Immunosorbent Assay
FOV	Field Of View
GY	Gray
Hg	Mercury
HRCT	High Resolution Computed Tomography
HU	Hounsfield Unit
IV	Intra Venous
IVC	Inferior Vena Cava
ISSPE	Isolated Subsegmental Pulmonary Embolism
KV	Kilovoltage
MA	Milliampere
MAA	Macro Aggregate Of Albumin
MDCT	Multidetector Computed Tomography

MHCT	Multislice Helical Computed Tomography
MIP	Maximum Intensity Projection
MPR	Multiplanar Reconstruction
MPVR	Multiplanar Volume Reconstruction
MR	Magnetic Resonance
MRA	Magnetic Resonance Angiography
MRI	Magnetic Resonance Imaging
MSCT	Multislice Computed Tomography
PACS	Picture-Archiving and Communications System
PAH	Pulmonary Artery Hypertension
PCO ₂	Partial Pressure of Carbon Dioxide
PE	Pulmonary Embolism
PERF	Perfusion
PIOPED	Prospective Investigation Of Pulmonary Embolism Diagnosis
PO ₂	Partial Pressure of Oxygen
PTE	Pulmonary Thromboembolism
RLL	Right Lower Lobe
LLL	Left Lower Lobe
SCTA	Spiral Computed Tomography Angiography
SPECT	Single Photon Emission Computed Tomography
T _C	Technitium
TREAT	Time Resolved Echo Shared Angiography Technique
RV	Right Ventricle
SD	Standard Deviation
US	Ultra Sound
VENT	Ventilation
VQ	Ventilation/Perfusion
VTE	Venous Thromboembolism