AORTIC HOMOGRAFT

Thesis Submitted In Partial Fulfillment Of The MD Degree In Cardiothoracic Surgery

Ву

Ashraf Abdalla Elsebaie

5461

Supervisors

Prof. Dr.
Mohamed F. Bassiouny
Head of Cardiothoracic surgery Dept.
Faculty Of Medicine, Ain Shams
University

Prof. Dr.
Tarek Zaghloul
Professor of Cardiothoracic surgery
Faculty Of Medicine, Ain Shams
University

Prof. Dr. Eglal Abdel Razik
Professor of Pathology
Faculty Of Medicine,
Ain Shams University

7.412 . A

> Faculty Of Medicine Ain Shams University 1997







Acknowledgment

Words fail to express my supreme appreciation to professor Dr. Mohamed F. Bassiouny head of cardiothoracic surgery Dept. Ain Shams University who was quite a tutor, guide and instructor throughout my work. His attitude has been my inspiration during the preparation of this thesis.

I wish also to thank professor Dr. Eglal Abdel Razik Prof. of pathology at Ain shams University who has been always ready with help and advice throughout this work.

I would like also to express my sincere thanks and appreciation to Prof. Dr. Tarek Zaghloul Prof. of cardiothoracic surgery at Ain Shams University for his great help and suggestions in preparing this thesis

Finally, I would like to thank all members and stuff of our Dept. for their help and cooperation.



List Of Figures

rig. (1) I he interior of the left ventricle.	4
Fig. (2): Short-axis view from above of the aortic valve.	5
Fig. (3): Development of aortic valve.	6
Fig. (4): The extra length of aortic cusps.	7
Fig. (5): Contact between aortic cusps during systole and diastole.	8
Fig. (6): Structural relationships of the left ventricular outflow tract.	9
Fig. (7) : Cardiac skeleton.	10
Fig. (8): The ascending aorta.	11
Fig. (9): Relations of the right aortic sinus.	12
Fig. (10) : Relation of the posterior aortic sinus.	13
Fig. (11): Relation of the aorta to the left ventricle	14
Fig. (12): Relation of the left aortic sinus.	15
Fig. (13): Normal aortic valve viewed from above.	16
Fig. (14): A bicuspid aortic valve stenosis viewed from above.	17
Fig. (15): A tricuspid aortic valve stenosis viewed from above.	18
Fig. (16): A unicuspid aortic valve stenosis viewed from above.	18
Fig. (17): A quadricuspid aortic valve stenosis viewed from above.	19
Fig. (18): Rheumatic aortic valve show commissural fusion and	
leaflet fibrosis.	20
Fig. (19): Rheumatic aortic valve stenosis showing	
calcification at commissural sites.	20
Fig. (20): Calcific aortic stenosis in bicuspid aortic valve.	21
Fig. (21) : Calcific aortic stenosis in tricuspid aortic valve.	22
Fig. (22): Posteroanterior view X-ray of the chest in a	
62-year-old male with calcific aortic stenosis shows a	
normal-sized heart with a slightly enlarged ascending aorta.	24
Fig. (23): A preoperative ECG of an 32-years old male	
with aortic stenosis showing left ventricular hypertrophy.	25
Fig. (24): Parasternal long-axis view echocardiography	
in systole show calcific aortic stenosis.	26
Fig. (25): Sever aortic stenosis is evaluated from the	
apical approach using a non-imaging, continuous- wave	
Doppler device.	26
Fig. (26): frontal-View angiogram of a patient with	
valvar aortic stenosis.	27
Fig. (27): Infective endocarditis of the aortic valve cusps.	28
Fig. (28): Aortic root dilation in a patient with Marfan's	
disease causes aortic regurgitation.	29

Fig. (29): Barium Opacifies the esophagus on the chest	
films of a patient with aortic regurgitation.	33
Fig. (30): Continuous-wave Doppler recording from the	
apex toward the aortic valve shows the high velocity	
diastolic flow jet due to aortic regurgitation.	34
Fig. (31): Vertical laminar airflow cabinet.	47
Fig. (32): Harvested aortic conduit and pulmonary valve conduit.	50
Fig. (33): Control freezing pouch with temperature probe	
placed through a watertight portal.	65
Fig. (34): Parasternal long axis two-dimensional	
echocardiographic view for allograft sizing.	79
Fig. (35): Trimming of aortic homograft valve pillars.	81
Fig. (36): Aortic homograft with non-coronary sinus intact.	81
Fig. (37): Reverse "S" aortotomy for freehand aortic	
valve replacement with allograft aortic valve.	83
Fig. (38): Three sutures are placed for the proximal suture	
line so as to "line up" the allograft coronary sinuses	
to the recipient coronary ostia.	84
Fig. (39): Continuos proximal suture line for free-hand	
insertion of aortic homograft.	85
Fig. (40): Allograft trimming below the coronary ostia.	86
Fig. (41): The distal suture line for homograft insertion.	86
Fig. (42-A): Trimming of homograft non-coronary	
sinus in flange technique.	87
Fig. (42-B): The distal suture line is continued to the	
points where the aortotomy closure sutures meet them.	88
Fig. (43): In classic technique all three sinuses of	
the homograft are excised.	88
Fig. (44): The aortotomy suture incorporates and	
approximates the intact noncoronary sinus.	90
Fig. (45): The aortotomy is closed and homograft is in position.	90
Fig. (46): interrupted proximal suture line technique for	
freehand aortic valve replacements.	92
Fig. (47): Aortotomy for annulus enlargement when	
using the Manouguian approach.	94
Fig. (48): Pledgetted technique in the proximal suture line used	
for mitral to mitral valve tissue closure (Manouguian Technique)	94
Fig (49): Aortic Annulus enlargement.	95
Fig. (50): Closure of the aortic root following	
freehand aortic valve insertion.	97

Fig. (51) :Aortic root geometry and the size of the	
noncoronary sinus are maintained using the classic technique	
plus augmentation using Dacron patch for a small aortic root.	98
Fig. (52): Placement of sutures to manage coronary ostia	
approaching 180° orientation.	100
Fig. (53): Allograft intra-aortic cylinder with side to side	
coronary artery anastomoses.	101
Fig. (54): If coronary ostia are made to approximate one another	
and are too high, the leaflet may be deformed and may prolapse.	102
Fig. (55): The aorta has been transected and the aortic buttons	
around the coronary ostia have been created.	106
Fig (56): Interrupted proximal suture line technique in	
aortic root replacement.	107
Fig. (57): Suturing of the coronary buttons.	108
Fig. (58): Distal suture line of an allograft aortic root	
replacement using Teflon felt strip.	109
Fig. (59): Aortic root replacement with homograft and the aortic	
root abscess has been obliterated.	110
Fig. (60): Use of anterior mitral leaflet of the homograft to repair	
the aortic sub-annular defect in cases of aortic root abscess.	111
Fig. (61): The septal defect is filled with the entire anterior leaflet	
of the mitral valve.	114
Fig. (62): trimming of aortic homograft before insertion on the	
operating room.	121
Fig. (63): Aortic homograft in place after compilation of proximal	
suture line.	123
Fig. (64): A pie-chart showing percentage of Egypt and	
Houston patients.	128
Fig. (65): A pie-chart showing percentage of males and females.	129
Fig. (66): A bar showing age and sex distribution in relation	
to years.	129
Fig. (67): A pie-chart showing the preoperative percentage of	
NYHA function class.	130
Fig. (68): A preoperative ECG of an 19 old male (No. 11) with	
aortic stenosis showing left ventricular hypertrophy and strain pattern.	131
Fig. (69): Chest X-ray Shows left ventricular hypertrophy.	131
Fig. (70): A pie-chart showing the etiology of aortic valve disease	
in our study.	132

Fig. (71): A pie-chart showing the pathology of aortic valve	
disease in our patient.	133
Fig. (72): preoperative transthoracic 2-D Echo. in patient	124
No. 15. Showing thickened aortic valve leastlet. Fig. (73): Preoperative transthoracic Doppler in patient	134
No. 15 Showing sever aortic regurgitation	134
Fig. (74): A pie-chart showing percentage of cryopreserved	
and antibiotic sterilized homografts which were used in our patients.	136
Fig. (75): A pie-chart showing number, percentage, and	
cause of discarded antibiotic sterilized homografts.	137
Fig. (76): A pie-chart showing number, percentage and type	120
of the surgical techniques that were done in our patient. Fig.(77) Shows aortic valve area by preoperative ECHO. in	139
relation to aortic ring diameter measured intraoperative in mm.	140
Fig. (78): A chart showing number, percentage and	110
results of transesophageal echocardiography.	142
Fig (79): A chart Showing the comparison between the mean of	
LVEDD and LVESD in the preoperative and postoperative period.	145
Fig. (80): A chart Comparing the mean of fractional shortening	140
and ejection fraction during preoperative and postoperative periods. Fig. (81) shows the preoperative EF% compared to postoperative	145
EF% in survival.	146
Fig. (82): A chart Showing the number and percentage	
of survival in both group and the number and percentage	
of the mortality on each group.	148

List Of Tables

	Page
- Table 1 Showing different protocols for aortic homograft	
sterilization.	55
- Table 2 Showing calcification of patients according to	
etiology, pathology and associated lesions.	133
- Table 3 Showing Preoperative echocardiography finding.	
LVEDD = Lt. Ventricular end diastolic dimension.	
LVESD = Lt. Ventricular end systolic dimension.	
EF = ejection fraction. FS = fraction shortening.	135
- Table 4 Showing Preoperative echocardiography aortic	
diameters.	135
- Table 5 Showing characteristics of antibiotic sterilized	
homografts.	137
- Table 6 shows aortic valve area by preoperative echo in relation	
to aortic ring diameter measured intraoperatively in mm.	140
Table 7 Showing the echocardiographic data of the postoperative	
patients in the 1 st postoperative, after 4 weeks and after 6 months.	
LVEDD = left ventricular end diastolic diameter. LVESD = left	
ventricular end systolic diameter. FS = fractional shortening.	
EF = ejection fraction.	144
- Table (8) shows the preoperative EF% compared to	
postoperative EF% in survival.	146
- Table (9) Showing all patients data pre & postoperative.	149



List of Abbreviations

- RCA = right coronary artery
- -A = anterior
- AATB = American Association of Tissue Banks
- AI = Aortic insufficiency
- AIDS = acquired immune deficiency disease
- -AO = aorta
- ARC = AIDS-related complex
- -AV = Aortic valve
- CJD = Creutzfeldt-Jakob disease
- DMS0 = dimethyl sulfoxide
- EF = Ejection fraction
- FCS = fetal calf serum
- FDA = United States Food and Drug Administration
- Fig. = Figure
- FS = Fraction shortening
- L = left,
- LC = Left coronary ostia
- LCA = left coronary artery
- LV = left ventricle
- LV = Left ventricle
- LVEDD = Left ventricular end diastolic dimension
- LVEDP = Left ventricular end diastolic pressure
- LVESD = Left ventricular end systolic dimension
- mg = milligram
- ml = millilitre

- mm = millimetre
- MS = Membranous septum
- MV = Mitral valve

- MS = Membranous septum

- NYHA = New-York Heart Association
- -P = posterior
- PPF = plasma protein fraction
- -R = right- RA = Right atrium
- RC = Right coronary ostia
- RV = Right ventricle
- RV = Right ventricle
- SD = Standard deviation - ST = tricuspid valve
- VS = ventricular septum
- VTB = Virginia Tissue Bank

Contents

** Introduction and aim of the work	l
** Anatomy of aortic root and valve	4
** The pathophysiology of aortic valve diseases 1	6
** Historical background	35
** Homograft valve banking	40
** Effects of preimplantation processing on homograft morphology 7	10
** Left ventricular outflow tract reconstructions	
using aortic homograft	77
** Material and methods1	16
** Results 12	28
** Discussion 15	1
** Summary and conclusion 10	59
** Reference 17	71
** Arabic summary	

