

STUDIES ON THE PRODUCTION AND STABILITY
OF BACILLUS THURINGIENSIS ENDOTOXIN

By

MAGDA ABD EL-GHAFFAR EL-BENDARY

B.Sc. (MICROBIOLOGY)

AIN-SHAMS UNIVERSITY

(1988)

A THESIS

SUBMITTED IN PARTIAL FULFILLMENT

OF THE REQUIREMENTS FOR

THE DEGREE OF

MASTER OF SCIENCE

IN

(MICROBIOLOGY)

UNDER SUPERVISION OF

PROF. DR. MOHAMED ABU-SHADY

PROF. OF MICROBIOLOGY
AND VICE DEAN OF
THE FACULTY OF SCIENCE
FACULTY OF SCIENCE
AIN SHMS UNIVERSITY

PROF. DR. MOHAMED SALAH FODA

PROF. OF MICROBIAL CHEMISTRY
AND VICE DEAN OF
BASIC SCIENCE DIVISION
NATIONAL RESEARCH CENTRE

DEPARTMENT OF MICROBIOLOGY

FACULTY OF SCIENCE

AIN SHAMS UNIVERSITY

1994

5849
M.A



بِسْمِ اللَّهِ الرَّحْمَنِ الرَّحِيمِ

وَالْوَفِيَّ الْإِبْرَاهِيمَ

حَدَّثَنَا أَبُو بَكْرِ بْنُ أَبِي شَيْبَةَ



ACKNOWLEDGEMENT

The author would like to express her thanks and gratitude to **Dr. Mohamed Abu-Shady**, Professor of Microbiology, Department of Microbiology and Vice Dean of the Faculty of Science, Ain-Shams University for his kindly supervision, encouragement and support and continuous advice.

The author sincerely presents her gratefulness and deepest feelings to **Dr. Mohamed Salah Foda**, Professor of Microbiology, Department of Microbial Chemistry, National Research Centre, Cairo, Egypt, for suggesting the point and for his instructive discussions, supervision, continuous encouragement, valuable advice and sincere guidance throughout this work.

The author is also grateful to **Dr. H.S. Salama**, and **Dr. Aziza El-Sharaby**, Professors of Entomology , Plant Protection Department, National Research Centre for their kindly supervision, encouragement and continuous advice.

The Co-operation and help of my colleagues and all members of Microbial Chemistry Department are gratefully acknowledged.

The thanks are also due to the National Research Centre for the facilities that enabled me to accomplish this work.

APPROVAL SHEET

NAME: MAGDA ABD EL-GHAFFAR EL-BENDARY

TITLE: *STUDIES ON THE PRODUCTION AND STABILITY OF
BACILLUS THURINGIENSIS ENDOTOXIN.*

SCIENTIFIC DEGREE : M.SC.

This thesis has been approved by supervisors:

Prof. Dr. **Mohamed Abu shady**

prof of microbiology and vice
dean of the faculty of science,
Ain -Shams University



Prof. Dr. **Mohamed Salah Fouda**

prof of microbiology and vice dean
of the basic science division,
National Research Centre.

Date of Research : 9/4/1991

CONTENTS

	Page
PREFACE.....	I
1. REVIEW OF LITERATURE.....	3
1.1. Discovery and History of Entomo-Pathogenicity of <u>Bacillus thuringiensis</u> (<u>B.t.</u>).....	3
1.2. Cell Morphology and Classification of <u>B.t.</u>	4
1.3. Toxins Produced by <u>B.t.</u>	5
1.4. Nature of the Parasporal Crystalline- δ -Endotoxin....	7
1.5. Role of Spores in Pathogenicity of <u>B.t.</u> Against Insects.....	10
1.6. The Combined Action of Spores and Crystals in <u>B.t.</u> Pathogenicity.....	14
1.7. Mode of Action of- δ -Endotoxin of <u>B.t.</u> in Susceptible Insects.....	15
1.8. Physiology of <u>B.t.</u> Growth and- δ -Endotoxin Production	26
1.8.1. Metabolic Activities and Growth Kinetics.....	26
1.8.2. Conditions for Growth, Sporulation and Crystal Formation.....	27
1.8.2.1. Effect of pH, Carbon and Nitrogen Sources.....	27
1.8.2.2. Utilization of Complex Media and Agro-Industrial Byproducts for <u>B.t.</u> Production.....	34
1.8.2.3. Effect of Aeration.....	40

	Page
1.9. Factors Affecting Stability and Potency of <u>B.t.</u>	
δ -Endotoxin.....	42
1.9.1. Effect of Temperature.....	43
1.9.2. Effect of Radiations.....	47
1.9.3. Enhancement of <u>B.t.</u> Potency by Chemical Additives.....	50
 2. MATERIALS AND METHODS.....	 54
2.1. Materials.....	54
2.1.1. Organisms Used and Sources of Cultures.....	54
2.1.2. Media.....	55
2.1.3. Chemicals and Reagents.....	61
2.2. Methods.....	64
2.2.1. Growth Conditions and Physiology of <u>B.t.</u>	64
2.2.2. Staining Procedure of Crystalline- δ -Endotoxin of <u>B.t.</u>	69
2.2.3. Light Microscopy, Electron Microscopy and Histological Procedures.....	70
2.2.4. Isolation of Mutants of <u>B.t.</u> Resistant to Physical Factors.....	74
2.2.5. Recovery of the Spore- δ -Endotoxin Complex of <u>B.t.</u> var. <u>aizawai</u> HD-282 for Bioassay Purposes	78
2.2.6. Insect Culture Breeding.....	79
2.2.7. Bioassay of Spore- δ -Endotoxin Complex of <u>B.t.</u> var. <u>aizawai</u> HD-282.....	80
2.2.8. Studies on Increasing Potency of Spore- δ -Endotoxin Complex by Chemical Additives.....	81

	Page
3. RESULTS AND DISCUSSION.....	84
3.1. Physiology of <u>B.t.</u>	84
3.1.1. Establishment of Growth Curves of <u>B.t.</u> Cul- tures Understudy.....	86
3.1.2. Effect of Initial pH and Medium Buffering on Growth pattern and Sporulation Yield.....	89
3.1.3. Effect of Medium Supplementation with Diffe- rent Carbon Sources on Growth, CFU, and Spor- ulation Yields of <u>B.t.</u> var. <u>aizawai</u> HD-282.....	102
3.1.4. Effect of Medium Supplementation with Amino Acids on Growth, Viable Counts and Sporulation Titers of <u>B.t.</u> var. <u>aizawai</u> HD-282.....	107
3.1.5. Effect of L-aspartic Acid and L-cysteine con- centration on Growth, CFU and Sporulation Yields of <u>B.t.</u> var. <u>aizawai</u> HD-282.....	111
3.1.6. Effect of Supplementation of NBM with Yiest Extract on Growth and Sporulation Yields of <u>B.t.</u> var. <u>aizawai</u> HD-282.....	112
3.1.7. Effect of Extent of Aeration on Growth, CFU, and Sporulation Titer of <u>B.t.</u> var. <u>aizawai</u> HD-282.....	117
3.1.8. Effect of Inoculum Size on Growth, Viable and Spore Counts of <u>B.t.</u> var. <u>aizawai</u> HD-282.....	125
3.2. Potentials of Local Agroindustrial Byproducts and Plant Proteins as Media for Production of <u>B.t.</u> Spore- -Endotoxin Complex.....	128

	Page
3.2.1. Evaluation of Agroindustrial Byproduct for <u>B.t.</u> Production.....	128
3.2.2. Utilization of Legumes as Major Medium Components for <u>B.t.</u> Production.....	129
3.2.3. Development of Mono Component Media Based on Local Byproducts.....	132
3.3. Microscopic Studies on Growth, Sporulation and Endotoxin Formation.....	145
3.3.1. Morphological Changes During Growth, Sporogenesis and Crystalline- δ -Endotoxin Formation in <u>B.t.</u> var. <u>aizawai</u> HD-282.....	145
3.3.2. Observation of Sporogenesis and Crystalline- δ -Endotoxin Formation by Transmission Electron Microscopy.....	153
3.4. Assessment of Mode of Action of Crystalline- δ -Endotoxin and Histological Changes Induced by <u>B.t.</u> var. <u>aizawai</u> HD-282 in larvae of <u>Spodoptera littoralis</u>	165
3.4.1. Symptoms of Infection.....	165
3.4.2. Histopathological Effects of <u>B.t.</u> var. <u>aizawai</u> HD-282 in <u>Spodoptera littoralis</u>	168
3.5. Isolation of Mutants Resistant to Physical Factors... ..	188
3.5.1. Isolation and Characterization of Ultraviolet Resistant Mutants.....	189
3.5.2. Isolation and Characterization of Heat-Resistant Mutants.....	208
3.6. Potency Increase of δ -Endotoxin by chemical Additives	222

	Page
4. GENERAL DISCUSSION.....	233
5. SUMMARY.....	267
6. REFERENCES.....	287
7. ARABIC SUMMARY.....	316

List of Tables

	Page
Table (1) Screening of some strains of <u>B.t.</u> against <u>Sp. littoralis</u> larvae.	88
Table (2) Establishment of growth parameters of the three selected <u>B.t.</u> strains.	90
Table (3) Effect of initial pH of phosphate buffered NBM on growth, CFU and sporulation yields of <u>B.t.</u> var. <u>entomocidus</u> HD-635 grown in shake culture.	96
Table (4) Effect of initial pH of phosphate buffered NBM on growth, CFU and sporulation yields of <u>B.t.</u> var. <u>aizawai</u> HD-134 grown in shake culture.	98
Table (5) Effect of initial pH of phosphate buffered NBM on growth, CFU and sporulation yields of <u>B.t.</u> var. <u>aizawai</u> HD-282 grown in shake culture.	100
Table (6) Effect of supplementation of NBM with different carbon sources on growth, viable count and sporulation yield of <u>B.t.</u> var. <u>aizawai</u> HD-282.	103
Table (7) Effect of concentration of glucose and wheat flour in NBM on growth, viable count and sporulation yield of <u>B.t.</u> var. <u>aizawai</u> HD-282.	105
Table (8) Effect of supplementation of NBM with amino acids on growth, CFU and sporulation titer of <u>B.t.</u> var. <u>aizawai</u> HD-282.	108

	Page
Table (9) Effect of L-aspartic acid concentration on growth parameters of <u>B.t.</u> var. <u>aizawai</u> HD-282.	113
Table (10) Effect of cysteine-Hcl concentration on growth parameters of <u>B.t.</u> var. <u>aizawai</u> HD-282.	115
Table (11) Effect of supplementation of NBM with yeast extract on growth, viable count and sporulation yield of <u>B.t.</u> var. <u>aizawai</u> HD-282.	118
Table (12) Effect of aeration level on growth parameters and crystalline- δ -endotoxin formation of <u>B.t.</u> var. <u>aizawai</u> HD-282 under shake incubation conditions.	121
Table (13) Effect of aeration level on growth parameters and crystalline- δ -endotoxin formation under static conditions.	123
Table (14) Effect of inoculum size on growth, viable count and sporulation titer of <u>B.t.</u> var. <u>aizawai</u> HD-282 in NBM.	126
Table (15) Growth, sporulation and crystalline- δ -endotoxin formation of <u>B.t.</u> var. <u>aizawai</u> HD-282 grown in BM supplemented with some Major byproducts of agricultural industries in Egypt.	130
Table (16) Growth, sporulation and crystalline- δ -endotoxin formation of <u>B.t.</u> var. <u>aizawai</u> HD-282 grown in BM supplemented with various leguminous seeds as the major nitrogen source in the growth medium.	131

	Page
Table (17) Effect of fodder yeast concentration on growth parameters of <u>B.t.</u> var. <u>aizawai</u> HD-282.	135
Table (18) Effect of offals meal concentration on growth parameters of <u>B.t.</u> var. <u>aizawai</u> HD-282.	137
Table (19) Effect of feather meal concentration on growth parameters of <u>B.t.</u> var. <u>aizawai</u> HD-282.	140
Table (20) Effect of soybean concentration on growth parameters of <u>B.t.</u> var. <u>aizawai</u> HD-282.	143
Table (21) UV-inactivation of vegetative cells (five hours grown culture) of <u>B.t.</u> var. <u>aizawai</u> HD-282.	191
Table (22) UV-inactivation of spores of <u>B.t.</u> var. <u>aizawai</u> HD-282.	194
Table (23) Comparative survivors of UV-resistant mutants derived from <u>B.t.</u> var. <u>aizawai</u> HD-282.	198
Table (24) Growth patterns, viable count, sporulation titer and ability of crystalline- δ -endotoxin formation of some selected UV-resistant mutants derived from <u>B.t.</u> var. <u>aizawai</u> HD-282.	200
Table (25) Comparison between the wild type strain (<u>B.t.</u> var. <u>aizawai</u> HD-282) and six selected UV-resistant mutants, (with different abilities to synthesize crystals) with respect to UV-radiation tolerance.	204

	Page
Table (26) Comparative evaluation of the potencies of UV-resistant mutants against the target insect species <u>S . littoralis</u> .	209
Table (27) Survivors after heating treatments of a full grown culture of <u>B.t.</u> var. <u>aizawai</u> HD-282.	211
Table (28) Comparative survivors counts of heat-resistant mutants derived from <u>B.t.</u> var. <u>aizawai</u> HD-282.	213
Table (29) Growth patterns, viable counts, sporulation titers and crystalline- δ -endotoxin formation of some selected heat-resistant mutants derived from <u>B.t.</u> var. <u>aizawai</u> HD-282.	216
Table (30) Comparative UV-resistance patterns of some selected heat resistant mutants of <u>B.t.</u> var. <u>aizawai</u> HD-282 as compared to their wild type strain.	220
Table (31) Comparative evaluation of the potencies of heat-resistant mutants against larva of <u>S . littoralis</u> .	223
Table (32) Effect of lipid emulsifying agents on the potency of <u>B.t.</u> var. <u>aizawai</u> HD-282 VS. <u>S . littoralis</u> larvae.	226
Table (33) Effect of protein solubilizing agents on the potency of <u>B.t.</u> var. <u>aizawai</u> HD-282 VS. <u>S . littoralis</u> larvae.	228
Table (34) Effect of amino acids on the activity and potency of <u>B.t.</u> var. <u>aizawai</u> HD-282 VS. <u>S . littoralis</u> larvae.	230

LIST OF FIGURES

	Page
Fig. (1) Changes in optical density (OD 600 nm) and pH of the three selected <u>B.t.</u> strains during 4 days incubation period in NBM.	92
Fig. (2) Viable counts (CFU) of the three selected <u>B.t.</u> strains during 4 days incubation period in NBM.	93
Fig. (3) Sporulation titers of the three selected <u>B.t.</u> strains during 4 days incubation period in NBM.	94
Fig. (4) Effect of initial pH of phosphate buffered NBM on growth pattern and sporulation yield of <u>B.t.</u> var. <u>entomocidus</u> HD-635 grown in shake cultures.	97
Fig. (5) Effect of initial pH of phosphate buffered NBM on growth and sporulation yield of <u>B.t.</u> var. <u>aizawai</u> HD-134 grown in shake cultures.	99
Fig. (6) Effect of initial pH of phosphate buffered NBM on growth and sporulation yield of <u>B.t.</u> var. <u>aizawai</u> HD-282 grown in shake cultures.	101
Fig. (7) Effect of concentration of glucose and wheat flour on viable count and sporulation yield of <u>B.t.</u> var. <u>aizawai</u> HD-282.	106
Fig. (8) Effect of NBM supplementation with amino acids on viable count and sporulation titer of <u>B.t.</u> var. <u>aizawai</u> HD-282.	110