

**DIAGNOSTIC AND THERAPEUTIC  
USES OF LAPAROSCOPY  
WITH EVALUATION OF LAPAROSCOPIC  
CHOLECYSTECTOMY IN AIN SHAMS  
UNIVERSITY HOSPITALS**

**THESIS**

Submitted For the partial  
Fulfillment of M.S Degree in **general Surgery**

617.556  
K.R. M

By

**Khaled Mohamed Sayed Ahmed**  
MB.B,Ch.

Supervised By

**Prof. Dr. Abou Bakr El-Sedik Moustafa**

*Professor of general surgery  
Faculty of medicine  
Ain Shams university*

**Dr. Gamal Abdel-Rahman El-Mowalled**

*Lecturer of general surgery  
Faculty of medicine  
Ain Shams university*


Faculty of medicine  
Ain Shams university

\*\*\* 1996 \*\*\*












## **ACKNOWLEDGMENT**

I would like to express my deepest gratitude and my heartfelt admiration for **Prof. Dr. Abou Bakr El-Sedik Moustafa** , Professor of General Surgery, Ain Shams University, whose fruitful suggestions and honest criticism were extremely important factors in the accomplishment of this thesis.

I would also like to thank intensely, **Dr. Gamal Abdel-Rahman El-Mowalled**, Lecturer of General Surgery, Ain Shams University, for his immense effort and advice and for the guidance that he gave me throughout this work.

*Khaled Mohamed Sayed Ahmed*



## **CONTENTS**

	Page
Introduction	1
Aim of the work.	2
Review of literature.	
History of laparoscopic surgery	3
History of laparoscopic cholecystectomy	8
Laparoscopic Equipment and Instrumentation	9
Technique of laparoscopy	33
Diagnostic indications of laparoscopy	42
Therapeutic indications of laparoscopy	62
Contraindications of laparoscopy	139
Complications of laparoscopy	140
Management of chronic calcular cholecystitis	149
Materials and Method	159
Subjects	165
Assignment	169
Evaluation	175
Conclusion	176
References	177
Arabic summary	





## **LIST OF FIGURES**

	page
<b>Fig. (1):</b> The verres needle	10
<b>Fig. (2):</b> Hasson Cannula	10
<b>Fig. (3):</b> Modern high-flow electronic pressure controlled insufflator for laparoscopic surgery	11
<b>Fig. (4):</b> Reusable Cannula	13
<b>Fig. (5):</b> Disposable Cannulae	14
<b>Fig. (6):</b> There are multiple heads available for the unipolar cautery devices: hook, needle, spatula and blunt	15
<b>Fig. (7):</b> Unipolar scissors and forceps	15
<b>Fig. (8):</b> Aquapurator	17
<b>Fig. (9):</b> The laparoscope consists of an eyepiece and a light source	18
<b>Fig. (10):</b> A typical multilevel video cart	22
<b>Fig. (11):</b> Commonly used laparoscopic instruments. (a) top to bottom, 10 mm trocar and 5 mm trocar, reducer tube, suture applicator, veress needle (b) 10 and 5 mm telescopes (c) curved grasping forceps (d) atraumatic grasping forceps (e) heavy duty traumatic grasping forceps	25
<b>(12):</b> (f) Clip applicators: (left) non-disposable for metal clips, (right) non-disposable for absorbable clips (g) Berci dissecting spatula (insulated) and electro-surgical hook knife (insulated) (h) suction/irrigation cannula (i) cholangiographic cannula	26
<b>(13):</b> The Endobag is a small plastic bag inserted through a cannula with a special inserter	29

<b>Fig. (14):</b>	(a) Injected saline cannot be aspirated from the peritoneal space, but (b) will be aspirated if injected into preperitoneal space, (c) a droplet of saline is aspirated into the peritoneal cavity, but not the preperitoneal space, (d) the tip should "fan" freely if in the peritoneal cavity	36
<b>Fig. (15):</b>	Two techniques of inserting the initial trocar. (a) after obtaining a pneumoperitoneum lifting the hypogastric abdominal wall gives counter traction against the considerable force necessary to insert the trocar. (b) an alternative method is to apply epigastric compression this elevates the lower abdomen, gives counter pressure, and creates a much safer angle of penetration	38
<b>Fig. (16):</b>	Transillumination of the abdominal wall reveals a blood vessels to be avoided during cannula insertion	39
<b>Fig. (17):</b>	The Hasson cannula is held to the fascia with sutures to prevent air loss. It is placed through a minifascial and peritoneal incision under direct visualization rather than abland puncture	40
<b>Fig. (18):</b>	Normal, pale-blue gallbladder in the typical location at the hepatic incisura, projecting somewhat beyond the edge of the liver	
<b>Fig. (19):</b>	Chronic cholecystitis with whitish, thickened wall, increase vascularity, and adhesions to the omentum	
<b>Fig. (20):</b>	Cirrhosis with micro to medium sized nodules with varying inflammatory activity	
<b>Fig. (21):</b>	Hydatid cyst on the surface of the right lobe	
<b>Fig. (22):</b>	Hepatocellular carcinoma in underlying liver cirrhosis of medium sized nodularity with typical whitish-red coloration and abnormal vascularity	
<b>Fig. (23):</b>	(a) Lines of traction are shown during laparoscopic cholecystectomy, (b) lateral and inferior traction of Harman's pouch	68

<b>Fig. (24):</b> The peritoneum over the anterior surface of Calot's triangle is opened, starting over Hartmann's pouch and staying close to the gall bladder	69
<b>Fig. (25):</b> Turning Hartmann's pouch medially permits posterolateral dissection of the serosa of the gall bladder	69
<b>Fig. (26):</b> Separating the neck of the gall bladder from the hepatic bed permits clear visualization of the junction of the cystic duct and infundibulum of the gallbladder in all of its circumference	70
<b>Fig. (27):</b> The cholangiogram catheter secured in place by a partially closed titanium clip	70
<b>Fig. (28):</b> Dissection of the gallbladder from the liver bed using hook diathermy	71
<b>Fig. (29):</b> Bleeding from a trocal side is controlled by two sutures placed with a Keith needle in the path of the vessel at both sides of the trocar	83
<b>Fig. (30):</b> Placement of trocars for laparoscopic treatment of non parasitic liver cysts	95
<b>Fig. (31):</b> (a top left) The appendix is grasped by the first assistant while traction of the cecum cephalad maintains exposure. (b top center) a very inflamed and rigid or friable appendix may be handled more easily by encircling it with a pre-tied chromic ligature. The first assistant grasps the tail of the ligature rather than the appendix. (c top right) the mesentery is serially clipped and divided by the surgeon. (d bottom left) two pre-tied chromic ligatures are placed at the base of the appendix where it merges with the cecum, and a third ligature is placed approximately 1 cm distally. (e bottom center) alternatively, an endoscopic stapler is placed across the base of	

	the appendix after the mesentery has been divided. (f bottom right) a retrocecal appendix may be exposed by incising the white line and rolling the cecum cephalad and medially	100
<b>Fig. (32):</b>	Trocar placement	103
<b>Fig. (33):</b>	Transabdominal preperitoneal approach. (a) peritoneal incision; (b) peritoneal flap and positioning of the mesh; (c) closure of the peritoneum using continuous sutures	105
<b>Fig. (34):</b>	(a) Small indirect hernia; (b) intraperitoneal onlay mesh	107
<b>Fig. (35):</b>	A balloon dissection device	108
<b>Fig. (36):</b>	(a) The dissector is advanced within the rectus sheath to reach the pubic tubercle. (b) the balloon is filled with 600ml of normal saline to create an extraperitoneal cavity. (c,d) after removal of the balloon, the cannula is advanced into cavity and connected to Co <sub>2</sub> . The cavity is inflated to a pressure of 8-10 mmHg	109
<b>Fig. (37):</b>	Closure of the ring for type II hernias (small indirect hernias in the young adult). A: Cooper's ligament; B: epigastric vessels; C: Sutures on the internal ring; D: spermatic vessels; E: vas deferens; F: iliac vessels	110
<b>Fig. (38):</b>	Positioning of ports	111
<b>Fig. (39):</b>	Abdominal port placement	112
<b>Fig. (40):</b>	The fan retractor elevate the left lobe of the liver. The hepatogastric ligament is sharply dissected for mobilization of gastroesophageal junction	118
<b>Fig. (41):</b>	The gastroesophageal junction is grasped at point • and pulled in direction A: for dissection of the left crus and direction B: for dissection of the right crus	119

<b>Fig. (42):</b>	A tape pulling the oesophagus forward	120
<b>Fig. (43):</b>	The anterior wall of the fundus, at approximately the fundus / body junction is picked up and passed to the forceps behind the oesophagus	121
<b>Fig. (44):</b>	The wrap is complete and secured with two interrupted sutures	122
<b>Fig. (45):</b>	Port placement sites with the two operative fields requiring consideration	123
<b>Fig. (46):</b>	Completed mobilization of the distal oesophagus with preservation of the hepatic branches of the anterior vagus	124
<b>Fig. (47):</b>	(a) The longitudinal representation of the blood supply within the falciform ligament and the plane of dissection used to preserve the blood supply to the round ligament. (b) the round ligament is detached from the umbilicus after ligation (c) completed mobilization with the tail of the extracorporeal ligature left long to act as a "tow rope"	125
<b>Fig. (48):</b>	Removal of the nasogastric tube and oesophageal mobilization with delivery of the "tow rope" behind the oesophagus from right to left	126
<b>Fig. (49):</b>	Downward traction on the sling is manometrically controlled to increase the distal sphincter pressure to 15-20 mmHg. The initial non-absorbable anchoring mattress suture is placed	127
<b>Fig. (50):</b>	Completed fixation with multiple interrupted sutures.	127
<b>Fig. (51):</b>	Sites of insertion of ports. 1 and 2: 10-11 mm ports; 3,4 and 5: 5 mm ports	128
<b>Fig. (52):</b>	Exposure of the lower oesophagus and cardio-oesophageal junction. A gentle traction on the anterior gastric wall allows the thoracic oesophagus to descent for some distance into the abdomen. The myotomy is performed using the electro-surgical hook	129

<b>Fig. (53):</b>	X-scope sites      Y-10/11 mm      O-optional Trocscar site placement	132
<b>Fig. (54):</b>	Schematic drawing of anterior wall of stomach. Two posterior T-Fasteners have been placed into the stomach to secure the wall. A long spinal needle is being inserted for introduction of a J- wire to be followed by subsequent dilatation of the gastrostomy	135
<b>Fig. (55):</b>	The schematic drawing. Four T-Fasteners shown with gastrostomy tube being introduced over a J- wire into the stomach	136
<b>Fig. (56):</b>	The laparoscope is inserted through the cannula at the pararectal line left to the umbilicus. The other three or four cannulae are placed at the epigastric area, lateral eighth intercostal space, midclavicular line of the left subcostal space, and midaxillar line of the left subcostal space	137