



BIOSTRATIGRAPHIC STUDY OF SOME CRETACEOUS EXPOSURES IN NORTHERN SINAI, EGYPT.


BY MOHAMED MOHAMED ABDOU EL SAADANY (B.Sc. IN GEOLOGY, 1986)

A THESIS SUBMITTED IN PARTIAL FULFILMENT OF THE REQUIRMENTS OF THE M.Sc. DEGREE IN GEOLOGY

TO
THE GEOLOGY DEPARTMENT
FACULTY OF SCIENCE
AIN SHAMS UNIVERSITY

1995





To the Teacher of Mankind, to the trustworthy

Leader Mohamed, the Prophet of Allah, God's blessing

and peace be upon Him.

O God! Consider this work as part of my good deeds on Judgement Day.

Amen

Mohamed El Saadany

NOTE

esis is submitted to the Faculty of Science, Ain Shams University in partial fulfilment of uirements of the Master's Degree of Science in Geology. In addition to the research naterialised in this dissertation, the candidate has attended and successfully passed ration in the following M. Sc. courses:

ad geology and mapping
statistics and basic language
ropaleontology
soecology
tostratigraphy
stratigraphy
vanced structural geology
sotectonics

, the student has also passed the final examination of a course in the English Language

Prof. Wagih Abdelmalek.
Head of Geology Department

LIST OF CONTENTS

Մևոր	oter	Page
No.		No.
	LIST OF FIGURES	IV
	ACKNWLEDGEMENT	VIII
	ABSTRACT	Х
I-	INTRODUCTION	1
	1- Generalites.	1
	2- Methodology.	3
IJ-	THE PREVIOUS WORK	4
	1- Generalites.	4
	2- Previous Work	4
III-	LITHOSTRATIGRAPHY	17
	1- The Malha Formation.	17
	2- The Risan Aneiza Formation.	31
	3- The Halal Formation.	34
	4- The Abu Qada Formation.	39
	5- The Wata Formation.	42
	6- Sudr Formation (Sudr Chalk).	43
ĮV-	DIOSTRATIGRAPHY	49
	1- Gabal Manzour.	. 49
	a) Ammobaculites spp./ Haplophragmoides globasa.	54
	b) Barren Interzone (1st).	54
	c) Orbitolina cf. discoidea.	. 55
	d) Barren Interzone (2nd).	55
	e) Flabellammina spp.	55
	f) Thomasinella spp.	56
	g) Barren Interzone (3rd).	56
	h) Arenaceous foraminifera.	57
	i) Barren Interzone (4th).	_ 57
	j) Orbitolina cf. conçava.	57
	k) Barren Interzone (5th),	58
	2- Gabal El Minsherah (El Minsherah ridge).	58
	a) Barren luterzoue (1st).	58
	b) Flabellanımına spp.	59
	c) Barren Interzone (2nd).	59
	d) Thomasinella spp.	6
	e) Barren luterzone (3rd).	(
	f) Heterohelix spp. / Hedbergella spp.	(

t	er	Page
		<u>No.</u>
	g) Barren Interzone (4th).	61
	h) Discorbis turonicus / D. minutus.	62
	i) Barren Interzone (5th).	62
	3- El Hasana composite Section.	62
	a) Globotruncanella havanensis.	63
	b) Globotruncana aegyptiaca.	65
	c) Gansserina gansseri.	66
	d) Abathomphalus mayaroensis.	68
	CRETACEOUS BOUNDARIES WITHIN THE STUDIED SECTIONS	71 ,
	1- The Malha Formation.	71
	2- The Risan Aneiza Formation.	71
	3- The Halal Formation.	74
	4- The Abu Qada Formation.	75
	5- The Wata Formation.	76
	6- Sudr Chalk.	77
	PALEOECOLOGY	78
	1- Malha Formation.	79
	2- Risan Aneiza Formation.	80
	3- Halal Formation.	84
	4- Abu Qada Formation.	86
	5- Wata Formation.	86
	6- Sudr Chalk,	87
	TAXONOMY	91
	1- Generalites.	91
	2- Systematic	91
	Superfamily: Hormosinacea.	92
	Superfamily: Lituolacea.	94
	Superfamily: Creatobuliminacea.	99
	Superfamily: Spiroplectamminacea.	100
	Superfamily: Verueuilinacea.	101
	Superfamily: Textulariacea.	104
	Superfamily: Nodosariacea.	108
	Superfamily: Heterohelicacea.	118
	Superfamily: Planomalinacea.	121
	Superfamily: Rotaliporacea.	122
	Superfamily: Globotruncanacea.	104

Chapte	er	Pag
No.	•	<u>No.</u> 138
	Superfamily: Bolivinacea.	
	Superfamily: Buliminacea.	. 141
	Superfamily: Pleurostomellacea.	143
	Superfamily: Discorbacea.	146
	Superfamily: Planorbulinacea.	[49
	Superfamily: Nonionacea.	152
	Superfamily: Chilostomellacea.	154
	Superfamily: Orbitolinacea.	157
VIII-	SUMMARY AND CONCLUSIONS	162
	REFERENCES.	166
	VITAE.	188
,	FOSSIL INDEX.	1.
	PLATES.	,
	ADADIC SUMMADV	

LIST OF FIGURES

:e		Page
	•	No.
	Regional geological map showing the location of the studied	
	sections.	2
	A rock stratigraphic clssification for the Upper Cretaceous	•
	(Ghorab, 1961).	8
	Field photo (Panoramic view)of Gabal Manzour looking NW,	
	showin the Lower Cretaceous rocks, (A), conformably overlain by	
	Cenomanian rocks, (C),	18
	Field photo (Panoramic view) of Gabal El Minsherah, looking SE,	1
	showing the Jurassic rocks (J), and the overlyin Cretaceous rocks.	ł
	(CR)	19
	Lithostratigraphic column of Gabal Manzour measured section.	20-21
	Lithostratigraphic column of Gabal El Minsherah measured	
	section.	22
	Lithostratigraphic column of El Hasana composite section.	23
	Stratigraphic correlation between the rock units of the studied	
	sections.	24
	Lithostratigraphic column of the Malha and Risan Aneiza	
	formations at Gabal Manzour.	26
	Field photo showing general view of the Risan Aniza Fm. (R) &	
	the Halai Fm. (H), at Gabal Manzour.	27
	Field photo showing cross-bedded & bioturbated sandstone in the	
	upper part of the Mallia Fin. (Bed-1, MZ-9), at Gabai Manzour.	27
12-	Field photo showing fine grained, bioturbated & cross bedded	
	sandstone in the upper part of the Malha Fm. (Bed-1, MZ-8&9), at	
	Gabal Manzour.	28
13-	Field photo showing mudstone (Bed-3, MZ-12), rich in small	
	gastropod in the lower part & dolomitic sandstone ledge (Bed-4,	
	MZ-13) in the middle part of the photo, in the lower part of the	
	Risan Aneiza Fm. at Gabal Manzour.	-28
	Field photo showing variegated & laminated shale in the lower	
	part of the Risan Anciza Fm. (Bed-7, MZ-25), at Gabal Manzour.	28
	Field photo showing sandstone with bard ferruginous bands in the	
	wer part of the Malha Fm. (Bed-1, MZ-5&6), at Gabal	
	uzour.	29

Figu	re	Page
<u>No.</u>		No.
16-	Field photo showing siltstone with nodular gypsum (see the	
	hanuner), followed by paleosol with sulphur (yellow color) in the	
	middle part of the Risan Anciza Fin. (Bed-15,MZ-49), at Gabal	
	Manzour.	29
17-	Field photo showing 2nd Actonella bed in the upper part of the	•
	Risan Aneiza Fm. (Bed-22, MZ-58), at Gabal Manzour.	29
18-	Lithostratigraphic column of the Mallia and Risan Aneiza	
	formations at Gabal El Minsherah.	30
19-	Field photo showing general view of the Risan Aneiza Fm. (R),	
	and the Halal Fm., (H), at Gabal El Minsherah.	32
20-	Field photo showing the songes of rudist-sponge reef-like complex	
	in the upper part of the Risan Aneiza Fm., (Bed-16, MS-23), at	
	Gabal El Minsherah.	32
21-	Field photo showing rudist in the upper part of the Risan Aneiza	
	Fm., (Bed-16, MS-23), at Gabal El Minsherah.	32
22-	Lithostratigraphic column of the Halal Formation at Gabal	
	Manzour.	35
23-	Field photo showing bioturbated limestone with Orbitolina cf.	
	concava (Lamarck), rudist and bivalvia in the lower part of the	
	Halal Fm., (Bed-26, MZ-68), at Gabal Manzour.	36
24-	Field photo showing rudist in the middle part of the Halal Fm,	
	(Bed-31, MZ-75), at Gabal Manzour.	36
25-	Field photo showing Orbitolina bank (O. cf. concava (Lamarck))	
	in the middle part of the Halal Fm., (Bed-36, MZ-81), at Gabal	
	Manzour.	36
26-	Lithostratigraphic column of the Halal Formation at Gabal El	
	Minsherah.	37
27-	Field photo showing dolomitic limestone with bands filled by	
	pelecypod shell fragments in the middle part of the Halal Fm.,	
	(Bed-25, MS-38), at Gabal El Minsherali.	38
28-	Field photo showing reefal facies (Millestromal reef), in the upper	_
	part of the Halal Fm., (Bed-27, MS-56), at Gabal El Minsherah.	38
29-	Lithostratigraphic column of the Abu Qada and Wata Formations	_
	at Gabal El Minsherab.	40

Figu	re	Page
No.		No.
30-	Field photo showing general view of the Abu Qada Fm. (A) & the	
	Wata Fnt. (W), at Gabal El Minsherah.	41
31-	Field photo showing rudist bank in the Wata Fm., (Bed-40, MS-	
	81), at Gabai El Minsherah.	41
32-	Field photo (panoramic view) of Gabal El Hasana (not studied),	
	showing from base to top, the Sudr Chalk, Esna Shale and Thebes	
	lintestone.	45
33-	Field photo showing the contact between Sudr Chalk at base and	
	and Esna Shale at top at West Safra El Hasna (WSF).	45
34-	Field photo showing the base of the Sudr Chalk filled with	
	Pycnodonta vesicularis (Lamarck), northeast Gabal El Minsherab	
	(NEM).	46
35-	Field photo (close-up view) showing the Scattered Pycnodonta	
	vesicularis (Lamarck), at Base of the Sudr Chalk, Northeast Gabal	
	Minsherah (NEM).	46
36-	Field photo (close-up view) showing sponges in the Sudr Chalk, at	
	Southeast El Minsherah (SEM).	46
37-	The relation of different stratigraphic subdivisions given by	
	different authors for the Cretaceous in Sinai and neighbouring	
	areas.	48
38-	Distribution chart of the foraminiferal and some megafossils	
	species in Gabal Manzour.	50
39-	Distribution chart of the foraminiferal and some megafossils	
	species in Gabal El Minsheralı.	51
40-	Distribution chart of the foraminiferal species in El Hasaua	
	composite section.	52
41-	Correlation between the proposed biozones in Gabal Manzour and	
	those recognized in Gabal El Minsherah studied sections.	53
42-	Correlation of the Maastrichtain planktic foraminiferal zones with	
	those recorded from other parts of the World.	64
43-	Correlation of the Creataceous zones in the studied sections with	
	those proposed by different authors in Gulf of Suez Region and	
	Sinai.	70
44-	Fisher a Diversity paleoenvironmental indices of the Malha and	
	Risan Aneiza formations.	81

	e	Page	
		<u>No.</u>	
	The paleoecologic parameters and interpreted paleoenvironment at		
	Gabal Manzour section.	82	
46-	The paleoecologic parameters and interpreted paleoenvironment at		
	Gabal El Minsherah section.	83	
47-	Fisher α Diversity paleoenvironmental indices of the Sudr Chalk of		
	El Ilasana composite section.	88	
48-	The paleoecologic parameters and interpreted paleoenvironment of		
	El Hasana composite section.	89	