

C D

**EFFECT OF PREOPERATIVE INSTRUCTION
ON THE POSTOPERATIVE OUTCOMES FOR
CHILDREN UNDERGOING HEART SURGERY**

Thesis
Submitted for Partial Fulfillment of the
Requirements of the Master Degree
in Pediatrics Nursing

By

AMAL AHMED NAWAL EL SAYED MORSE

B.Sc. Nursing
Demonstrator in High Institute of Nursing
Ain Shams University

Supervised by

PROF. DR. ZEINAB LOUTFI
Chairman of Psychiatric Nursing Dept.,
Director of High Institute of Nursing
Ain Shams University

PROF. DR. AYMAN SHOEB
Professor of Cardiothoracic Surgery
Faculty of Medicine
Ain Shams University

DR. WAFAA EL SAYED ABD EL GILEEL

Lecturer of Pediatrics Nursing
High Institute of Nursing
Ain Shams University

**Ain Shams University
High Institute of Nursing
1997**



بسم الله الرحمن الرحيم

قالوا سبحانك لا علم لنا إلا ما علمتنا،
إنك أنت العليم الحكيم

صدق الله العظيم

سورة البقرة، الآية ٣٢

ACKNOWLEDGMENT

I wish to express my deep appreciation and gratitude to Prof. Dr. Zeinab Loutfy, Professor of Psychiatric Nursing, High Institute of Nursing, Ain Shams University, for giving me the privilege of working under her supervision and for her helpful guidance. Her advice was always stimulating and especially essential to complete this work.

I wish to express my sincere thanks and gratitude to Prof. Dr. Ayman Shoeib, Professor of Cardiothoracic Surgery, Faculty of Medicine, Ain Shams University, for his great support, encouragement, patience, enthusiasm, criticism, fruitful comments and help without which this work would have never seen light.

I am also grateful to Dr. Wafaa El-Sayed Abdel Gileel, Lecturer of Pediatrics Nursing, High Institute of Nursing, Ain Shams University, for her meticulous revision of this work, her great cooperation and generous help.

To my father, my mother, my brother, my sister, and to everyone who participated in some way or other to let this work come to such a final picture, I owe my thanks and gratitude.

CONTENTS

| | page |
|------------------------|-------------|
| INTRODUCTION | 1 |
| AIM OF THE STUDY | 3 |
| REVIEW OF LITERATURE | 4 |
| SUBJECTS & METHODS | 45 |
| RESULTS | 53 |
| DISCUSSION | 89 |
| SUMMARY AND CONCLUSION | 99 |
| RECOMMENDATIONS | 102 |
| REFERENCES | 103 |
| APPENDICES | 114 |
| ARABIC SUMMARY | |

LIST OF TABLES

| | Page |
|---|------|
| Table (I): Number and percentage distribution of parents in both groups according to their biosocial characteristics | 54 |
| Table (II): Number and percentage distribution of children in both groups according to their biosocial characteristics | 56 |
| Table (III): Number and percentage distribution of children in both groups according to their anxiety state pre- and postoperatively | 57 |
| Table (IV): Number and percentage distribution of children in both groups according to their anxiety trait pre- and postoperatively | 59 |
| Table (V): Number and percentage distribution of children in both groups according to their hemodynamic stability in relation to vital signs as estimated hourly after complete recovery | 60 |
| Table (VI): Number and percentage distribution of children in both groups according to their hemodynamic stability in relation to pH of the blood as estimated hourly after complete recovery | 61 |
| Table (VII): Number and percentage distribution of children according to length of postoperative stay in ICU | 62 |
| Table (VIII): Number and percentage distribution of children in both groups according to their reaction stability as estimated hourly after complete recovery | 63 |

| | |
|--|----|
| Table (IX): Number and percentage distribution of children in both groups according to their reaction stability as estimated hourly after complete recovery | 69 |
| Table (X): Number and percentage distribution of children in both groups according to their reaction stability as estimated hourly after complete recovery | 73 |
| Table (XI): Number and percentage distribution of children in both groups according to their reaction stability as estimated hourly after complete recovery | 76 |
| Table (XII): Number and percentage distribution of children in both groups according to their reaction stability as estimated hourly after complete recovery | 80 |
| Table (XIII): Number and percentage distribution of children in both groups according to their cooperation in postoperative activities | 74 |

LIST OF FIGURES

| | Page |
|---|-------------|
| Fig. (I): Flow of blood through CPB circuit | 9 |
| Fig. (II): Comparison between study and control groups as regard postoperative anxiety state | 58 |
| Fig. (III): Comparison between study and control groups as regards facial expression as estimated hourly after complete recovery | 65 |
| Fig. (IV): Comparison between study and control groups as regard restlessness as estimated hourly after complete recovery | 66 |
| Fig. (V): Comparison between study and control groups as regard stretching muscles as estimated hourly after complete recovery | 67 |
| Fig. (VI): Comparison between study and control groups as regard striking legs as estimated hourly after complete recovery | 68 |
| Fig. (VII): Comparison between study and control groups as regard weeping as estimated hourly after complete recovery | 71 |
| Fig. (VIII): Comparison between study and control groups as regard crying as estimated hourly after complete recovery | 72 |
| Fig. (IX): Comparison between study and control groups as regard biting on the endotracheal tube as estimated hourly after complete recovery | 74 |
| Fig. (X): Comparison between study and control groups as regard trying to talk aggressively while endotracheal tube is in as estimated hourly after complete recovery | 75 |

| | |
|--|----|
| Fig. (XI): Comparison between study and control groups as regards frightened to move while the chest tube is in as estimated hourly after complete recovery | 78 |
| Fig. (XII): Comparison between study and control groups as regard move wrong movement while chest tube is in as estimated hourly after complete recovery | 79 |
| Fig. (XIII): Comparison between study and control groups as regard pushing nurses' hands (during painful procedures) as estimated hourly after complete recovery | 82 |
| Fig. (XIV): Comparison between study and control groups as regard trying to pull connections as estimated hourly after complete recovery | 85 |
| Fig. (XV): Comparison between study and control groups as regard cooperation during breathing exercises after complete recovery | 86 |
| Fig. (XVI): Comparison between study and control groups as regard cooperation during positioning after complete recovery | 87 |
| Fig. (XVII): Comparison between study and control groups as regard cooperation during suctioning after complete recovery | 88 |

LIST OF ABBREVIATIONS

| | |
|-------|---|
| CABG: | Coronary artery bypass graft |
| CPB: | Cardiopulmonary bypass |
| ICU: | Intensive care unit |
| V.S.: | Vital signs |
| PT.: | Prothrombin time |
| PTT: | Partial thromboplastin time |
| DIC: | Disseminated intravascular coagulopathy |
| ECG: | Electrocardiogram |

