

**STUDIES  
ON THE SYNTHESIS AND STRUCTURE ELUCIDATION  
OF  
SOME NEW ORGANOMETALIC COMPOUNDS**


5247.2  
A. El

A Thesis

Submitted  
for the Requirements of  
Ph.D degree in Chemistry

Presented by  
Attia El- Sayed Attia Auf  
M.SC. (1990)

57121



Ain Shams University  
Faculty of Science  
Chemistry Department

(1994)

**STUDIES**  
**ON THE SYNTHESIS AND STRUCTURE ELUCIDATION**  
**OF**  
**SOME NEW ORGANOMETALIC COMPOUNDS**

**THESIS ADVISORS**

*Prof. Dr. M. F. El-Shahat*  
Ain Shams University at Cairo

*Prof. Dr. Cortlandt G. Pierpont*  
University of Colorado at Boulder

**APPROVAL**

*El-Shahat*

*Cortlandt G. Pierpont*



*Prof. Dr. F. Fahmy*  
*A. F. M. Fahmy*  
Head of Chemistry Department



*To my wife Holly Jones.  
You have always been there for me;  
this one's for you.*

## *ACKNOWLEDGEMENT*

First and foremost I would like to thank God for giving me the opportunity and willpower to accomplish my Ph.D.

I would like to acknowledge the following people for their help, encouragement, and friendship during my graduate studies: *Dr. M. F. El-Shahat*, professor of inorganic and analytical chemistry at Ain Shams University, and *Dr. Cortlandt G. Pierpont*, Professor of inorganic and biochemistry at C. U. Boulder.

I would like to thank my family for their support and tolerance during my lengthy absence. Especially my mother who has stuck with me through the hard times. I love you !

Finally, I want to thank my friends in the Arab Student Club at C. U. Boulder for being there for me in times of need. God bless all of you.

*Attia S. Attia*

## CONTENTS

<b>LIST OF TABLES</b>	v
<b>LIST OF FIGURES</b>	vii
<b>SUMMARY</b>	xii
<b>I. INTRODUCTION</b>	1
<b>II. EXPERIMENTAL</b>	10
2.1 Instrumentation and Methods	10
2.2 Solvents and Reagents	13
2.3 Synthesis and Studies of Manganese Quinone Complexes	14
2.3.1 Synthesis of $\text{Mn}(3,6\text{-DBSQ})_3$	14
2.3.2 Synthesis of $\text{Mn}(\text{Bupy})_2(3,6\text{-DBCat})_2$	15
2.3.3 Synthesis of $\text{Mn}(\text{py})_2(3,6\text{-DBCat})_2$	17
2.3.4 Synthesis of $\text{Mn}(\text{bpy})(3,6\text{-DBCat})_2$	17
2.3.5 Synthesis of $\text{Mn}(\text{phen})(3,6\text{-DBCat})_2$	18
2.3.6 Synthesis of $[\text{Mn}(\text{pz})(3,6\text{-DBSQ})(3,6\text{-DBCat})]_n$ polymer	19
2.3.7 Synthesis of $\text{Mn}(\text{thf})_2(3,6\text{-DBSQ})(3,6\text{-DBCat})$	20
2.3.8 Synthesis of $\text{Mn}(4,4'\text{-bpy})_2(3,6\text{-DBSQ})(3,6\text{-DBCat})$	21
2.3.9 Synthesis of $\text{Mn}(\text{NO}_2)(3,6\text{-DBSQ})_2$	22
2.4 Synthesis and Studies of Iron Quinone Complexes	23
2.4.1 Synthesis of $\text{Fe}(3,6\text{-DBSQ})_3$	23
2.4.2 Synthesis of $[\text{Fe}(\text{OMe})(3,6\text{-DBSQ})_2]_2$	25

2.4.3	Synthesis of $\text{Fe}(\text{Bupy})_2(3,6\text{-DBSQ})(3,6\text{-DBCat})$	25
2.4.4	Synthesis of $\text{Fe}(\text{py})_2(3,6\text{-DBSQ})(3,6\text{-DBCat})$	26
2.4.5	Synthesis of $\text{Fe}(\text{tmeda})(3,6\text{-DBSQ})(3,6\text{-DBCat})$	27
2.4.6	Synthesis of $\text{Fe}(\text{tmeda})(3,5\text{-DBSQ})(3,5\text{-DBCat})$	28
2.4.7	Synthesis of $\text{Fe}(\text{bpy})(3,6\text{-DBSQ})(3,6\text{-DBCat})$	29
<b>2.5</b>	<b>X-Ray Crystallographic analysis</b>	30
2.5.1	Structure Determination of $\text{Mn}(3,6\text{-DBSQ})_3$	30
2.5.2	Structure Determination of $\text{Mn}(\text{Bupy})_2(3,6\text{-DBCat})_2$	31
2.5.3	Structure Determination of $\text{Mn}(\text{bpy})(3,6\text{-DBCat})_2$	31
2.5.4	Structure Determination of $[\text{Mn}(\text{pz})(3,6\text{-DBSQ})(3,6\text{-DBCat})]_n$	32
2.5.5	Structure Determination of $\text{Mn}(\text{thf})_2(3,6\text{-DBSQ})(3,6\text{-DBCat})$	33
2.5.6	Structure Determination of $\text{Mn}(4,4'\text{-bpy})_2(3,6\text{-DBSQ})(3,6\text{-DBCat})_2$	34
2.5.7	Structure Determination of $\text{Mn}(\text{NO}_2)(3,6\text{-DBSQ})_2$	35
2.5.8	Structure Determination of $\text{Fe}(3,6\text{-DBSQ})_3$	36
2.5.9	Structure Determination of $[\text{Fe}(\text{OMe})(3,6\text{-DBSQ})_2]_2$	36
2.5.10	Structure Determination of $\text{Fe}(\text{Bupy})_2(3,6\text{-DBSQ})(3,6\text{-DBCat})$	37
2.5.11	Structure Determination of $\text{Fe}(\text{tmeda})(3,5\text{-DBSQ})(3,5\text{-DBCat})$	38
2.5.12	Structure Determination of $\text{Fe}(\text{tmeda})(3,6\text{-DBSQ})(3,6\text{-DBCat})$	39
2.5.13	Structure Determination of $\text{Fe}(\text{bpy})(3,6\text{-DBSQ})(3,6\text{-DBCat})$	37

<b>III. RESULTS AND DISCUSSION</b>	41
<b>3.1 Quinone Complexes of Manganese</b>	41
3.1.1 Quinone Complexes of Manganese(IV)	43
3.1.1.1 Characterization of $Mn(Bupy)_2(3,6-DBCat)_2$ and $Mn(py)_2(3,6-DBCat)_2$	43
3.1.1.2 Characterization of $Mn(bpy)(3,6-DBCat)_2$ and $Mn(phen)(3,6-DBCat)_2$	52
3.1.2 Quinone Complexes of Manganese(III)	63
3.1.2.1 Characterization of $Mn(3,6-DBSQ)_3$	63
3.1.2.2 Characterization of $Mn(4,4'bpy)_2(3,6-DBSQ)(3,6-DBCat)$	73
3.1.2.3 Characterization of $[Mn(pz)(3,6-DBSQ)(3,6-DBCat)]_n$	79
3.1.2.4 Characterization of $Mn(thf)_2(3,6-DBSQ)(3,6-DBCat)$	86
3.1.3 Quinone Complex of Manganese(II)	92
3.1.3.1 Characterization of $Mn(NO_2-phen)(3,6-DBSQ)_2$	92
<b>3.2 Quinone Complexes of Iron(III)</b>	96
3.2.1 Characterization of $Fe(3,6-DBSQ)_3$	98
3.2.2 Characterization of $[Fe(OMe)(3,6-DBSQ)_2]_2$	106
3.2.3 Characterization of $Fe(bpy)(3,6-DBSQ)(3,6-DBCat)$	112
3.2.4 Characterization of $Fe(tmeda)(3,5-DBSQ)(3,5-DBCat)$ and $Fe(tmeda)(3,6-DBSQ)(3,6-DBCat)$ Complexes	124
3.2.5 Characterization of $Fe(Bupy)_2(3,6-DBSQ)(3,6-DBCat)$	136
<b>3.3 Characterization of Some Iron and Manganese Complexes by Infrared Spectroscopy</b>	147



3.3.1	Infrared Spectra of $\text{Fe}(\text{3,6-DBSQ})_3$ and $\text{Mn}(\text{3,6-DBSQ})_3$	148
3.3.2	Infrared Spectra of $[\text{Fe}(\text{OMe})(\text{3,6-DBSQ})_2]_2$ and $\text{Mn}(\text{NO}_2\text{-phen})(\text{3,6-DBSQ})_2$	148
3.3.3	Infrared Spectra of $\text{Fe}(\text{tmeda})(\text{3,6-DBSQ})(\text{3,6-DBCat})$ $\text{Fe}(\text{bpy})(\text{3,6-DBSQ})(\text{3,6-DBCat})$ and $\text{Mn}(\text{bpy})(\text{3,6-DBCat})_2$	151
<b>IV. GENERAL DISCUSSION AND CONCLUSION</b>		153
<b>REFERENNCES</b>		158
<b>APPENDIXES</b>		165
<b>ARABIC SUMMARY</b>		

## LIST OF TABLES

3.1	Selected Bond Lengths and Angles for $\text{Mn}(\text{Bupy})_2(3,6\text{-DBCat})_2$	46
3.2	Representative M-O and M-N Bond Lengths of Selected Manganese-Quinone Complexes	44
3.3	Selected Bond Lengths and Angles for $\text{Mn}(\text{bpy})(3,6\text{-DBCat})_2$	57
3.4	Selected Bond Lengths and Angles for $\text{Mn}(3,6\text{-DBSQ})_3$	65
3.5	Selected Bond Lengths and Angles for $\text{Mn}(4,4'\text{-bpy})_2(3,6\text{-DBSQ})(3,6\text{-DBCat})$	77
3.6	Selected Bond Lengths and Angles for $[\text{Mn}(\text{pz})(3,6\text{-DBSQ})(3,6\text{-DBCat})]_n$	83
3.7	Selected Bond Lengths and Angles for $\text{Mn}(\text{thf})_2(3,6\text{-DBSQ})(3,6\text{-DBCat})$	90
3.8	Selected Structural Data for $\text{trans-Mn}^{\text{III}}$ Complexes	88
3.9	Selected Bond Lengths and Angles for $\text{Mn}(\text{NO}_2\text{-phen})(3,6\text{-DBSQ})_2$	95
3.10	Selected Bond Lengths and Angles for $\text{Fe}(3,6\text{-DBSQ})_3$	100
3.11	Selected Bond Lengths and Angles for $[\text{Fe}(\text{OMe})(3,6\text{-DBSQ})_2]_2$	110
3.12	Selected Bond Lengths and Angles for $\text{Fe}(\text{bpy})(3,6\text{-DBSQ})(3,6\text{-DBCat})$	117

3.13	Mossbauer Least-Squares Fitting Parameters for Some Fe(bpy)(SQ)(Cat) Complexes	119
3.14	Selected Bond Lengths and Angles for Fe(tmeda)(3,6-DBSQ)(3,6-DBCat)	127
3.15	Selected Bond Lengths and Angles for Fe(tmeda)(3,5-DBSQ)(3,5-DBCat)	129
3.16	Selected Bond Lengths and Angles for Fe(Bupy) <sub>2</sub> (3,6-DBSQ)(3,6-DBCat)	141

## LIST OF FIGURES

1.1	Redox series of 1,2-benzoquinone	1
1.2	Relative ordering of metal and quinone $\pi$ orbital energy	2
1.3	General synthetic procedure of metal quinone complexes	4
1.4	Spin states arising from coupling of Fe(III) ion to $S=1/2$ semiquinone ligands	7
3.1	Crystal structure of $\text{Mn}(\text{Bupy})_2(3,6\text{-DBCat})_2$	45
3.2	Packing diagram for $\text{Mn}(\text{Bupy})_2(3,6\text{-DBCat})_2$	48
3.3	Temperature dependent effective magnetic moment for $\text{Mn}(\text{Bupy})_2(3,6\text{-DBCat})_2$	49
3.4a	Electronic spectrum of $\text{Mn}(\text{Bupy})_2(3,6\text{-DBCat})_2$ recorded on a solid sample at 298 K prepared as a KBr pellet.	50
3.4b	Electronic spectrum of $\text{Mn}(\text{py})_2(3,6\text{-DBCat})_2$ recorded on a solid sample at 298 K prepared as a KBr pellet.	51
3.5	Temperature-dependent changes in the intensity of the $\text{Mn(III)} \rightarrow \text{SQ}$ charge transfer band of $\text{Mn}(\text{Bupy})_2(3,6\text{-DBCat})_2$ recorded on a solid sample at temperatures 298 K (1), 327 K (2), 343 K (3), 285 K (4), and 243 K (5).	53
3.6	The cyclic voltammogram of $\text{Mn}(\text{Bupy})_2(3,6\text{-DBCat})_2$ in pyridine- $\text{CH}_2\text{Cl}_2$	54
3.7	Crystal structure of $\text{Mn}(\text{bpy})(3,6\text{-DBCat})_2$	56
3.8	The stacking interaction between bipyridine ligands of adjacent $\text{Mn}(\text{bpy})(3,6\text{-DBCat})_2$ molecules.	58

3.9	Streo view showing The stacking interaction between bipyridine ligands of adjacent $\text{Mn}(\text{bpy})(3,6\text{-DBCat})_2$ molecules.	59
3.10	Temperature dependent effective magnetic moment for $\text{Mn}(\text{phen})(3,6\text{-DBCat})_2$ .	61
3.11	Temperature-dependent changes in the intensity of the $\text{Mn}(\text{III})\rightarrow\text{SQ}$ charge transfer band of $\text{Mn}(\text{bpy})(3,6\text{-DBCat})_2$ recorded on solid sample at temperatures 298 K (1), 325 K (2), and 343 K (3) in the NIR region.	62
3.12	Crystal structure of $\text{Mn}(3,6\text{-DBSQ})_3$	64
3.13	Packing digram for $\text{Mn}(3,6\text{-DBSQ})_3$	67
3.14	The bond lengths and angles within the 5-member ring formed by the semiquinone ligands with the central Mn atom in $\text{Mn}(3,6\text{-DBSQ})_3$	68
3.15	Temperature dependent effective magnetic moment for $\text{Mn}(3,6\text{-DBSQ})_3$ .	70
3.16	The low-lying energy states for $\text{Mn}^{\text{III}}(\text{SQ})_3$ resulting from an intramolecular antiferromagnetic exchange interaction between the unpaired electrons of the o-semiquinone ligands and the metal ion.	71
3.17a	Infrared spectrum in KBr showing $\text{Mn}(\text{III})\rightarrow\text{SQ}$ CT transition in the NIR region.	72
3.17b	Electronic spectrum of $\text{Mn}(3,6\text{-DBSQ})_3$ recorded on a solid sample at 298 K prepared as a KBr pellet.	72
3.17c	Temperature-dependent changes in the intensity of the charge transfer bands of $\text{Mn}(3,6\text{-DBSQ})_3$ recorded on solid sample at temperatures between 298 K (1), 318 K (2), and 350 K (3).	74
3.18	Crystal structure of $\text{Mn}(4,4'\text{-bpy})_2(3,6\text{-DBSQ})(3,6\text{-DBCat})$	76
3.19	Temperature dependent effective magnetic moment for $\text{Mn}(4,4'\text{-bpy})_2(3,6\text{-DBSQ})(3,6\text{-DBCa})$ .	78

3.20	Electronic spectrum of $\text{Mn}(4,4'\text{-bpy})_2(3,6\text{-DBSQ})(3,6\text{-DBCat})$ recorded on a solid sample at 298 K prepared as a KBr pellet.	80
3.21	Crystal structure of a monomeric unit of the $[\text{Mn}(\text{pz})(3,6\text{-DBSQ})(3,6\text{-DBCat})]_n$ polymer.	82
3.22:	View of the polymeric chains in the crystal lattice of the $[\text{Mn}(\text{pz})(3,6\text{-DBSQ})(3,6\text{-DBCat})]_n$ polymer.	84
3.23	Temperature dependent effective magnetic moment for $[\text{Mn}(\text{pz})(3,6\text{-DBSQ})(3,6\text{-DBCat})]_n$ .	85
3.24	Temperature-dependent changes in the intensity of the charge transfer band of $[\text{Mn}(\text{pz})(3,6\text{-DBSQ})(3,6\text{-DBCat})]_n$ recorded on solid sample at temperatures 298 K (1), 320 K (2), and 345 K (3).	87
3.25	Crystal structure of $\text{Mn}(\text{thf})_2(3,6\text{-DBSQ})(3,6\text{-DBCat})$ .	89
3.26	Packing diagram for $\text{Mn}(\text{thf})_2(3,6\text{-DBSQ})(3,6\text{-DBCat})$ .	91
3.27	Electronic spectrum of $\text{Mn}(\text{thf})_2(3,6\text{-DBSQ})(3,6\text{-DBCat})$ recorded on a solid sample at 298 K prepared as a KBr pellet.	93
3.28	Crystal structure of $\text{Mn}(\text{NO}_2\text{-phen})(3,6\text{-DBSQ})_2$ .	94
3.29	Temperature dependent effective magnetic moment for $\text{Mn}(\text{NO}_2\text{-phen})(3,6\text{-DBSQ})_2$ .	97
3.30	Crystal structure of $\text{Fe}(3,6\text{-DBSQ})_3$	99
3.31	Packing diagram for $\text{Fe}(3,6\text{-DBSQ})_3$	101
3.32	Temperature dependent effective magnetic moment for $\text{Fe}(3,6\text{-DBSQ})_3$ .	103
3.33a	Cyclic voltammogram of $\text{Fe}(3,6\text{-DBSQ})_3$ in $\text{CH}_2\text{Cl}_2$ .	105
3.33b	Cyclic voltammogram of $\text{Fe}(3,6\text{-DBSQ})_3$ in $\text{CH}_3\text{CN}$ .	107

3.34	Proposed reaction scheme for the formation of $[\text{Fe}(\text{OMe})(3,6\text{-DBSQ})_2]_2$ from $\text{Fe}(3,6\text{-DBSQ})_3$ .	108
3.35	Crystal structure of $[\text{Fe}(\text{OMe})(3,6\text{-DBSQ})_2]_2$ .	109
3.36:	Observed (o) and calculated (–) effective magnetic moment for $[\text{Fe}(\text{OMe})(3,6\text{-DBSQ})_2]_2$ .	113
3.37	$^1\text{H}$ NMR spectrum of $[\text{Fe}(\text{OMe})(3,6\text{-DBSQ})_2]_2$ in d-chloroform.	114
3.38	Crystal structure of $\text{Fe}(\text{bpy})(3,6\text{-DBSQ})(3,6\text{-DBCat})$ .	116
3.39:	$^{57}\text{Fe}$ Mossbauer spectrum for 110 K sample of $\text{Fe}(\text{bpy})(3,6\text{-DBSQ})(3,6\text{-DBCat})$ .	118
3.40	Temperature dependent effective magnetic moment for $\text{Fe}(\text{bpy})(3,6\text{-DBSQ})(3,6\text{-DBCat})$ .	120
3.41	Stacking interaction between bipyridine ligands of adjacent $\text{Fe}(\text{bpy})(3,6\text{-DBSQ})(3,6\text{-DBCat})$ molecules.	122
3.42	The electronic spectra of $\text{Fe}(\text{bpy})(3,6\text{-DBSQ})(3,6\text{-DBCat})$ recorded in tetrahydrofuran (a), $\text{CH}_2\text{Cl}_2$ (b), and toluene (c) solvents.	123
3.43	Cyclic voltammogram of $\text{Fe}(\text{bpy})(3,6\text{-DBSQ})(3,6\text{-DBCat})$ in $\text{CH}_2\text{Cl}_2$ .	125
3.44	Crystal structure of $\text{Fe}(\text{tmeda})(3,6\text{-DBSQ})(3,6\text{-DBCat})$ .	126
3.45	Crystal structure of $\text{Fe}(\text{tmeda})(3,5\text{-DBSQ})(3,5\text{-DBCat})$ .	128
3.46	C-C bonds within the semiquinone ring (a) $\text{Fe}(\text{tmeda})(3,6\text{-DBSQ})(3,6\text{-DBCat})$ (b) $\text{Fe}(\text{tmeda})(3,5\text{-DBSQ})(3,5\text{-DBCat})$ .	131
3.47	Temperature dependent effective magnetic moment for $\text{Fe}(\text{tmeda})(3,6\text{-DBSQ})(3,6\text{-DBCat})$ .	132
3.48	$^1\text{H}$ NMR spectrum of in $\text{Fe}(\text{tmeda})(3,5\text{-DBSQ})(3,5\text{-DBCat})$ d-chloroform.	133
3.49	Electronic spectra of $\text{Fe}(\text{tmeda})(3,5\text{-DBSQ})(3,5\text{-DBCat})$ recorded in $\text{CH}_3\text{CN}$ (a), $\text{CH}_2\text{Cl}_2$ (b), hexane (c), and toluene (d) solvents.	135