

AIN SHAMS UNVIERSITY
FACULTY OF MEDICINE
DEPARTMENT OF OBSTETRICS AND GYNAECOLOGY

A Thesis

Submitted for Partial Fulfillment of
Master Degree
in Obstetrics and Gynaecology

**Comparative Study of Short Term
Morbidity Between Closure and
Non-Closure of Parietal peritoneum
during Caesarean section**

By

Ahmed Mohamed Ibrahim
M.B., B.Ch.

Under Supervision Of:

PROF. DR. IBRAHIM EL-MITWALLY SAMANA
Professor of Obstetrics and Gynaecology,
Faculty of Medicine,
Ain Shams University


DR. KHALED HUSSEIN ABDEL-RAZEK
Lecturer of Obstetrics and Gynaecology,
Faculty of Medicine,
Ain Shams University

612.86

A.M

1997





INDEX

Part	Subject	Page
I	Introduction	1
II	Aim of the work	4
II	Review of literature:	
	♦ Caesarean section	5
	♦ Short term postoperative complications	66
	♦ Peritoneum	108
III	Subjects and methods	131
IV	Results	138
V	Discussion	170
VI	Summary and Conclusions	176
VII	Recommendations	181
VIII	References	182
IX	Arabic summary	----

LIST OF TABLES

No.	Title	Page
1	Clinical qualitative characteristics of the non closure (I) and closure (II) group	138
2	Indications of C.S. in both groups	142
3	Clinical quantitative characteristics of the non closure group	144
4	Clinical quantitative characteristics of the closure group	145
5	Comparison between group I and group II regarding age, parity and gestational age	146
6	Comparison between group I and group II regarding operative time in minutes.	150
7	Comparison between group I and group II regarding PIQS	152
8	Comparison between group I and group II regarding number of analgesic doses given as needed	154
9	Comparison between group I and group II regarding time needed to restore bowel motions in hours	156
10	Comparison between group I and group II regarding postoperative hospital stay in days	158
11	Comparison between group I and group II regarding presence or absence of postoperative febrile morbidity	160
12	Causes of febrile morbidity in both groups	162
13	Comparison between afebrile and febrile groups regarding operative time	165
14	Correlation between operative time, number of analgesic doses needed, time required to regain bowel motion, PIQS and postoperative hospital stay	167

LIST OF FIGURES

No.	Title	Page
1	Vertical abdominal incisions	19
2	Three types of lower abdominal transverse incisions	20
3	Modified Pfannenstiel incision	23
4	The visceral peritoneum is grasped just above the bladder and incised in the midline	29
5	The visceral peritoneum is undermined and incised laterally	30
6	Dissection of urinary bladder	31
7	Uterine incision	33
8	Extension of uterine incision by fingers or scissor	34
9	Delivery of the baby	39
10	Delivery of the head assisted by fundal pressure	40
11	Laparoscopy	42
12	Delivery of deeply engaged head	44
13	Delivery of the head by Kielland forceps	45
14	Splint procedure	46
15	Delivery of placenta	48
16	Closure of the uterus	51
17	Visceral peritoneal closure	55
18	Parietal peritoneal closure	56
19	Evaluation of possible pulmonary embolism	80
20	Pathogenesis of puerperal infection	89
21	Technique of enbloc wound reclosure	101
22	Mass abdominal closure	102

(continue)

LIST OF FIGURES

(Continue)

No.	Title	Page
23	Comparison between group I and group II regarding whether C.S. was primary or repeat	139
24	Comparison between group I and group II regarding whether C.S. was elective or indicated	140
25	Comparison between group I and group II regarding whether B.T.L. was done or not	141
26	Indications of C.S.	143
27	Comparison between group I and group II regarding the maternal age	147
28	Comparison between group I and group II regarding the parity	148
29	Comparison between group I and group II regarding the gestational age	149
30	Comparison between group I and group II regarding the operative time	151
31	Comparison between group I and group II regarding PIQS	153
32	Comparison between group I and group II regarding the number of analgesic doses given as needed	155
33	Comparison between group I and group II regarding the time needed to restore bowel motions in hour	157
34	Comparison between group I and group II regarding the postoperative hospital stay in days	159
35	Number and percentage of cases with febrile morbidity in group I and group II	161
36	Causes of febrile morbidity in group I	163
37	Causes of febrile morbidity in group II	164
38	Comparison between patients with febrile morbidity and patients without regarding operative time in minutes	166
39	Correlation between operative time and regain of bowel motions	168
40	Correlation between operative time and postoperative hospital stay	169

LIST OF ABBREVIATIONS

BTL	Bilateral tubal ligation
C.S.	Caesarean section
CMC	Carboxymethylcellulose
DIC	Disseminated intravascular coagulation
DVT	Deep venous thrombosis
ECG	Electrocardiogram
HA	Hyaluronic acid
HPF	High power field
IU	International unit
IV	Intravenous
KD	Kilodalton
NSAID	Nonsteroidal anti-inflammatory drugs
PIQS	Pain Intensity Quantitative Reference Scale
Pts	Patients
ROM	Rupture of membranes
SC	Subcutaneous
tPA	Tissue plasminogen activator
UTI	urinary tract infection

