

**CHARACTERIZATION OF MURINE
HUMORAL RESPONSE AGAINST
SCHISTOSOMA HAEMATOBII
ANTIGENS**

THESIS

Submitted to The
Faculty of Science
Ain Shams University
In Partial Fulfillment of The
Requirements for The Degree of
**MASTER OF SCIENCE
IN BIOCHEMISTRY**

BY

AMR MEDHAT MOHAMMED GHALEB

Supervisors

Prof. Dr. MOHAMMED MOHAMMED ABDEL-FATTAH

Biochemistry Dept.

Faculty of Science

Ain Shams University

Dr. KAMAL ALY FATHI SHALABY

Lecturer in the Biochemistry Dept.

Faculty of Science

Ain Shams University

Dr. BEVERLY L. MANGOLD

Immunologist

Naval American Medical Research Unit-3 (NAMRU-3)

1995



**CHARACTERIZATION OF MURINE
HUMORAL RESPONSE AGAINST
SCHISTOSOMA HAEMATOBIMUM
ANTIGENS**

THESIS

Submitted to The

Faculty of Science

Ain Shams University

In Partial Fulfillment of The

Requirements for The Degree of

MASTER OF SCIENCE

IN BIOCHEMISTRY

BY

AMR MEDHAT MOHAMMED GHALEB

Supervisors

Prof. Dr. MOHAMMED MOHAMMED ABDEL-FATTAH

Biochemistry Dept.

Faculty of Science

Ain Shams University

Dr. KAMAL ALY FATHI SHALABY

Lecturer in the Biochemistry Dept.

Faculty of Science


Ain Shams University

Dr. BEVERLY L. MANGOLD

Immunologist

Naval American Medical Research Unit-3 (NAMRU-3)

1995



**TO MY DEAR
MOTHER AND
MY WIFE**

Scientific Board:

Prof. Dr. Mohammed Mohammed Abdel-Fattah

Biochemistry Department,

Faculty of Science,

Ain Shams University.

Dr. Kamal Ali Fathi Shalaby

Lecturer in the Biochemistry Department,

Faculty of Science,

Ain Shams University.

Dr. Beverly L. Mangold

Immunologist,

Naval American Research Unit-3 (NAMRU-3).

ACKNOWLEDGMENT

I would like to express my gratefulness to Prof. Dr. Mohammed Abdel-Fattah, Biochemistry Dept., Ain Shams University, for his support and kind supervision for completing this work.

My great thanks to Dr. Kamal Shalaby, Lecturer in the Biochemistry Dept., Ain Shams University, for his suggestions, help, encouragement, supervision and efforts throughout the work.

I would also like to express my deepest and most sincere gratefulness to Dr. Beverly Mangold, Immunologist, NAMRU-3, who encouraged me to start the whole work, supervised supported and guided me and to whom I owe all what I have learned.

My special thanks to Dr. David A. Dean, Immunologist, NAMRU-3, for approval to start the work and for his assistance and advice.

I would also like to thank Dr. Moustafa Mansour, Head of Basic Sciences Department, NAMRU-3, Dr. Karim Kamal, Head of Technical Support Branch, NAMRU-3, and Dr. Sam Lewis, Technical Support Branch, NAMRU-3, for their kindness, generosity, help and assistance.

TABLE OF CONTENTS:

AIM OF WORK	1
INTRODUCTION:	2
1. Historical Background	2
2. Life Cycle, Infection and Illness	4
3. Human Immunity to Schistosomiasis	7
4. Prevention and Control	9
5. Immunity in Experimental Animal Models to Schistosomiasis	11
6. Schistosome Antigens	13
MATERIALS AND METHODS:	21
1. Parasites	21
2. Labeling male worms	23
3. Extraction of soluble proteins	23
4. Separation of worm glycoproteins	25
4.1. Preparation of cross-linked Con A-Sepharose	26
4.2. Glycoprotein extraction	27
5. Preparation and collection of mice sera	28
5.1. Mice groups	28
5.2. Separation of serum	30
5.3. Efficiency of mice immunization	30
6. Radioimmunoprecipitation	30
6.1. Radioimmunoprecipitation buffers	31
6.2. Activation of <i>Staphylococcus aureus</i>	32
6.3. Preparation of siliconized glass tubes	33

6.4. Preclearing of ³⁵ S-methionine labeled <i>S. haematobium</i> adult male worms lysate	33
6.5. Radioimmunoprecipitation	34
7. Electrophoretic Separation of Immunoprecipitated Antigens	35
7.1. One dimensional SDS-PAGE	35
7.1.1. Principle and apparatus used	35
7.1.2. Assembling the glass plates	36
7.1.3. Separating gel	37
7.1.4. Stacking gel	37
7.1.5. The electrode buffers	38
7.1.6. Running conditions	38
7.1.7. Gel staining	39
7.1.8. Detection of separated radiolabeled antigens	40
7.2. Two dimensional SDS-PAGE	40
7.2.1. Principle and apparatus used	40
7.2.2. Buffers used	41
7.2.3. First dimension	42
7.2.4. Second dimension	45
8. Electrophoretic transfer of proteins on nitrocellulose paper (Western Blots)	46
8.1. Principle	46
8.2. Buffers used	47
8.3. Electrophoretic transfer	47
8.4. Screening of transferred immunogenic proteins using enzyme-conjugated antibodies	48
RESULTS	50

DISCUSSION	96
REFERENCES	104
ABSTRACT.	
ENGLISH SUMMARY.	
ARABIC SUMMARY.	

LIST OF TABLES:

Table 1.	Immunoinefficiency experiment.	62.
Table 2.	³⁵ S-methionine labeled total and glycoproteins immuno-precipitated by infected and by immunized mice sera.	63.
Table 3.	³⁵ S-methionine labeled total and glycoproteins immuno-precipitated by infected and by immunized mice sera.	65.
Table 4.	³⁵ S-methionine labeled total and glycoproteins immuno-precipitated by immunized and challenged mice sera.	66.
Table 5.	Total and glycoproteins recognized by infected and by immunized mice sera on NC paper after electrophoretic transfer (Western Blot).	68.
Table 6.	Total and glycoproteins recognized by infected and by immunized mice sera on NC paper after electrophoretic transfer (Western Blot).	70.
Table 7.	Total and glycoproteins recognized by immunized and challenged mice sera on NC paper after electrophoretic transfer (Western Blot)..	72.

Table 8.	The sizes and pI's of spots that showed on autoradiographs (Fig. 13) of 2D SDS-PAGE after radioimmunoprecipitation of ³⁵ S-methionine labeled <i>S. haematobium</i> adult male worm total proteins using 20 weeks-infected female mice sera.	74.
Table 9.	The sizes and pI's of spots that showed on autoradiographs (Fig. 14) of 2D SDS-PAGE after radioimmunoprecipitation of ³⁵ S-methionine labeled <i>S. haematobium</i> adult male worm total proteins using 21 weeks-immunized female mice sera.	75.
Table 10.	The sizes and pI's of spots that showed on autoradiographs (Fig. 15) of 2D SDS-PAGE after radioimmunoprecipitation of ³⁵ S-methionine labeled <i>S. haematobium</i> adult male worm total proteins using 12 weeks-immunized and challenged female mice sera.	76.
Table 11.	The sizes and pI's of spots that showed on autoradiographs (Fig. 16) of 2D SDS-PAGE after radioimmunoprecipitation of ³⁵ S-methionine labeled <i>S. haematobium</i> adult male worm glycoproteins using 20 weeks-infected female mice sera.	77.
Table 12.	The sizes and pI's of spots that showed on autoradiographs (Fig. 17) of 2D SDS-PAGE after radioimmunoprecipitation of ³⁵ S-methionine labeled <i>S. haematobium</i> adult male worm	

glycoproteins using 21 weeks-immunized female mice sera. 78.

Table 13. The sizes and pI's of spots that showed on autoradiographs (Fig. 18) of 2D SDS-PAGE after radioimmunoprecipitation of ³⁵S-methionine labeled *S. haematobium* adult male worm glycoproteins using 12 weeks-immunized and challenged female mice sera. 79.

List of figures:

- Fig. 1. 1D SDS-PAGE autoradiograph of ^{35}S -methionine labeled *S. haematobium* adult male worm total proteins immunoprecipitated by infected (weeks 2, 4, 6 & 8 after infection) and by immunized (weeks 2, 4, 6 & 8 after first immunization) mice sera. 80
- Fig. 2. 1D SDS-PAGE autoradiograph of ^{35}S -methionine labeled *S. haematobium* adult male worm total proteins immunoprecipitated by infected (weeks 10, 12, 16 & 20 after infection) and by immunized (weeks 11, 13, 17 & 21 after first immunization) mice sera. 81
- Fig. 3. 1D SDS-PAGE autoradiograph of ^{35}S -methionine labeled *S. haematobium* adult male worm total proteins immunoprecipitated by immunized and challenged (weeks 2, 4, 8 & 12 after challenge) mice sera 82
- Fig. 4. 1D SDS-PAGE autoradiograph of ^{35}S -methionine labeled *S. haematobium* adult male worm glycoproteins immunoprecipitated by infected (weeks 2, 4, 6 & 8 after infection) and by immunized (weeks 2, 4, 6 & 8 after first immunization) mice sera. 83