# Comparison of various imaging modelities in patients with avascular necrosis of femoral head

## Essay

# Submitted for partial Fullfilment of the master degree of radiodiagnosis

 $\mathcal{B}y$ 

Dr. Amany abd el - Fattah El-Sayed

(M. B; B. ch)

Faculty of Medicine

Ain Shams University

52689


Under supervision of


# Professor Dr. Ahmed Talaat Professor of radiodiagnosis

Faculty of Medicine Ain Shams university

> Faculty of Medicine Ain Shams University 1996







### يرفع الله الذين آمنوا والذين أتوا العلم منكم درجات .


صدق الله العظيم

#### Acknowledgment

It is a pleasant obligation to express my deepest thanks to Dr. Ahmed Talaat, professor of radiodiagnosis, Ain Shams University for his sincere guidance, valuable instructions, meticulous supervision and continuous encouragement throughout this work.

Much credit also goes to my colleagues who so willingly help me to achieve this work

Amany Abd El Fattah



#### Contents

Introduction
Aim of the work
Review
Anatomical structure of the hip joint 1
Pathology 15
Conventional plain X-ray of the hip Joint 35
C. T. examination of the hip Joint 49
MR Imaging of the hip Joint60
Scintigraphy of avascular necrosis of femoral head 76
Discussion 92
Summary and conclusien 102
References 104
Arabic Summary

#### Table of figures

Fig. Page
1- Anatomy of head of femur
2- Sagittal section in hip joint
3- Comparison between normal and osteonecrotic femoral head $\dots$ 16
4- Coronal section of femoral head with osteonecrosis 16
5- Fatty marrow in normal and osteonecrotic femoral head 24
6- Core biopsy of osteonecrotic femoral head
7- Thromboembolism as etiological factor of osteonecrosis
8- Microscopic section of thrombus as etiological factor
9- Antero posterior view of normal hip joint
10- Stage II of osteonecrosis by X-ray
11- The radiolucent crescent line (early diagnostic sign)
12- Stage IV of osteonecrosis by X- ray
13- Stage IV of osteonecrosis "classic crescent sign"
14- Classic saddle shape (pathognomonic in X- ray)47
15- Bite sign (radiographic finding) late sign47
16- Different modalities to diagnose early osteonecrosis 50
17- Subchondral fracture and collapse in hip joint (different modalities)52
18- Different modalities to diagnose early osteonecrosis 55
19- Different modalities to diagnose advanced osteonecrosis 57
20- Different modalities to diagnose advanced osteonecrosis 58

21- Coronal T1 weighted MRI stage (0) osteonecrosis	61
22- MRI findings in early osteonecrosis	63
23- Sagittal T1 weighted MRI , classic location of osteonecrosis	. 6
24- MRI findings in early osteonecrosis in Lt. hip	<b>7</b> 1
25- MRI findings in advanced osteonecrosis in Rt. hip	72
26- Stage I ( photopenic region in femoral head)	82
27- Stage II (by X-ray and scintigraphy)	85
28- Stage III (by x-ray and scintigraphy)	85
29- Stage IV (by X- ray and scintigraphy)	88
30- Stage IV (by X- ray and scintigraphy )	88
31- Stage V (by X - ray and scintigraphy)	90