PREVALENCE OF HEPATITIS-E VIRUS IN CHRONIC RENAL FAILURE PATIENTS UNDER REGULAR HAEMODIALYSIS

Thesis

Submitted for Partial Fulfillment of Master Degree in Internal Medicine

RAED M. A. AL-A

M.B., B.CH

616.614 R.M

Supervised by:

54(3)

Prof. Dr. AMR FATEEN

Professor of Internal Medicine & Gastroenterology Faculty of Medicine - Ain Shams University

Prof. Dr. ESSAM KHEDRE

Professor of Internal Medicine & Nephrology Faculty of Medicine - Ain Shams University

Prof. Dr. LAILA AL-SHAWARBY

Professor of Clinical Pathology
Faculty of Medicine - Ain Shams University

Faculty of Medicine Ain Shams University



CONTENTS

	page
Acknowledgment.	
• Introduction.	1
• Aim of the study.	3
• Review of literature.	_
- Hepatitis - A.	4
- Hepatitis - B.	8
- Hepatitis - D.	23
- Hepatitis - C.	25
- Hepatitis - E.	29
- Hepatitis - F.	48
- Hepatitis - G.	50
• Chronic renal failure & dialysis.	52
 Patients & Methods. 	65
• Results.	77
• Discussion.	80
• Summary.	86
• Conclusion & Recommendation.	88
• Tables & Figures.	89
• References.	100
Arabic Summary.	115

LIST OF TABLES

- Table (1): Relation between the patients group & the control group as regard age.
- Table (2): Relation between the patients group & the control group as regard sex.
- Table (3): Laboratory data (mean \pm S.D) of the patients group & the control group.
- Table (4): Distribution of HBsAg, HCVAB, and both in positive cases of HEV.
- Table (5): Distribution of hepatitis markers in the patients group.
- **Table (6):** Relation between number of blood transfusions & presence of hepatitis marker in positive cases for HEV.
- Table (7): Distribution of the mean value in the patients group & in the positive cases for HEV as regard; number of blood transfusions, duration of haemodialysis & hours/week on haemodialysis.

LIST OF FIGURES

- Figure (1): Distribution of the mean value of age in the patients group & the control group.
- Figure (2): Relation between the patients group & the control group as regard sex.
- Figure (3): Distribution of HBsAg, HCVAB, both (HBsAg & HCVAB) and HEVIgG in the patients group.
- Figure (4): Distribution of HBsAg, HCVAB, and both in positive cases of HEV.

ACKNOWLEDGMENT

Acknowledgment

first and foremost, I feel always indebted to allah, the most beneficent and merciful.

I would like to express my sincere gratitude to **Prof. Dr. Amr Fateen**, Prof. of Internal Medicine & gastroentrology. Faculty of Medicine, Ain-Shams University. whose guidance, brotherly attitude, moral support and follow up of the progress of this work were certainly the most real steps in accomplishing this work.

Profound gratitude is expressed to **Prof. Dr. Essam Khedre**, Prof. of Internal Medicine & Nephrology. Faculty of Medicine, Ain-Shams University. For his continuous guidance, his fruitful criticism and generous advice throughout the whole work.

Great indebtedness is offered to **Prof. Dr. Laila Al-Shawarby**, Prof. of Clinical Pathology. Faculty of Medicine, Ain-Shams University. For her generous efforts she had made in the practical part of this work. Without her assisstance this work could not be done.

INTRODUCTION