

Exercise-Induced Proteinuria in Healthy Children & in Children With Nephropathy


Thesis

Submitted for partial fulfillment of the master degree in Pediatrics

618.5261
H.M.

By

Hasan Mohamed Aydoos Aljefri M.B.E.Ch



Under Supervision of

Prof. Dr. Saadeya Mohamed Abdel Fattah

Professor of Pediatrics

Faculty of Medicine-Ain Shams University

Prof. Dr. Sawsan Sayed El-Moselhi

Professor of Pediatrics

Faculty of Medicine-Ain Shams University

60872

Dr. Ahmed M. El-Kahky

Lecturer of Physical Therapy

Med. Dept. Institute of Post Graduate Childhood Studies

Faculty of Medicine - Ain Shams University

Cairo


1999

[Handwritten signature]

[Handwritten signature]

[Handwritten signature]
9/1/10

10



بِسْمِ اللَّهِ الرَّحْمَنِ الرَّحِيمِ

"قالوا سبحانك لا علم لنا إلا

ما علمتنا إنك أنت العليم

الحكيم"

صدق الله العظيم

سورة البقرة

آية ٣٢

ACKNOWLEDGMENT

I would like to express my deepest gratitude and appreciation to my dear **Professor Dr. Saadia Mohamed Abdel Fatah**, professor of Pediatrics, Ain Shams University, for her continuous guidance, expert advice and valuable suggestions that greatly enriched this work.

I am deeply grateful to the kindness of **Dr. Sawsan Sayed El Moselhi**, professor of Pediatrics, Ain Shams University for her keen support, sincere help and valuable efforts throughout this study.

I am much obliged to **Dr. Ahmed M. El Kahky**, Lecturer of Physical therapy, Medical department, Institute of postgraduate Childhood studies, Ain Shams University, for his keen supervision, also for his continuous generous and sincere contributions of this work.

I would like also to thank specially **Dr. Mona Mostafa Othman** Lecturer of Clinical Pathology for her great effort in this study.

LIST OF ABBREVIATIONS

δ	Gamma globulin.
α	Alpha globulin.
β_2/cr	Beta 2 microglobulin creatinine.
$\beta_2\text{MG}$	Beta 2 microglobulin.
ADH	Antidiuretic hormone.
ELISA	Enzyme linked immunosorbent assay.
FA	Fatty acid.
GFR	Glomerular filtration rate.
gm	Gram.
gm/d	Gram per deciliter.
HLA	Human leukocyte antigen.
HMW	High molecular weight.
IgA	Immunoglobulin A.
IgG	Immunoglobulin G.
IgM	Immunoglobulin M.
LMW	Low molecular weight.
m	Meter.
mg	Milligram.
mg/d	Milligram per deciliter.
min.	Minutes.
mmHg	Millimeter mercury.
MW	Molecular weight.

No.	Number of cases.
Pr/cr	Protein creatinine ratio.
RBF	Renal blood flow.
RPF	Renal plasma flow.
ug	Microliter.
ug/ml	Microgram per milliliter.

List of Abbreviations

TABLE OF CONTENTS

Introduction	1
Aim of The Work	3
Review Of Literature	3
Proteinuria	3
Normal handling of proteins by the kidney	7
Protein content of normal urine	9
Pathophysiology of proteinuria	10
Functional proteinuria	12
Diagnosis of proteinuria	19
Effect of proteinuria on renal function	23
Subjects & methods	31
Results	39
Discussion	71
Summary	77
Conclusion	81
Recommendations	83
References	85
Appendix	95
Arabic summary	

LIST OF TABLES

<i>Table</i>	<i>Comment</i>	<i>P</i>
Table 1	Classification of proteinuria.	6
Table 2	Protein composition of normal urine.	9
Table 3	Qualitative tests for urine protein.	21
Table 4	Clinical data of patients and controls.	40
Table 5	Laboratory data of patients and controls.	41
Table 6	Urinary protein creatinine and β_2 microglobulin excretion for all groups before and after exercise.	43
Table 7	Post exercise urinary protein excretion in all patients (group A) in comparison with the control (group B).	49
Table 8	Post exercise urinary creatinine excretion in all patients (G A) in comparison with the control (G B).	52
Table 9	Post exercise urinary protein/creatinine excretion in all patient (G A) in comparison with the control (G B).	55
Table 10	Post exercise urinary β_2 microglobulin in all patients (G A) in comparison with the control (G B).	56

Table	Comment	P
Table 11	Post exercise urinary β_2 MG creatinine ration in all patients (G A) in comparison with the control (G B).	59
Table 12	Comparison between nephrotic patients in relapse, partially treated nephrotic patients and the control group.	60
Table 13	Comparison between nephrotic patients in remission, glomerulonephritis patients in remission and the control group.	62
Table 14	Urine analysis before and after exercise (group A1).	95
Table 15	Urine analysis before and after exercise (group A2).	96
Table 16	Urine analysis before and after exercise (group A3).	97
Table 17	Urine analysis before and after exercise (group A4).	98
Table 18	Urine analysis before and after exercise (group B).	99

List of Tables

LIST OF FIGURES

<i>Figure</i>	<i>Comment</i>	<i>P</i>
Figure 1	Mechanism of post exercise proteinuria.	18
Figure 2	The effect of endocytosis of fatty acid bearing albumin by the proximal tubule.	25
Figure 3	Urinary protein excretion before and after exercise in all groups.	46
Figure 4	Urinary creatinine excretion before and after exercise in all groups.	47
Figure 5	Urinary β_2 MG excretion before and after exercise in all groups.	48
Figure 6	Urinary protein excretion before and after exercise in the control group B and all patients group A.	50
Figure 7	Urinary protein excretion (the differences between pre and post exercise protein excretion in healthy group B and all patients group A).	51
Figure 8	Urinary creatinine excretion before and after exercise in the control group B and all patients group A.	53
Figure 9	Post exercise creatinine excretion in healthy children group B and all patients group A.	54

Figure	Comment	P
Figure 10	Urinary β_2 MG excretion before and after exercise in the control group B and all patients group A.	57
Figure 11	Post exercise β_2 MG excretion in healthy group B and all patients (group A).	58
Figure 12	Post exercise protein excretion (the difference between pre and post exercise) in all patients (group A).	64
Figure 13	Post exercise creatinine excretion in all patients (group A).	65
Figure 14	The correlation between post exercise urinary β_2 microglobulin/creatinine ratio and serum creatinine in the control group.	66
Figure 15	The correlation between post exercise urinary β_2 microglobulin excretion and urinary protein creatinine ratio in glomerulonephritis patients in remission.	67
Figure 16	The correlation between post exercise urinary protein excretion and protein/creatinine ratio in glomerulonephritis patients in remission.	68
Figure 17	The correlation between post exercise protein excretion and serum creatinine ratio in nephrotic patients in remission.	69

List of Figures

Introduction

