
EFFECT OF PROPOFOL-FENTANYL VERSUS ISOFLURANE-FENTANYL ANESTHESIA ON MYOCARDIAL CONTRACTILITY AND PERIPHERAL HEMODYNAMICS DURING CORONARY ARTERY BYPASS GRAFTING



Thesis Submitted for Partial Fulfillment of M.D. Degree in Anesthesiology

Comic Legisland

BY

617.962

Amira Fathy Hefni

M.B.B.Ch., M.Sc., Anesthesiology

57117

Supervised BY

Prof. Dr. Yousri Rubin Ghatas

Professor of Anesthesia and Intensive Care Faculty of Medicine - Ain Shams University

Prof. Dr. Bahira Mohammed Tawfik

Professor of Anesthesia and Intensive Care Faculty of Medicine - Ain Shams University

Dr. Amin Yacin Abd El-Salam

Assistant Professor of Anesthesia and Intensive Care Faculty of Medicine - Ain Shams University


> Faculty of Medicine Ain Shams University *** 1998 ***




الله المحالية

قالوا سبحانك لا علم لنا إلا ما علمتنا إنك أنت العليم الحكيم

صدق الله العظيم





Acknowledgment

First, thanks are all to **God** for blessing me this work until it reached its end, as a little part of his generous help throughout life.

I would like to express my sincere appreciation and deep gratitude to **Prof. Dr. Yousri Rubin Ghatas**, Professor of Anesthesia and Intensive Care, Faculty of Medicine, Ain Shams University, for his kind moral support and continuous encouragement throughout this whole work.

Prof. Dr. Bahira Mohammed Tawfik, Professor of Anesthesia and Intensive Care, Faculty of Medicine, Ain Shams University, for her great support, valuable supervision, precious instructions and encouragement throughout this work.

My deepest appreciation and grateful thanks are due to **Dr. Amin Yacin Abd El-Salam**, Assistant Professor of Anesthesia and Intensive Care, Faculty of Medicine, Ain Shams University, for his sincere help, supervision and cooperation.

AMIR FATHY HEFNY

List of Contents

	PAGE
Introduction	1 – 2
Review of Literature	3 – 71
Aim of the Work	72 – 73
Patients and Methods	74 – 82
Results	83 – 114
Discussion	115 – 127
Conclusion and Recommendations	128
Summary	129 – 131
References	132 – 153
Arabic Summary	1 – 3

List of Figures

No.	Figure	Page
Figure (1):	Characteristics of a sound wave.	4
Figure (2):	Schematic of a modern ultrasound scanner.	6
Figure (3):	M-mode display.	7
Figure (4):	Pulse transmission of image recording and display in the cross-sectional format.	8
Figure (5):	Transducers for esophageal echocardiographic imaging.	9
Figure (6):	Short axis view at the papillary muscle level.	12
Figure (7):	Diagram of loading and trailing edges of the endocardial border.	14
Figure (8):	The end-systolic pressure-volume relaxation is obtained by connecting all the end systolic points measured during a rapid decrease in preload.	19
Figure (9):	Diagram of the end systolic wall stress / rate-corrected velocity of fiber shortening relation.	20
Figure (10):	Circumferential, meridional, and radial components of left ventricular wall stress for an ellipsoid model.	22
Figure (11):	Fentanyl	24
Figure (12):	2,6 di-isopropofol (propofol)	33
Figure (13):	Isoflurane	38
Figure (14):	The definition of coronary dominance	44
Figure (15):	Relative importance of variables that determine myocardial oxygen consumption.	48,
Figure (16):	Lipoid plaque lesion of a coronary artery.	50
Figure (17):	Myocardial oxygen balance between the determinants of myocardial oxygen supply and demand.	52
Figure (18):	Conditions for coronary steal between different areas of the heart and between the subendocardial and subepicardial layers of the left ventricle.	53

Typical plasma concentrations of cardiac enzymes following the onset of chest pain associated with acute myocardial infarction.	62
Typical evolutionary changes of the T wave, ST segment and Q wave following myocardial infarction.	63
Segmental wall motion abnormalities as opposed to normal wall motion.	66
A typical method for defining specific left ventricular segments for wall motion analysis.	67
Normal pressure values and waveforms as a pulmonary artery catheter is advanced from the right atrium to a wedged position in a pulmonary artery.	76
Echocardiogram of the left ventricle at the papillary muscle level. A = LVEDA (left ventricular end-diastolic area).	79
Echocardiogram of the left ventricle at the papillary muscle level A = LVESA (left ventricular end-systolic area).	79
Echocardiogram of the left ventricle at the papillary muscle level. A = the total area enclosed by left ventricular epicardium and the right septum.	8 0
Comparison between P-F group and I-F group as regard velocity of circumferential fiber shortening.	86
Comparison between P-F group and I-F group as regard left ventricular end systolic wall stress.	88
Left ventricular end systolic wall stress versus velocity of circumferential fiber shortening corrected for heart rate relationship obtained in the propofol – fentanyl group.	90
Left ventricular end systolic wall stress versus velocity of circumferential fiber shortening corrected for heart rate relationship obtained in the isoflurane – fentanyl group	90
	following the onset of chest pain associated with acute myocardial infarction. Typical evolutionary changes of the T wave, ST segment and Q wave following myocardial infarction. Segmental wall motion abnormalities as opposed to normal wall motion. A typical method for defining specific left ventricular segments for wall motion analysis. Normal pressure values and waveforms as a pulmonary artery catheter is advanced from the right atrium to a wedged position in a pulmonary artery. Echocardiogram of the left ventricle at the papillary muscle level. A = LVEDA (left ventricular end-diastolic area). Echocardiogram of the left ventricle at the papillary muscle level. A = LVESA (left ventricular end-systolic area). Echocardiogram of the left ventricle at the papillary muscle level. A = the total area enclosed by left ventricular epicardium and the right septum. Comparison between P-F group and I-F group as regard velocity of circumferential fiber shortening. Comparison between P-F group and I-F group as regard left ventricular end systolic wall stress. Left ventricular end systolic wall stress versus velocity of circumferential fiber shortening corrected for heart rate relationship obtained in the propofol – fentanyl group. Left ventricular end systolic wall stress versus velocity of circumferential fiber shortening corrected for heart rate relationship obtained in the propofol – fentanyl group.

Figure (31):	Comparison between P-F group and I-F group as regard heart rate.	92
Figure (32):	Comparison between P-F group and I-F group as regard systolic arterial blood pressure.	94
Figure (33):	Comparison between P-F group and I-F group as regard mean arterial blood pressure.	96
Figure (34):	Comparison between P-F group and I-F group as regard diastolic arterial blood pressure.	98
Figure (35):	Comparison between P-F group and I-F group as regard central venous pressure.	100
Figure (36):	Comparison between P-F group and I-F group as regard.	102
Figure (37):	Comparison between P-F group and I-F group as regard mean pulmonary arterial pressure.	104
Figure (38):	Comparison between P-F group and I-F group as regard pulmonary capillary wedge pressure.	106
Figure (39):	Comparison between P-F group and I-F group as regard coronary perfusion pressure.	108
Figure (40):	Comparison between P-F group and I-F group as regard systemic vascular resistance index.	110
Figure (41):	Comparison between P-F group and I-F group as regard pulmonary vascular resistance index.	112
Figure (42):	Comparison between P-F group and I-F group as regard postoperative results.	114

