


**INTERACTION BETWEEN NICOTINE AND
CAFFEINE ADMINISTRATION
EXPERIMENTAL STUDIES ON BLOOD AND
HAEMATOPOIETIC SYSTEM**

THESIS

**Presented to Department of Zoology, Faculty of Girls
Ain Shams University**



By

AMAL HASSAN AZMY

**Assistant Lecturer in Zoology Department
Faculty of Girls
Ain Shams University
(B.Sc.) 1980**

**Submitted For The Degree of
DOCTOR OF PHILOSOPHY
(In Zoology)**


1994



INTERACTION BETWEEN NICOTINE AND CAFFEINE
ADMINISTRATION: EXPERIMENTAL STUDIES ON BLOOD AND
HAEMATOPOIETIC SYSTEM

SCIENTIFIC SUPERVISION:

Dr. Sanaa M.R. Wahba (Ph. D) Professor of Histology and Histochemistry, Department of Zoology, Faculty of Girls, Ain Shams University, Cairo, Egypt.

A handwritten signature in black ink, reading "Sanaa M.R. Wahba", written over a horizontal line.

Head of Department



ABSTRACT

The present investigation is a trial to throw more light on the side effects of interaction between nicotine and caffeine administration. The study was carried out on blood and haematop-oeitic system including the morphological, haematological and biochemical, histological and bone marrow studies on male albino rats.

LIST OF TABLES

	Page
Table (1): Group distribution.....	58
Table (2): Averages of total body weights in control and experimental groups (grams).....	62
Table (3): Averages of absolute spleen weights in control and experimental groups (grams).	65
Table (4): Averages of relative spleen weights to body weights in control and experimental groups (%).....	68
Table (5): Averages of absolute thymus weights in control and experimental groups (grams).	70
Table (6): Averages of relative thymus weights to body weights in control and experimental groups (%).....	73
Table (7): Averages of R.B.C. counts in control and experimental groups (million/mm ³ blood).	76
Table (8): Averages of haemoglobin content in control and experimental groups (gram/100 ml blood).	78
Table (9): Averages of packed cell volume in control and experimental groups (%).....	81
Table (10): Averages of total leucocytic counts in control and experimental groups (thousand/mm ³ blood).....	83
Table (11): Averages of differential leucocytic counts in control and experimental groups (%).....	86

CONTENTS

	Page
LIST OF TABLES	
LIST OF FIGURES	
INTRODUCTION	1
<u>CHAPTER I: REVIEW OF LITERATURE</u>	4
A. Nicotine Administration:	4
I. Morphological Studies:	4
1. Growth rates and food intake	4
2. General behaviour	11
3. Distribution in tissues	16
II. Haematological and Biochemical Studies	18
III. Histological and Histochemical Studies	29
B. Caffeine Administration	33
I. Morphological Studies	33
1. Growth rates and food intake	33
2. General behaviour	35
3. Distribution in tissues	36
II. Haematological and Biochemical Studies	37
III. Histological and Histochemical Studies	43
C. Interaction Administration	45
<u>CHAPTER II: MATERIAL AND METHODS</u>	49
MATERIAL:	49
1. Experimental Drugs	49
2. Drug Administration	51
3. Experimental Animals	51

CONTENTS

	Page
METHODS:	52
1. Housing of Experimental Animals	52
2. Total Body Weights	52
3. Organ Weights	52
4. Histological Methods	53
5. Haematological Methods	53
a. Evaluation of total erythrocytic count,	54
b. Determination of haemoglobin	
concentration	54
c. Determination of the packed cell	
volume (P.C.V.)	54
d. Evaluation of total leucocytic count ..	54
e. Differential leucocytic count	55
6. Bone Marrow Methods	55
7. Statistical Analysis	55
Experimental Design	58
CHAPTER III: <u>RESULTS</u> :	61
I. Morphological Investigation:.....	61
1. General behaviour.....	61
2. Growth rates.....	62
3. Absolute spleen weights.....	64
4. Relative spleen weights.....	67
5. Averages of absolute thymus weights.....	67
6. Relative thymus weights.....	72

CONTENTS

	Page
II. Haematological Studies:	75
1. Evaluation of total red blood cell count.	75
2. Determination of haemoglobin content	75
3. Determination of P.C.V. (haematocrit value).	80
4. Evaluation of total leucocytic count	80
5. Differential count of leucocytes	85
III. Histological and Histochemical Studies	89
IV. Bone Marrow Studies	156
<u>CHAPTER IV: DISCUSSION</u>	188
I. Morphological Investigation	188
II. Haematolgoical Studies	192
III. Histological and Histochemical Studies	197
IV. Bone Marrow Studies	206
<u>CHAPTER V: SUMMARY</u>	211
<u>BIBLIOGRAPHY</u>	219
<u>ARABIC SUMMARY</u>	

ACKNOWLEDGEMENT

Thank you Allah for your great help to do this work.

No words can express my great thanks and gratitude to my supervisor Dr. Sanaa R. Wahba, Professor of Histology and Histochemistry, Department of Zoology, Faculty of Girls, Ain Shams University, for suggesting and planning this work also her creative guidance and assistance. Without her help the present thesis would not have been accomplished.

I would like also to express my ample thanks for Dr. Madiha M. Ashry, Professor of Histochemistry for her encouragement and continuous interest.

LIST OF FIGURES

	Page
Fig.(1): Average of body weights in grams	63
Fig.(2): Averages of absolute spleen weights in grams	66
Fig.(3): Averages of relative spleen weights to	
body weights.....	69
Fig.(4): Averages of absolute thymus weights in grams	71
Fig.(5): Averages of relative thymus weights to body	
weights.....	74
Fig.(6): Averages of total R.B.C. counts (million/mm ³)	77
Fig.(7): Averages of haemoglobin content (grams/100 ml	
blood).....	79
Fig.(8): Averages of packed cell volume (P.C.V.) %	82
Fig.(9): Averages of total leucocytic counts.....	
(thousands/mm ³).....	84
Fig.(10): Averages of differential leucocytic count of:	
8-a): Lymphocytes.	
8-b): Monocytes.	
8-c): Neutrophils.	
8-d): Eosinophils.....	87
Fig.(11): A section of normal control spleen from rat	
showing white and red pulp. (HX-E, X 60).	91
Fig.(12): A section of normal control spleen from rat	
showing splenic cords within the red pulp.	
(HX-E, X 60).....	91

LIST OF FIGURES

	Page
Fig.(13): Section of spleen from control rat to show normal reticular architecture. (Gomori's technique, X 60).....	92
Fig.(14): Section of spleen from rat after one week of 2 mg/kg nicotine injection showing hyaline material within the congested sinusoids and aggregations of deeply staining lymphocytes within the red pulp. (HX-E, X 100).....	94
Fig.(15): Spleen section of rat treated with 2 mg/kg nicotine for two weeks showing large lymphoid follicle harboring deeply stained lymphocytes and encircled by bands of paler staining epithelial cells. (HX-E, X 100).....	94
Fig.(16): Three weeks of 2 mg/kg nicotine administration showing wedge shaped infarct areas of splenic tissue stuffed with R.B.Cs. (HX-E, X 120)....	95
Fig.(17): Spleen section from rat at the third week post injection with 2 mg/kg nicotine showing thickened trabeculae lined by inflammatory cells. (HX-E, X 60).....	95
Fig.(18): Section of spleen of rat treated with 2 mg/kg nicotine showing necrotic Malpighian corpuscle, small blood vessels and sinusoids within the red pulp with hyalinized walls. (HX-E, X 60).	96

LIST OF FIGURES

	Page
Fig.(19): Spleen section from rat after four weeks of 2 mg/kg nicotine injection to show dilated trabecular venule, filled with inflammatory cells, mainly monocytes. (HX-E, X 100).....	96
Fig.(20): Five weeks of 2 mg/kg nicotine treatment showing thrombosis in the spleen of rat where aggregation of blood platelets, erythrocytes and leucocytes are trapped in solid masses and deposit as eosinophilic material in the vessels walls. (HX-E, X 100).....	98
Fig.(21): Section of spleen from rat treated with 2 mg/kg nicotine for five weeks showing encapsulated Malpighian follicle which attained well-defined boundaries of small compact lymphocytes. (HX-E, X 100).....	98
Fig.(22): Spleen section from rat after six weeks of 2 mg/kg nicotine administration to show necrotic tissue areas with pale staining cytoplasmic cells and some cells showing pyknosis. (HX-E, X 160).....	99
Fig.(23): Section in rat spleen after eight weeks of 2 mg/kg nicotine injection showing signs of recovery in some areas represented by the reappearance of defined Malpighian corpuscle., but still surrounded by degenerative areas. (HX-E, X 60).....	99

LIST OF FIGURES

	Page
Fig.(24): Section of rat spleen after injection with 100 mg/kg caffeine for one week to show degenerative rad pulp, distended and dilated sinuses containing pale staining histiocytes and deeply staining lymphocytes. (HX-E, X 100).....	101
Fig.(25): Rat spleen section post injection with 100 mg/kg caffeine for one week showing infarcted degenerative area demarcated by bands of cells of acute inflammatory exudate. (HX-E, X 100).....	101
Fig.(26): Section of spleen of rat administered 100 mg/kg caffeine after three weeks to show thickenced splenic capsule. (HX-E, X 100).....	102
Fig.(27): Rat spleen section post treatment with 100 mg/kg caffeine for three weeks showing heterogenously weakly stained subcapsular area, necrotic cells with scanty cytoplasm and deeply stained nuclei and pleomorphic cells. (HX-E, X 100).....	102
Fig.(28): Section of rat spleen after four weeks of injection with 100 mg/kg caffeine to show large Malpighian follicle surrounded by concentric mantles of small lymphocytes and hyalinized walls of penecillary arteries. (HX-E, X 60).....	103

- Fig.(29): Spleen section from rat treated with 100 mg/kg caffeine after four weeks to show internal haemorrhage within the red pulp. Red blood cells invaded sinusoidal spaces intermingled with plasma cells and histiocytes. (HX-E, X 100)..... 103
- Fig.(30): After five weeks of injection with 100 mg/kg caffeine, a section of rat spleen showing degenerative changes and necrosis where cells were devoid of cytoplasmic material. (HX-E, X 100)..... 105
- Fig.(31): Section of rat spleen post treatment with 100 mg/kg caffeine for six weeks to show thickened arteriole. The internal lamina was lined by histiocytes. (HX-E, X 100)..... 105
- Fig.(32): Spleen section of rat injected with 100 mg/kg caffeine for six weeks to show detached tissue area within the red pulp where such tissue appeared vacuolated and margined by lymphocytes. (HX-E, X 100)..... 106

LIST OF FIGURES

	Page
Fig.(33): Section of rat spleen after injection with 100 mg/kg caffeine for seven weeks showing thickened trabeculae and degenerative tissue areas. (HX-E, X 60).....	106
Fig.(34): After eight weeks of treatment with 100 mg/kg caffeine, a section of rat spleen showing necrotic areas and dark pigments possibly of haemosiderin invading several parts. (HX-E, X 100).....	107
Fig.(35): Section of spleen of rat treated with the double stress for one week showing faintly stained areas of complete degenerative cells (HX-E, X 100).....	107
Fig.(36): Spleen section of rat treated with the double stress after three weeks showing thickened arteriolar walls invaded by pale eosinophilic stained histiocytes (HX-E, X 240).....	109
Fig.(37): Section of rat spleen after five weeks of treatment with nicotine and caffeine to show haemorrhagic rad pulp areas, separated by broad bands of eosinophilic collagenous fibrous tissue intermixed with the thickened trabeculae (HX-E, X 100).....	109