

Soft Tissue

Management of Soft Tissue Defects of the Lower Third Leg

Lower Third Leg

A comparative study for different modalities of management

Thesis submitted for partial fulfillment of the Master
Degree in General Surgery

Presented by

RASIM MOHAMMED ZAKI SALEM

MB, B.Ph. (1993)

Ain Shams University

6/7.95
B. H

54701

Under the supervision of

PROF. DR. **MUSTAFA MOHAMED HEMEDA**

*Professor of Plastic and Reconstructive Surgery
Faculty of Medicine-Ain Shams University*

DR. **AYMAN ABU EL-MAKAREM SHAKER**

*Lecturer in Plastic and Reconstructive Surgery
Faculty of Medicine-Ain Shams University*

FACULTY OF MEDICINE
AIN SHAMS UNIVERSITY
1998








بِسْمِ اللَّهِ الرَّحْمَنِ الرَّحِيمِ
قَالُوا سُبْحَانَكَ لَا عِلْمَ لَنَا إِلَّا مَا عَلَّمْتَنَا
إِنَّكَ أَنْتَ الْعَلِيمُ الْحَكِيمُ

صدق الله العظيم
(سورة البقرة الآية ٣٢)

...

Dedication.....

*In Memory of my
Father with
Appreciation*



*To my Mother with
Love and
Gratitude*

Acknowledgment

Thanks to God firstly and lastly.

*I wish to offer my profound gratitude and acknowledgment to my supervisor Prof. Dr. **Mustafa M. Hemeda**, Professor and Head of the Plastic and Reconstructive Surgery Department, Ain Shams University, for his continuous guidance, kindness in accepting to pursue this work and unfailing support. I will always remain indebted.*

*I wish also to express my sincere appreciation to Dr. **Ayman A. Shaker** for his great help, assistance and unique cooperation throughout this study.*

*I also thank all members of the Plastic and Reconstructive Surgery Department at Ain Shams University for their great help with special thanks to Dr. **S. El-Fayoumy**, Dr. **H. Maher**, Dr. **A. Hanafy** and Dr. **S. Nasser**.*

Basim Zaki
1998

List of Contents

	<i>Page No.</i>
<i>Introduction and Aim of the Work</i>	<i>1-3</i>
<i>Review of Literature</i>	<i>4-137</i>
I) <i>Anatomy of the Leg</i>	<i>4-37</i>
<i>Vascular supply of the leg</i>	
<i>Nerve supply of the leg</i>	
<i>Vascular supply of the skin of the leg</i>	
II) <i>Pathophysiology of Leg Ulcers</i>	<i>38-54</i>
III) <i>Methods of Reconstruction of Soft Tissue Defects of the Lower 1/3 of the Leg</i>	<i>55-137</i>
<i>Reconstruction by primary approximation</i>	<i>57</i>
<i>Reconstruction by skin grafting</i>	<i>58</i>
<i>Reconstruction by skin flaps</i>	<i>61</i>
<i>Reconstruction by muscle flaps</i>	<i>65</i>
<i>Reconstruction by fasciocutaneous flaps</i>	<i>86</i>
<i>Reconstruction by adipofascial flaps</i>	<i>113</i>
<i>Reconstruction by distant flaps</i>	<i>119</i>
<i>Reconstruction by free tissue transfer</i>	<i>124</i>
<i>Materials and Methods</i>	<i>138-148</i>
<i>Results</i>	<i>149-156</i>
<i>Discussion</i>	<i>157-169</i>
<i>Summary and Conclusion</i>	<i>170-174</i>
<i>References</i>	<i>175-199</i>
<i>Arabic Summary</i>	

List of Figures

Fig. (1): Deep fascia of the leg.

Fig. (2): Variations in the manner of division of popliteal artery.

Fig. (3): Branches of the popliteal and posterior tibial arteries.

Fig. (4): Posterior tibial artery.

Fig. (5): Peroneal artery.

Fig. (6): Anterior tibial artery.

Fig. (7): Branches of the anterior tibial artery.

Fig. (8): Variations of the source arteries of the leg.

Fig. (9): Routes by which vessels reach the skin.

Fig. (10): Direct cutaneous system of vessels.

Fig. (11): Fasciocutaneous system.

Fig. (12): Schematic representation of a septocutaneous pedicle.

Fig. (13): Musculocutaneous system.

Fig. (14): Byrd's Classification of compound fractures of the leg.

Fig. (15): Recommended muscles for distal third leg coverage and reach of these flaps.

Fig. (16): Soleus vascular anatomy.

Fig. (17): Distally based hemisoleus muscle flap.

Fig. (18): Flexor digitorum longus muscle flap

Fig. (19): Peroneus brevis muscle flap.

Fig. (20): Extensor hallucis longus muscle flap.

Fig. (21): Flexor hallucis longus muscle flap.

Fig. (22): Transverse section through the lower third of

the leg.

Fig. (23): Vasculature near the skin.

Fig. (24): Distally based peroneal artery fasciocutaneous flap.

Fig. (25): Reverse flow anterior tibial artery fasciocutaneous flap.

Fig. (26): Distally based posterior tibial artery fasciocutaneous flap.

Fig. (27): Reversed flow posterior tibial artery fasciocutaneous flap.

Fig. (28): A Marjolin ulcer.

Fig. (29): Elevation of a soleus muscle flap.

Fig. (30): Cross leg musculocutaneous flap.

Fig. (31): Application of a free latissimus dorsi myocutaneous flap.

Fig. (32): Rectus abdominus free muscle transfer.

Fig. (33): Large leg defect after failed free flap.

Fig. (34): STSG to the medial malleolus in a patient with chronic venous ulcer (one week post-operative).

Fig. (35): STSG one month post-operative showing partial graft loss.

Fig. (36): STSG applied to the medial malleolus (one month post-operative).

Fig. (37): STSG in a patient with chronic leg ulcers showing a near total graft loss 1 month post-operative.

Fig. (38): Compound fractures both bones leg.

Fig. (39): Elevation of the soleus muscle after debridement.

Fig. (40): The soleus muscle in place two weeks postoperative.

Fig. (41): Skin grafted soleus muscle two weeks after

