

# **THE ROLE OF ULTRASONOGRAPHY IN ACUTE PELVIC PAIN IN FEMALES**

**ESSAY**

**Submitted in Partial Fulfilment of Master Degree  
in Radio Diagnosis**

**By**

**ENAS AHMED AZAB**

**M.P., B.Ch.**

**SUPERVISOR**


**Prof. Dr. SALWA T. AHMED ISMAIL**

Prof. of Radio Diagnosis  
Faculty of Medicine  
Ain Shams University

*Faculty of Medicine  
Ain Shams University*

**1994**





## **ACKNOWLEDGEMENT**

*I am deeply indebted to Prof. Dr. SALWA T. AHMED ISMAIL, Professor of Radiology, Faculty of Medicine, Ain Shams University for her unlimited support, patience, guidance and spirit of cooperation which provide me with great help and assistance to continue this work.*

*My sincere gratitude for all my Professors and Colleagues of Radiodiagnosis Departments, Faculty of Medicine, Ain Shams University for their kind help and support they offered.*

*Finally, no words can express the warmth of my feelings to my Father and Mother for their continuous encouragement, my Husband for sincere help and also everyone involved in making contributions to this work.*



## LISTS OF FIGURES

|           | Title                                                                            | Page |
|-----------|----------------------------------------------------------------------------------|------|
| Fig. (1)  | The uterus and left broad ligament                                               | 5    |
| Fig. (2)  | The left uterine and ovarian arteries                                            | 8    |
| Fig. (3)  | Parasagittal single sweep of the midline.                                        | 12   |
| Fig. (4)  | Real-time sector transverse scan                                                 | 14   |
| Fig. (5)  | Transverse sweep of the pelvis through the mid portion of the uterus.            | 16   |
| Fig. (6)  | Limitations of transabdominal scanning                                           | 22   |
| Fig. (7)  | Limitations of transvaginal scanning                                             | 23   |
| Fig. (8)  | Different transvaginal transducers                                               | 25   |
| Fig. (9)  | Transvaginal image display options                                               | 25   |
| Fig. (10) | TVS cervical canal and lower uterine segment                                     | 30   |
| Fig. (11) | TVS Endometrium of uterine fundus                                                | 32   |
| Fig. (12) | Examination of the ovaries                                                       | 34   |
| Fig. (13) | Tubo-ovarian abcess                                                              | 49   |
| Fig. (14) | Transverse sonogram of a patient with severe chronic pelvic inflammatory disease | 49   |
| Fig. (15) | Endometritis                                                                     | 51   |
| Fig. (16) | Acute salpingitis                                                                | 51   |
| Fig. (17) | Hydrosalpinx                                                                     | 53   |
| Fig. (18) | Chronic hydrosalpinx                                                             | 53   |
| Fig. (19) | Pyosalpinx                                                                       | 55   |
| Fig. (20) | Tubo-ovarian abcess                                                              | 57   |
| Fig. (21) | Endometrioma                                                                     | 61   |
| Fig. (22) | Endometrioma                                                                     | 61   |



|                 | Title                                                          | Page |
|-----------------|----------------------------------------------------------------|------|
| Fig. (23)       | Ruptured dermoid                                               | 63   |
| Fig. (24)       | An unusual ectopic pregnancy                                   | 66   |
| Fig. (25)       | An intrauterine fluid collection                               | 66   |
| Fig. (26)       | Ectopic pregnancy with an interauterine decidual cast          | 68   |
| Fig. (27)       | An extrauterine gestational sac                                | 68   |
| Fig. (28)       | Tubo-ovarian abcess                                            | 70   |
| Fig. (29)       | An echogenic clot                                              | 72   |
| Fig. (30)       | Pseudogestational sac                                          | 77   |
| Fig. (31)       | Pseudogestational sac versus non viable intrauterine pregnancy | 79   |
| Fig. (32)       | Living ectopic pregnancy                                       | 82   |
| Fig. (33)       | Extrauterine gestational sac                                   | 82   |
| Fig. (34)       | Extrauterine gestational sac                                   | 84   |
| Fig. (35)       | Ectopic pregnancy with hematosalpinx                           | 84   |
| Fig. (36)       | Ectopic pregnancy with hematosalpinx                           | 86   |
| Fig. (37)       | Hemoperitoneum                                                 | 88   |
| Fig. (38)       | Case No. 1: right sided. hydrosalpinx                          | 95   |
| Fig. (39)       | Case No. 2: moderately-sized right hydrosalpinx.               | 97   |
| Fig. (40)       | Case No. 3: Right-sided tubo-ovarian abcess                    | 99   |
| Fig. (41)       | Case No. 4: Pelvic abcess                                      | 101  |
| Fig. (42)       | Case No. 5: cul-de-sac abcess                                  | 103  |
| Fig. (43)       | Case No.6: Cul-de-sac abcess associated with IUD               | 105  |
| Fig. (44), (45) | Case No.7: Rapidly progressed pelvic inflammatory disease      | 108  |





|                 | <b>Title</b>                                                   | <b>Page</b> |
|-----------------|----------------------------------------------------------------|-------------|
| Fig. (46)       | Case No. 8: Pelvic abcess secondary to diverticular disease    | 110         |
| Fig. (47)       | Case No. 9: Pelvic abcess secondary to ulcerative colitis      | 110         |
| Fig. (48)       | Case No. 10: Drainage of tubo-ovarian abcess                   | 112         |
| Fig. (49)       | Case No. 11: bilateral endometrioma                            | 114         |
| Fig. (50)       | Case No. 12: Pelvic endometrioma                               | 116         |
| Fig. (51)       | Case No. 13: Dermoid cyst                                      | 118         |
| Fig. (52)       | Case No. 14: Interstitial ectopic pregnancy                    | 120         |
| Fig. (53)       | Case No. 15: Pseudogestational sac                             | 122         |
| Fig. (54)       | Case No. 16: Pseudogestational sac with echogenic fluid        | 124         |
| Fig. (55)       | Case No. 17: Pseudogestational sac simulating embryonic demise | 126         |
| Fig. (56), (57) | Case No. 18, 19: Ovarian Torsion                               | 128         |



## LISTS OF DIAGRAMS

|           | Title                                                                | Page |
|-----------|----------------------------------------------------------------------|------|
| Diag. (1) | Transverse scan of the pelvis                                        | 14   |
| Diag. (2) | Normal anteverted uterus                                             | 18   |
| Diag. (3) | Approach in transvaginal scanning                                    | 27   |
| Diag. (4) | Suggested algorithm for evaluating suspected ectopic pregnancy       | 75   |
| Diag. (5) | Drawing of the cul. dc-sac finding associated with ectopic pregnancy | 86   |



## LISTS OF TABLES

|           | Title                                                                                                 | Page |
|-----------|-------------------------------------------------------------------------------------------------------|------|
| Table (1) | Test performance of transvaginal ultrasound examination in the diagnosis of plasma cell endometritis. | 59   |
| Table (2) | Findings from transvaginal sonography in 47 ectopic tubal pregnancies.                                | 89   |

