

**EFFECT OF SUBCURATIVE DOSE AND REPEATED
EXPOSURE TO CERCARIAE OF *SCHISTOSOMA*
MANSONI ON TREATMENT OF MICE
WITH PRAZIQUANTEL**


By

SAMIA WILLIAM ABDEL-SAYED

B. Sc. (Zoology), Ain Shams University

Theodor Bilharz Research institute

591.0724
S.W



A THESIS

Submitted in Partial Fulfillment of
the Requirement for the Degree of
Master of Science

48269

In
Zoology

Department of Zoology
Faculty of Science
Ain Shams University

[Signature]



1993

SUPERVISORS

Dr. Nawal Nagib Nashed,

Assistant Professor , Zoology department


Faculty of Science

Ain Shams University

Prof. Dr. Hussein Mohamed Shoukry El-Nahal,

Head of Parasitology department

Theodor Bilharz Research Institute



Prof. Dr. Aisha Metwally,

Head of Pharmacology department

Theodor Bilharz Research Institute



TO

THE

LIBRARY OF THE

CONTENTS

	Page
- Introduction.....	1
- Aim of work.....	4
- Review of Literature.....	5
- Chemotherapy of Schistosomiasis.....	5
- Classification of antischistosomal drugs...	5
. Antimony compounds.....	5
. Non-antimony compounds.....	8
. Praziquantel (Biltricide).....	16
- Drug resistance.....	32
- Resistance to reinfection in schistosomiasis mansoni.....	36
- Pathology of liver in schistosomiasis.....	49
- Materials & Methods.....	55
- Results.....	75
- Discussion.....	138
- Summary.....	159
- References.....	165
- Arabic summary.....	

ACKNOWLEDGMENT

I am most grateful to Dr. Nawal Nagib Nashed, Assistant. Professor of Zoology department, Ain Shams university, for her initiating power, stimulating suggestions, and kind help and advice.

I wish to express my profound gratitude and sincere appreciation to prof. Dr. Hussein Mohamed Shoukry El-Nahal for his instructing guidance and valuable suggestions and supervision.

It is difficult to translate my feelings towards prof. Dr. Aisha Metwally who sacrificed a good deal of her valuable time to help me in preparation of the thesis.

I would like to thank Dr. Afkar Abdel-Ganay for kind help her valuable advises.

In fact, I owe much to the staff of the parasitology department and pharmacology department in Theodor Bilharz Institute.

LIST OF TABLES

	Page
Table 1: Effect of praziquantel on worm distribution in albino mice infected with 50 <u>S. mansoni</u> cercariae.....	76
Table 2: The effect of praziquantel on oogram in small intestine in albino mice infected with 50 <u>S. mansoni</u> cercariae.....	83
Table 3: The effect of praziquantel on the number of ova/gm tissues in albino mice infected with 50 <u>S. mansoni</u> cercariae.....	87
Table 4: Number of liver granulomas in 5 successive power field (10x 10).....	91
Table 5: Size of liver granuloma in relation to different doses of praziquantel therapy....	92
Table 6: The effect of praziquantel (2 x 500 mg/kg b.w) on worm distribution in albino mice infected with different doses of <u>S. mansoni</u> cercariae.....	101

LIST OF TABLES (Cont'd)

	Page
Table 7: The effect of praziquantel (2 x 500 mg/kg b.w) on oogram in small intestine in albino mice infected with different doses of <u>S. mansoni</u> cercariae.....	107
Table 8: The effect of praziquantel (2 x 500 mg/kg b.w) on the number of ova/gm liver and intestine in mice infected with different doses of <u>S. mansoni</u> cercariae...	109
Table 9: The effect of praziquantel (2 x 500 mg/kg b.w) on worm distribution in albino mice infected with different doses of <u>S. mansoni</u> cercariae given in a single dose or by repeated infection.....	113
Table 10: The effect of praziquantel (2 x 500 mg/kg b.w) on oogram in albino mice infected with different doses of <u>S. mansoni</u> cercariae given in a single dose or by repeated infection.....	116

LIST OF TABLES (Cont'd)

	Page
Table 11: The effect of praziquantel (2 x 500 mg/kg b.w) on the number of ova/gm tissues in albino mice infected with different doses of <u>S. mansoni</u> cercariae given in a single dose or by repeated infection..	118
Table 12: The effect of praziquantel (2 x 500 mg/kg b.w) on worm distribution in albino mice infected repeatedly with <u>S. mansoni</u> cercariae.....	120
Table 13: The effect of praziquantel (2 x 500 mg/kg b.w) on oogram in small intestine in albino mice infected repeatedly with <u>S. mansoni</u> cercariae.....	125
Table 14: The effect of praziquantel (2 x 500 mg/kg b.w) on the number of ova/gm tissues in mice infected repeatedly with <u>S. mansoni</u> cercariae.....	128

LIST OF FIGURES

	Page
(Fig.1): Experimental design of the work....	56
(Fig.2): Experimental design for single infection of mice with 50 <u>S. mansoni</u> Cercariae.....	57
(Fig.3): Experimental design for repeated infection of mice.....	60
(Fig.4): Oogram picture showing Immature eggs of the 1 st (A) and 2 nd (B) stages and mature ova (E) of <u>S. mansoni</u> (X 56)...	66
(Fig.5): Oogram picture showing Immature eggs of the 2 nd (B) and 3 rd (C) stages of <u>S. mansoni</u> (X 56).....	67
(Fig.6): Oogram picture showing Immature eggs of the 3 rd (C) stage, mature ova (E) and Dead egg of <u>S. mansoni</u> (X 56).....	68
(Fig.7): Oogram picture showing Immature eggs of the 4 th stage (D) of <u>S. mansoni</u> (X 56)	69

LIST OF FIGURES (Cont'd)

	Page
<p>(Fig.8): - Oogram picture showing dead egg (F) of <u>S. mansoni</u> (granular, darkened) and mature ova (E) of <u>S. mansoni</u> (X 56)....</p>	70
<p>(Fig 9): The effect of praziquantel on total number of schistosomes in albino mice infected with 50 <u>S. mansoni</u> cercariae..</p>	78
<p>(Fig 10): The effect of praziquantel on number of <u>schistosoma mansoni</u> worms in liver of mice treated with different regimens of praziquantel.....</p>	79
<p>(Fig 11): The effect of praziquantel on oogram in small intestine in albino mice infected with 50 <u>S. mansoni</u> cercariae.....</p>	81
<p>(Fig.12): The effect of praziquantel on the number of ova/gm liver in albino mice infected with 50 <u>S. mansoni</u> cercariae.....</p>	84

LIST OF FIGURES (Cont'd)

	Page
<p>(Fig 13): The effect of praziquantel on the number of ova/gm intestine in albino mice infected with 50 <u>S. mansoni</u> cercariae....</p>	85
<p>(Fig.14): Hepatic granuloma formed of ova surrounded by fibroblasts, collagen fibers with other inflammatory cells in a mice infected for 14 weeks. Masson Trichrome stain. (X 200).....</p>	88
<p>(Fig.15): Portal tract distended by multiple granulomas, collagen fibres and inflammatory cells in a mice infected for 14 weeks. Masson Trichrome stain (X 128).....</p>	90
<p>(Fig.16): Very small sized hepatic granulomas in a mice treated by 2 full doses of praziquantel and sacrificed at the 14th weeks after infection, consisting of ova shell surrounded by collagen fibers and pigmented macrophage cells. Masson Trichrome stain. (X 200).....</p>	93

LIST OF FIGURES (Cont'd)

	Page
<p>(Fig.17): A thin portal tract containing fibrotic granulomas and collagen fibers with pigmented macrophage cells in a mice treated by 2 full doses of praziquantel and sacrificed 14 weeks after infection. Masson Trichrome stain (X 128).....</p>	94
<p>(Fig.18): A worm granuloma consisting of dead worm surrounded by inflammatory cells and collagen fibers in a mice treated by praziquantel. HX and Eosin stain (X 128).....</p>	95
<p>(Fig.19): Fibrotic granuloma in a mice treated by full dose of praziquantel and sacrificed 14 weeks after infection. Masson Trichrome stain. (X 200).....</p>	97
<p>(Fig.20): Fibrocellular granuloma in a mice treated by half curative dose of praziquantel followed by the curative dose.</p> <p>Masson Trichrome stain (X 200).</p>	98

LIST OF FIGURES (Cont'd)

	Page
(Fig.21): Fibrocellular granuloma in a mice treated by half the curative dose of praziquantel. Masson Trichrome stain. (X 200).....	100
(Fig 22): The effect of praziquantel (2 x 500 mg/kg b.w) on total number of schistosomes in albino mice infected with different doses of <u>S. mansoni</u> cercariae.....	103
(Fig 23): The effect of praziquantel (2 x 500 mg/kg b.w) on number of schistosomes worm in liver of mice infected with different doses of <u>S. mansoni</u> cercariae.....	104
(Fig 24): The effect of praziquantel (2 x 500 mg/kg b.w) on oogram in small intestine in albino mice infected with different doses of <u>S. mansoni</u> cercariae.....	105
(Fig 25): The effect of praziquantel (2 x 500 mg/kg b.w) on the number of ova/gm liver in albino mice infected with different doses of <u>S. mansoni</u> cercariae.....	110

LIST OF FIGURES (Cont'd)

	Page
(Fig 26): The effect of praziquantel (2 x 500 mg/kg b.w) on the number of ova/gm intestine in albino mice infected with different doses of <u>S. mansoni</u> cercariae.....	111
(Fig 27): The effect of praziquantel (2 x 500 mg/kg b.w) on total number of schistosomes in albino mice infected repeatedly with <u>S. mansoni</u> cercariae.....	122
(Fig 28): The effect of praziquantel (2 x 500 mg/kg b.w) on number of schistosomes worm in liver of mice infected repeatedly with <u>S. mansoni</u> cercariae.....	123
(Fig 29): The effect of praziquantel (2 x 500 mg/kg b.w) on oogram in small intestine in mice infected repeatedly with <u>S. mansoni</u> cercariae.....	126
(Fig 30): The effect of praziquantel (2 x 500 mg/kg b.w) on the number of ova/gm liver in mice infected repeatedly with <u>S. mansoni</u> cercariae.....	130