

**GEOLOGICAL STUDIES ON THE AREA
AROUND GEBEL KULYIET, SOUTH
EASTERN DESERT**


By
ZEINHOM SAYED AHMED EL ALFY
B. Sc. in Geology

A THESIS
Submitted in Partial Fulfilment of
The Requirements for the Degree of
MASTER OF SCIENCE

in
GEOLOGY

Department of Geology
Faculty of Science
Ain Shams University

1986



TO
WHOM I AM TAUGHT BY



ACKNOWLEDGEMENTS

The present work is carried out in the Department of Geology, Ain Shams University, for such assistance, the writer wishes to express Deep thank for the staff members, who kindly offered laboratory facilities and unfailing help during the progress of the investigations.

The writer would like to express his deep gratitude to Prof. Dr. M. Y. Meneisy, Geology Department, Ain Shams University, (Now head of Geology Department, Faculty of Science, Qatar University), for his supervision, continuous guidance and fruitful discussions during the progress of this work.

The writer gratefully thanks Dr. Bahay Issawi, The Oil Vice Minister, for his supervision and encouragement.

Grateful thanks are also due to Dr. B. A. El Kaluobi, Geology Department, Ain Shams University, for his direct supervision, help and advice in each stage of the present work.

Chemical analyses were carried out in the laboratory of Institute Mineralogy and Petrology, Mainz University, for such assistance, the writer feels indebted.

The writer wishes to express his deep thanks for the Egyptian Geological Survey and Mineral projects Authority, who provided facilities during field studies.

I am also grateful to geologists Tarek Y. S. and Amir M. S., who typed the manuscript at various stages.

N O T E

The present thesis is submitted to Ain Shams University in partial fulfillment of the requirements for the Degree of Master of Science in Geology.

Besides the research work materialized in this thesis, the Candidate has attended eight graduate courses for one year in the following topics:

1. Field geology.
2. Laboratory techniques.
3. Mineralogy.
4. Crystallography.
5. Igneous petrology.
6. Metamorphic petrology.
7. Sedimentary petrology.
8. Geochemistry.

He has successfully passed the final examinations in these courses.

In fulfillment of the language requirements of the degree, he has also passed the final examination of a course in English language.

Prof. Dr.

Monamed A. Bassyuni
Head, Department of Geology

LIST OF CONTENTS

ABSTRACT

CHAPTER I	INTRODUCTION	1
1.1.	Previous Work	4
1.2.	Scope and Object of the Present Study	11
CHAPTER II	GEOLOGIC SETTING	13
II.1.	Volcanogenic Metasediments	15
II.1.1.	Phyllites	18
II.1.2.	Schists	20
II.2.	Metavolcanics and Associated Metapyroclastics	22
II.2.1.	Metapyroclastics	23
II.2.2.	Metatuffs	25
II.2.	Metaandesite	25
II.2.4.	Metadacite-rhyodacite Association	26
II.2.5.	Metarhyolite	26
II.3.	Metagabbro-diorite	27
II.4.	Syntectonic to Late Tectonic Granodiorite and Tonalite	30
II.5.	Younger Granites	30
II.5.1.	Seiga Granite	31
II.5.2.	Kulyiet Granite	31
II.6.	Post granitic Dykes and Veins	32
	General Structural Features in the Area	33

CHAPTER III	PETROGRAPHY	56
III.1.	Metasediments	56
III.1.1.	Phyllite	57
III.1.1.1.	Quartz Phyllite	57
III.1.1.2.	Chlorite Phyllite	59
III.1.1.3.	Biotite Phyllite (phyllitic schist)	59
III.1.2.	Schists	42
III.1.2.1.	Chlorite-biotite Schist	43
III.1.2.2.	Quartz-rich Schist	43
III.1.2.3.	Quartz-muscovite Schist	45
III.1.2.4.	Mica Schists	45
	A. Biotite Schist	45
	B. Muscovite-biotite Schist	46
III.1.2.5.	Garnetiferous-mica Schist	50
III.1.2.6.	Amphibole-bearing Schists	50
	A. Chlorite-actinolite Schist	52
	B. Biotite-hornblende Schist	52
	C. Hornblende Schist	53
SOURCE ROCKS AND PHYSICAL CONDITION OF METAMORPHISM		56
III.2.	Metavolcanics	58
III.2.1.	Metaandesite	58
III.2.2.	Metadacite-rhyodacite	62
III.2.3.	Metarhyolite	63
III.2.3.1.	Porphyritic Metarhyolite	63
III.2.3.2.	Mylonitized Metarhyolite	67
III.2.4.	Metapyroclastics and Tuffs	68
III.3.	Metagabbro-diorite Complex	75
III.4.	Syntectonic Plutonites (Tonalite, Granodiorite)	78
III.4.1.	Tonalite	78
III.4.2.	Granodiorite	80
III.4.3.	Myrmekitic Granodiorite	82

III.5.	Younger Granitoids	83
III.5.1.	Biotite Granite	83
III.5.2.	Muscovite Granite	84
III.6.	Post-Granite Dykes and Veins	86
III.6.1.	Basic Dykes	86
III.6.2.	Acidic Dykes	86
III.6.2.1.	Granophyre	86
III.6.2.2.	Porphyritic Rhyolite	87
III.6.2.	Trachyte	87
CHAPTER IV GEOCHRONOLOGY OF KULYIET GRANITE		90
Laboratory Techniques and Results		91
DISCUSSION		91
CHAPTER V PETROCHEMISTRY		94
V.1.	Petrochemistry of Metavolcanics	95
V.1.1.	Variation in Chemical Composition	102
	The Alkalies-Silica Diagrams	102
	FeO^t/MgO Versus SiO_2 , TiO_2 and FeO Diagram	103
	Al_2O_3 Versus Normative Plagioclase Diagram	105
	Variation of P_2O_5 Versus TiO_2 Diagram	106
V.1.2.	Ternary Variation Diagrams	106
	The Or-Ab-An Diagram	108
	The Q-Or-Ab Diagram	111
	AFM Diagram	111
V.1.3.	Classification of Metavolcanics	111
V.1.4.	Trace Elements	112
V.2.	Petrochemistry of Granites	116
V.2.1.	Chemical Features of The Studied Granites	129

V.2.2.	Magma Type	134
	a. Rittmann Suite Index	136
	b. Wright's Alkalinity Ratio	137
	c. K-C-N Ternary Diagram	139
	d. AFM Ternary Diagram	139
V.2.3.	Classification	144
V.2.4.	Distribution and Behaviour of Trace Elements	146
CHAPTER VI SUMMARY AND CONCLUSION		153
ACQUAINTANCESHIP		158
REFERENCES		159

LIST OF FIGURES

Fig. 1	Location map.	2
Fig. 2	Geologic map of the area around Gabal Kulyiet, South Eastern Desert, Egypt.	
Fig. 3	General view of the eastern side of Gabal Kulyiet, and the surrounding low hilly land metasediments and metavolcanics.	3
Fig. 4	Hunting photogeological map (1967).	8
Fig. 5	Metasediments (south of Gabal Seiga) northern part of the area showing marked foliation parallel to bedding.	19
Fig. 6	Sharp contact between granodiorite and metasediments north east of Gabal Kulyiet.	19
Fig. 7	General view of strongly sheared and foliated metatuffs.	24
Fig. 8a	Mylonitized metarhyolite.	29
Fig. 8b	Foliated and folded metarhyolite.	29
Fig. 9a	Fine grained phyllite consisting of quartz grains and wispy seams of chlorite matrix parallel to a faint foliation, and scattered magnetite. Notice the formation of magnetite. (P.P. Light, $\times 65$).	38

- Fig. 9b Fine grained phyllite showing chlorite (dark) in parallel and subparallel arrangement with quartz matrix and quartz veinlets. (C.N. Light, $\times 65$). 38
- Fig. 10 Chlorite phyllite with bands of calcite. (C.N. Light 6.3×10). 40
- Fig. 11a Phyllitic schist showing the formation of large biotite crystals in the fine grained phyllitic groundmass. (P.P. Light 4×10). 41
- Fig. 11b The formation of large biotite crystals in the fine grained phyllitic groundmass. (C.N. Light 4×10). 41
- Fig. 12a Chlorite biotite schist consisting of fine grained biotite and chlorite with large spots of biotite flakes. (P.P. Light 4×10). 42
- Fig. 12b Chlorite biotite schist consisting of fine grained biotite and chlorite with large spots of biotite flakes. (C.N. Light 4×10). 42
- Fig. 13 Quartz-rich schist with abundant quartz grains and minor mats of muscovite, flakes of chlorite and calcite crystals that appear between the quartz. (C.N. Light 2.5×6.3). 47
- Fig. 14 Quartz muscovite schist showing alternating thin bands of quartz lenticles and fine muscovite and sericite. (C.N. Light 2.5×6.3). 47

Fig. 15a	Mica schist showing pronounced formation of S planes. (P.P. Light 2.5×6.3).	48
Fig. 15b	Mica schist showing pronounced formation of S planes. (C.N. Light 2.5×6.3).	48
Fig. 16a	Muscovite biotite schist with formation and rotation of mica flakes that define early S_1 and later incipient S_2 . (C.N. Light 2.5×6.3).	49
Fig. 16b	Muscovite biotite schist with formation of S_1 and S_2 planes. (P.P. Light 2.5×6.3).	49
Fig. 17a	Garnetiferous mica schist showing euhedral garnet crystals with rhombdodecahedral form set in schistose aggregates of biotite and quartz in parallel alignment. (C.N. Light 2.5×6.3).	51
Fig. 17b	Garnetiferous mica schist showing garnet crystals with rhombdodecahedral form, biotite flakes and quartz grains. (C.N. Light 2.5×6.3).	51
Fig. 18a	Biotite hornblende schist showing intergrowth of biotite and hornblende. (P.P. Light 2.5×6.3).	54
Fig. 18b	Biotite hornblende schist showing intergrowth of biotite and hornblende. (C.N. Light 2.5×6.3).	54

Fig. 19a	The development of hornblende in two S directions in hornblende schist. (P.P. Light 2.5×6.3).	55
Fig. 19b	The development of elongate prismatic hornblende crystals in two S directions in hornblende schist. (C.N. Light 2.5×6.3).	55
Fig. 20a	Metaandesite showing partially altered plagioclase phenocrysts embedded in fine grained groundmass. (P.P. Light 2.5×6.3).	60
Fig. 20b	Porphyritic metaandesite showing partially altered plagioclase phenocrysts embedded in fine grained groundmass consisting of quartz, plagioclase, chlorite and actinolite. (C.N. Light 2.5×6.3).	60
Fig. 21	Porphyritic metaandesite showing plagioclase phenocrysts embedded in fine grained groundmass of quartz, mafics and iron oxides. (P.P. Light 2.5×6.3).	61
Fig. 22	Porphyritic metaandesite showing euhedral twinned plagioclase phenocrysts. (C.N. Light 2.5×6.3).	65
Fig. 23	Porphyritic metadacite. (C.N. Light 2.5×6.3).	65
Fig. 24	Showing segregation of quartz and muscovite rich layers. Quartz layers show deformation, and shearing that produce long attenuated lenses. (C.N. Light 2.5×6.3).	66

Fig. 25a	Quartz-augen porphyroclast in fine grained matrix of quartz and muscovite. (P.P. Light 2.5×6.3).	70
Fig. 25b	Quartz-augen porphyroclast in fine grained matrix of quartz and muscovite. (C.N. Light 2.5×6.3).	70
Fig. 26a	Lapilli tuffs. (P.P. Light 2.5×6.3).	71
Fig. 26b	Lapilli tuffs. (C.N. Light 2.5×6.3).	71
Fig. 27a	Lithic metatuffs showing rock fragments of schistose metasediments and quartz crystals embedded in coarse grained tuffaceous matrix. (P.P. Light 2.5×6.3).	72
Fig. 27b	Lithic metatuffs showing rock fragments of schistose metasediments and quartz crystals embedded in coarse grained tuffaceous matrix. (C.N. Light 2.5×6.3).	72
Fig. 28a	Very fine metatuffs consist of rock fragments, iron oxide grains enclosed in a very fine grained banded matrix. (P.P. light 2.5×6.3).	73
Fig. 28b	Fine grained metatuffs with interveining quartz laminae. (P.P. Light 2.5×6.3).	73
Fig. 29a	Lithic and crystal metatuffs showing euhedral igneous phenocrysts of feldspars in clastic matrix. (P.P. Light 2.5×6.3).	74

Fig. 29b	Lithic and crystal metatuffs showing euhedral igneous phenocrysts of feldspars in clastic matrix. (C.N. Light 2.5×6.3).	74
Fig. 30a	Metagabbro with plagioclase, hornblende and altered pyroxene. (P.P. Light 2.5×6.3).	77
Fig. 30b	Metagabbro with plagioclase, hornblende and altered pyroxene. (C.N. Light 2.5×6.3).	77
Fig. 31	Tonalite Consisting of plagioclase, hornblende and biotite with interstitial quartz. (C.N. Light 2.5×6.3).	79
Fig. 32	Granodiorite consisting of plagioclase, alkali feldspar and deformed quartz with mafics. (C.N. Light 2.5×6.3).	81
Fig. 33	Biotite granite showing deformed quartz and small crystals of kinked plagioclase enclosed in deformed alkali feldspar. (C.N. Light 2.5×6.3).	85
Fig. 34	Muscovite granite showing perthitic texture and deformed quartz, muscovite is found in clusters between the light minerals. (C.N. Light 2.5×6.3).	85
Fig. 35	Diabase showing partially altered plagioclase and chloritized pyroxene with ocleritic texture. (C.N. Light 2.5×6.3).	88