



Ain Shams University
Faculty of Science
Botany Department

**PHYSIOLOGICAL STUDIES ON THE
PHENOMENON OF THE APICAL
DOMINANCE IN PLANT**

THESIS
Submitted for the Degree of
Doctor of Philosophy
IN
BOTANY
(**PHYSIOLOGY**)

مكتبة المعلومات الجامعية
الدراسات والبحوث
توثيق البكروفيلم

رسالة

By

Hemmat Khattab Ibrahim Khattab
(B. Sc. &M. Sc.)

583.322
H.K

97983

Supervisors

Prof. Dr.
Hassan Anwar Foda
Hassan Foda

Dr.
Amira Ahmed Hassanein
Amira Ahmed



1994

بِسْمِ اللَّهِ الرَّحْمَنِ الرَّحِيمِ

وَقُلْ رَبِّ زِدْنِي عِلْمًا

سورة الفاتحة



This Thesis has not been
previously submitted for a
degree at this or at any
other University

Hemmat Khattab Ibrahim Khattab

CONTENTS

| | Page |
|---|------|
| ACKNOWLEDGEMENT | |
| LIST OF TABLES | i |
| LIST OF FIGURES | ii |
| LIST OF PLATES | iii |
| GENERAL INTRODUCTION AND AIM OF WORK..... | 1 |
| MATERIALS..... | 7 |
| PART I | |
| Histological and Cytological Studies of Apical Dominance in Soybean Plants | |
| Introduction | 8 |
| Material and Methods | 13 |
| Results | 15 |
| Discussion | 39 |
| Summary | 44 |
| PART II | |
| Apical Dominance and Vascular Connections in Soybean Plants. | |
| Introduction | 45 |
| Material and Methods | 49 |
| Results and Discussion | 50 |
| Summary | 59 |

| | Page |
|--|-------------|
| PART III | |
| Carbohydrate Metabolism in Relation to Apical Dominance in Soybean Plants | |
| Introduction | 60 |
| Material and Methods | 66 |
| Results | 72 |
| Discussion | 88 |
| Summary | 95 |
| PART IV | |
| N, P, K and Mg Distribution in Relation to Apical Dominance in Soybean Plants | |
| Introduction | 97 |
| Material and Methods | 100 |
| Results | 102 |
| Discussion | 123 |
| Summary | 128 |
| GENERAL DISCUSSION AND CONCLUSIONS | 129 |
| SUMMARY | 138 |
| LITERATURES | 141 |
| ARABIC SUMMARY | |

LIST OF TABLES

Page

PART I

- Table 1:** Effect of decapitation of the main stem terminal bud and replacement of the decapitated bud with 10 gm Kg⁻¹ IAA in lanolin on bud length (mm) of the cotyledonary buds (V0) and first node axillary buds (V1) of soybean seedlings. 19
- Table 2:** Effect of decapitation of the main stem terminal bud and replacement of the decapitated bud with 10 gm Kg⁻¹ IAA in lanolin on the fresh weight of cotyledonary (V0) and first-node (V1) axillary buds of soybean seedlings. 22
- Table 3:** Effect of decapitation of the main stem terminal bud... and replacement of the decapitated bud with 10 gm Kg⁻¹ IAA in lanolin on the dry weight of cotyledonary (V0) and first node (V1) axillary buds of soybean seedlings. 24
- Table 4:** Effect of decapitation of the main stem terminal bud and replacement of the decapitated bud with 10 gm Kg⁻¹ IAA in lanolin on the mean number of mitotic figures per bud at either the cotyledonary (V0) or first (V1) node of soybean seedlings. 27

| | |
|---|----|
| Table 5: Effect of decapitation of the main stem terminal bud and replacement of the decapitated bud with 10 gm Kg ⁻¹ IAA in lanolin on the area and height of apical dome from median longitudinal sections of cotyledonary (V0) and first node (V1) axillary buds of soybean seedlings. | 30 |
| Table 6: Effect of decapitation of the main stem terminal bud and replacement of the decapitated bud with 10 gm Kg ⁻¹ IAA in lanolin on the number of cells and the number of mitotic cells per median longitudinal section of the apical dome of axillary buds at the cotyledonary (V0) and first (V1) node of soybean seedlings. | 32 |
| Table 7: Effect of decapitation of the main stem terminal bud and its replacement with 10 gm Kg ⁻¹ IAA in lanolin on bud length of V0 & V1 axillary buds of soybean seedlings. | 89 |
| Table 8: Effect of the removal of the main stem terminal bud and its replacement with 10 gm Kg ⁻¹ IAA in lanolin on the carbohydrate content of the cotyledonary node (V0) segments including the two cotyledonary buds. | 73 |

| | |
|--|----|
| Table 9: Effect of the removal of the main stem terminal bud and its replacement with 10 gm Kg ⁻¹ IAA in lanolin on the carbohydrate content of the first node (V1) segments including the two axillary buds. | 77 |
| Table 10: Effect of the removal of the main stem terminal bud and its replacement with 10 gm Kg ⁻¹ IAA in lanolin on the reducing sugars: starch ratio of V0 and V1 segments that included the axillary buds. | 81 |
| Table 11: Effect of the removal of the main stem terminal bud and its replacement with 10 gm Kg ⁻¹ IAA in lanolin on the sucrose phosphate synthase (SPS) activity of the V0 and V1 segments that included the axillary buds. | 82 |
| Table 12: Effect of the removal of the main stem terminal bud and its replacement with 10 gm Kg ⁻¹ IAA in lanolin on the sucrose synthase (SS) activity of the V0 and V1 segments that included the axillary buds. | 85 |
| Table 13: Effect of the removal of the main stem terminal bud and its replacement with 10 gm Kg ⁻¹ IAA in lanolin on the acid invertase activity of the V0 and V1 segments that included the axillary buds. | 87 |

Table 14: Effect of removing the main stem terminal bud, adding 10 g Kg⁻¹ IAA in lanolin to the tip of decapitated stem, applying 200 ppm BAP directly to inhibited axillary buds and applying 5 g Kg⁻¹ TIBA in lanolin as a ring around the stem directly beneath the terminal bud on the length and width of axillary buds at both the cotyledonary (V0) or first (V1) nodes of soybean seedlings. 103

Table 15: Effect of removing the main stem terminal bud, adding 10 g Kg⁻¹ IAA in lanolin to the tip of the decapitated stem, applying 200 ppm BAP directly to inhibited axillary buds and applying 5 g Kg⁻¹ TIBA in lanolin as a ring around the stem directly beneath the terminal bud on the dry weight of the root systems of soybean seedlings. 106

Table 16: Effect of removing the main stem terminal bud, adding 10 gm Kg⁻¹ IAA in lanolin to the tip of the decapitated stem, applying 200 ppm BAP directly to the inhibited axillary buds and applying 5 gm Kg⁻¹ TIBA in lanolin as a ring around the stem directly beneath the terminal bud on the total nitrogen content of axillary buds at both the V0 or V1 nodes of soybean plants. 108

Table 17: Effect of removing the main stem terminal bud, adding 10 gm Kg⁻¹ IAA in lanolin to the tip of the decapitated stem, applying 200 ppm BAP directly to the inhibited axillary buds and applying 5 gm Kg⁻¹ TIBA in lanolin as a ring around the stem directly beneath the terminal bud on the total nitrogen content of the cotyledonary (V0) or first node (VI) segments of soybean plant. 109

Table 18: Effect of applying 200 ppm BAP directly to the inhibited axillary buds and applying 5 gm Kg⁻¹ TIBA as a ring directly around the stem beneath the terminal bud on the nitrogen content of terminal buds of 14-day old soybean seedlings. 112

Table 19: Effect of removing the main stem terminal bud, adding 10 g Kg⁻¹ IAA in lanolin to the tip of the decapitated stem, applying 200 ppm BAP directly to the inhibited axillary buds and applying 5 gm Kg⁻¹ TIBA in lanolin as a ring around the stem directly beneath the terminal bud on the nitrogen content of the root system of soybean seedlings..... 114

- Table 20:** Effect of applying 200 ppm directly to the inhibited axillary buds and applying 5 g Kg⁻¹ TIBA in lanolin on the K, P and Mg content of the terminal buds of soybean seedlings. 117
- Table 21:** Effect of removing the main stem terminal bud, adding 10 gm Kg⁻¹ IAA in lanolin to the tip of the decapitated stem, applying 5 gm Kg⁻¹ TIBA in lanolin as a ring around the stem directly beneath the terminal bud, and applying 200 ppm BAP directly to the inhibited axillary bud on the K,P,and Mg content of the V0 node segments including cotyledonary buds of soybean seedlings. 118
- Table 22:** Effect of removing the main stem terminal bud, adding 10 gm Kg⁻¹ IAA in lanolin to the tip of the decapitated stem, applying 5 g Kg⁻¹ TIBA in lanolin as a ring around the stem directly beneath the terminal bud and applying 200 ppm BAP directly to the inhibited axillary buds on the K, P and Mg content of V1 node segments including axillary buds of soybean seedlings. 121

LIST OF FIGURES

Page

- Figure 1:** The effect of removing the: 1) young expanding leaves beneath the terminal bud 2) the terminal bud, 3) the terminal portion of the stem 6 cm above the first (V1) node, or 4) all of the terminal portion of the stem but 1 cm above the first node (V1) on the length of the axillary buds at the cotyledonary or first node of soybean seedlings. 17
- Figure 2:** The effect of decapitation of the main stem terminal bud or replacement of the decapitated bud with 10 gm Kg⁻¹ IAA in lanolin on the bud length of the axillary branches at the cotyledonary (V0) or first node (V1) of soybean seedlings. 20
- Figure 3:** The effect of decapitation of the main stem terminal bud or replacement of the decapitated bud with 10 gm Kg⁻¹ IAA in lanolin on the fresh weight of the axillary buds at the cotyledonary (V0) or first node (V1) of soybean plants. 23
- Figure 4:** The effect of decapitation of the main stem terminal bud or replacement of the decapitated bud with 10 gm Kg⁻¹ IAA in lanolin on the dry weight of the axillary branches at the cotyledonary (V0) or first node (V1) of soybean seedlings. 25

Table 5: The effect of decapitation of the main stem terminal bud or replacement of the decapitated bud with 10 gm Kg⁻¹ IAA in lanolin on the mean number of mitotic figures in the axillary buds at the cotyledonary (V0) or first node (V1) of soybean seedlings. 28

Figure 6: The effect of decapitation of the main stem terminal bud or replacement of the decapitated bud with 10 gm Kg⁻¹ IAA in lanolin on area of median longitudinal sections of the apical dome of axillary buds at the V0 or V1 node; height of median longitudinal sections of apical domes of axillary buds at the V0 or V1 node of soybean seedlings. 31

Figure 7: The effect of decapitation of the main stem terminal bud or replacement of the decapitated bud with 10 gm Kg⁻¹ IAA in lanolin on: the number of cells per median longitudinal section of the apical dome of axillary buds at the V0 or V1 node; the number of mitotic cells per median longitudinal section of the apical dome of axillary buds at the V0 or V1 of soybean seedlings. 33

- Figure 8:** The effect of decapitation of the main stem terminal bud or replacement of the decapitated bud with 1% IAA in lanolin on hexose, sucrose, starch and total sugar content of the cotyledonary segment (V0) and its buds of soybean seedlings. 74
- Figure 9:** The effect of decapitation of the main stem terminal bud or replacement of the decapitated bud with 1% IAA in lanolin on hexose, sucrose, starch and total sugar content of the first node segment (V1) and its buds of soybean seedlings. 78
- Figure 10:** The effect of decapitation of the main stem terminal bud or replacement of the decapitated bud with 1% IAA in lanolin on the SPS activity, SS activity and acid invertase of the V0 segments and its buds of soybean seedlings. 83
- Figure 11:** The effect of decapitation of the main stem terminal bud or replacement of the decapitated bud with 1% IAA in lanolin on the, SPS activity, SS activity and acid invertase of the first segment (V1) and its buds of soybean seedlings. 86