
Correlation between protein patterns in serum and tears of normal and some infectious eye diseases

Thesis
Submitted in partial fullfilment
for the master degree of science in Biochemistry



By

MARGERITTE AZMY AZIZ

Research Institute of Ophthalmology
Laboratory of Biochemistry

Under Supervision of

Prof. Dr. MOHAMED M. ABD EL-FATTAH

Head of Biochemistry Department
Faculty of Science
Ain Shams University

Prof. Dr.
FAWZI EL- SHOBAKI
Professor of Biochemical Nutrition
National Research Center

Prof. Dr. ZEINAB ZAKI EL-DARDERY

Professor of Biochemistry Religional Consecutive Faculty of Science
Ain Shams University

Ain Shams University
Faculty of Science
Biochemistry Department

(1993) Cairo





To My Professor

Dr. Fawzi A. El-Shobaki

ACKNOWLEDGEMENT

I express my thanks and deep gratitude to **Prof. Dr. Mohamed M. Abd El-Fattah,** Head of Biochemistry Department, Faculty of Science,

Ain Shams University for giving every possible help, and for his continuous support and advices during this work.

I owe special gratitude and my thanks to **Prof. Dr. Fawzi A. El-Shobki,** Prof. of Biochemistry and Nutritional Chemistry, National Research center, for suggesting the point of research, supervising the whole work and for his consistent supervision, valuable suggestion, building scientifically and great effect throughout the work.

My deep thanks and gratitude to **Prof. Dr. Zeinab Zaki EL-Dardery**, Prof. of Biochemistry, Faculty of Science, Ain Shams
University, for giving every possible help, and for her continuous support
and advices during this work.

I express my thanks and deep gratitude to **Prof. Dr. Magdoline Y. El-Gammal,** Prof. of Microbiology, Research Institute of ophthalmology for giving every possible help, and for her continuous support and advices during this work.



CONTENTS

	Page
PREFACE	1
CHAPTERT I:	
1. LITERATURE REVIEW.	
1.1. THE EYE.	3
1.2. OCULAR SURFACE DEFENSE MECHANISM.	4
1.3. SOURCE AND FUNCTION OF TEARS.	9
1.4. COMPOSITION, PHYSICAL AND CHEMICAL PROPERTIES	
OF TEARS.	14
1.5. BIOCHEMICAL CHANGES ASSOCIATING	
CONJUNCTIVITIS.	28
1.5.1. I. LACTO FERRIN.	28
1.5.2. II . IMMUNOGLOBULINS.	36
1.5.3. III . LYSOZYME.	44
1.5.4. IV. PREALBUMIN.	48
1.5.5. V. COMPLEMENT.	49
1.6. CLINICAL NOTES ON CONJUNCTIVAL DISEASES.	50
1.6.1. BACTERIAL CONJUNCTIVITIS.	52
1. 6. 2. SPRING CATARRH VERNAL CONJUNCTIVITIS.	56
1. 6. 3. ACUTE OR SUBACUTE ALLERGIC CATARRHAL	
CONJUNCTIVITIS.	57
1. 6. 4. CHLAMYDIAL CONJUNCTIVITIS (TRACHOMA).	58



CHAPTER II:

2.	MATERIALS AND METHODS.	66
	2. 1. PRODUCTION OF ANTIBODIES AGAINST LACTOFERRIN.	69
	2. 2. ESTIMATION OF ANITIBODIES TITER IN ANTISERUM.	 7 0
	2. 3. DETERMINATION OF ANTIBODY TITER BY ROCKET	
	IMMUNOELECTROPHORESIS.	 71
	2. 4. DEVELOPMENT OF SINGLE RADIAL IMMUNODIFFUSION	
	TECHNIQUE FOR ESTIMATION OF LACTOFERRIN.	- 72
	2. 5. DETERMINATION OF BLOOD HAEMOGLOBIN	
	CONCENTERATION.	79
	2. 6. DETERMINATION OF SERUM IRON.	7 9
	2. 7. DETERMINATION OF TOTAL IRON BINDING CAPACITY.	82
	2. 8. DETERMINATION OF SERUM TOTAL PROTEIN.	83
	2. 9. DETERMINATION OF TOTAL PROTEIN IN TEARS.	84
	2. 10. ELECTROPHORETIC SEPARATION OF PROTEINS FOR	
	BOTH SERUM AND TEARS.	87
	2. 11. COLLECTION OF SAMPLES FOR BACTERIAL	
	IDENTIFICATION	O1

CHAPTER III:

3. RESULT
3. 1. BLOOD HAEMOGLOBIN, SERUM IRON AND TOTAL IRO
BINDING CAPACITY.
3. 2. THE LEVEL OF SERUM TRANSFERRIN AND TEAR
LACTOFERRIN.
3. 3. THE LEVEL OF IgG AND IgA IN SERUM AND TEARS.
3. 4. SERUM TOTAL PROTEIN AND ELECTROPHORETICALLY
SEPARATED FRACTIONS.
3. 5. TEAR TOTAL PROTEIN AND ELECTROPHORETICALLY
SEPARATED FRACTIONS
CHAPTER . IV:
DISCUSSION
CHAPTER V:
SUMMARY.
REFRENCES
KEFKENUES
ADADIC CUMMADV
ARABIC SUMMARY.



Preface

Tear fluid contains a high concenteration of proteins. The proteins in tears may be distinguished into 3 main types namely albumin, γ -globulins and lysozyme. It is established that the antimicrobial activity of tears is in the γ -globulins and lysozyme fractions (Vaughan and Asbury, 1980).

Lactoferrin which was discovered in bovine milk has been detected in the neutrophils and in tears. This protein molecule is similar to transferrin present in blood and can bind reversibly to two atoms of iron (Selinger, et al, 1979). The capacity of lactoferrin or transferrin to bind iron makes it important as a factor that inhibit proliferation of several bacteria especially Gram negative bacteria due to removal of iron from the medium (Reiter, 1983).

In addition to this function, lactoferrin was also reported to protect tissues from oxidative damage by inhibition of lipid peroxidation (Gutteridge et al, 1981). Several other functions were reported to be performed by lactoferrin dealing with the immunity of the eye, it functions as a natural killer cells (Nishiya and Horwitz, 1982), it suppresses the in vitro primary antibody response (Duncan and Mc Arthur, 1981), exhibits anticomplementary effects (Kijlstra and Jeurissen, 1982) and stimulates the leucocytes (Kijlstra and Broersma, 1984).

It is thus clear that lactoferrin has both bacteriostatic and bacteriocidal properties and may interact with specific antibody to produce an antimicrobial system more powerful than either lactoferrin or specific antibody alone.

On the other hand blood transferrin which is similar to lactoferrin also performs several of these functions performed by lactoferrin. It is known that

