

**COMPUTERIZED TOMOGRAPHIC  
EVALUATION AND STAGING  
OF  
COLORECTAL CARCINOMA**



**Thesis  
Submitted in Partial Fulfilment  
for the Degree of M.D.  
Radiodiagnosis**



616.07572  
B.H

**BY  
Badr Mohamed Hegazy  
M.B.,B.CH. , M.S. Radiodiagnosis**

**Supervisor  
Prof. Dr. Ahmed Abd El Tawab Mohamed  
Prof. of Radiodiagnosis  
Ain Shams University**

**Faculty of Medicine  
Ain Shams University**

**1996**



Central Library - Ain Shams University



## List of Figures and Diagrams

|                                                                                     | <u>Page</u> |
|-------------------------------------------------------------------------------------|-------------|
| Fig. (1) : Surfaces projection of the colon.                                        | 3           |
| Fig. (2) : A radiograph of the abdomen<br>after administration of barium enema.     | 4           |
| Fig.(3) : The interior of the caecum                                                | 7           |
| Fig. (4) : The various positions of<br>the appendix                                 | 8           |
| Fig. (5 A, B) : Normal colon.                                                       |             |
| (A) A cross section (10mm) of air<br>filled rectum.                                 |             |
| (B) Air filled transverse colon                                                     | 11          |
| Fig. (6) : A coronal section of the pelvis and rectum                               | 16          |
| Fig. (7) : Coronal and transverse sections of<br>the rectum and adjacent structures | 17          |
| Fig. (8) : The radiographic anatomy of male pelvis                                  | 19          |
| Fig. (9) : Superior mesenteric artery and its branches                              | 22          |
| Fig. (10) : Inferior mesenteric artery and its branches                             | 23          |
| Fig. (11) : Blood supply of the rectum and anal canal                               | 24          |
| Fig. (12) : Lymphatic drainage of the colon                                         | 27          |
| Fig. (13) : Lymphatic drainage of the rectum                                        | 28          |
| Fig. (14) : Cross section of the colon shows the<br>longitudinal muscle fibres      | 31          |
| Fig. (15) : The macroscopic varieties of colorectal<br>carcinoma                    | 34          |
| Fig. (16) : Stage II carcinoma of the caecum                                        | 88          |
| Fig.(17 A, B):Stage II carcinoma of the sigmoid colon                               | 90          |
| Fig. (18 A, B, C) : Stage II carcinoma of the<br>sigmoid colon                      | 93          |

|                                                                                  | <b>Page</b> |
|----------------------------------------------------------------------------------|-------------|
| <b>Fig (19 A, B) : Stage II carcinoma of the caecum</b>                          | <b>96</b>   |
| <b>Fig. (20) : Stage II carcinoma of the caecum</b>                              | <b>98</b>   |
| <b>Fig. (21) : Stage II carcinoma of the rectum</b>                              | <b>100</b>  |
| <b>Fig. (22) : Stage II carcinoma of the ascending colon</b>                     | <b>102</b>  |
| <b>Fig. (23 A, B) : Stage II carcinoma of<br/>the descending colon</b>           | <b>105</b>  |
| <b>Fig.(24 A, B): Stage II carcinoma of the rectosigmoid</b>                     | <b>108</b>  |
| <b>Fig. (25 A, B) : Stage IIIA carcinoma of the rectum</b>                       | <b>110</b>  |
| <b>Fig.(26 A, B, C): Stage IIIA carcinoma of the caecum</b>                      | <b>113</b>  |
| <b>Fig. (27) : Stage IIIA carcinoma of the rectum</b>                            | <b>115</b>  |
| <b>Fig. (28) : Stage IIIA carcinoma of the rectum</b>                            | <b>117</b>  |
| <b>Fig. (29) : Stage IIIA carcinoma of the rectum</b>                            | <b>119</b>  |
| <b>Fig.(30 A, B) : Stage IIIB carcinoma of the ascending<br/>colon</b>           | <b>121</b>  |
| <b>Fig. (31) : Stage IIIB carcinoma of the rectum</b>                            | <b>123</b>  |
| <b>Fig. (32 A, B) : Stage IIIB carcinoma of the transverse<br/>colon</b>         | <b>125</b>  |
| <b>Fig.(33 A, B, C, D): Stage IV carcinoma of the rectum</b>                     | <b>128</b>  |
| <b>Fig.(34 A, B, C): Stage IV carcinoma of the rectum</b>                        | <b>131</b>  |
| <b>Fig. (35 A, B) : Stage IV carcinoma of the caecum</b>                         | <b>133</b>  |
| <b>Fig. (36 A, B) : Uterine sarcoma (false positive)<br/>case</b>                | <b>135</b>  |
| <b>Fig. (37) : Normal CT appearance of the pelvis<br/>(false negative)</b>       | <b>137</b>  |
| <b>Fig. (38 A, B) : Positive recurrence<br/>Stage II carcinoma of the caecum</b> | <b>139</b>  |
| <b>Fig. (39) : Positive recurrence<br/>Stage II carcinoma of the rectum</b>      | <b>141</b>  |

|                                                                                                                       | <u>Page</u> |
|-----------------------------------------------------------------------------------------------------------------------|-------------|
| <b>Fig. (40) : Positive recurrence</b>                                                                                |             |
| Stage IIIA carcinoma of the rectum                                                                                    | 143         |
| <b>Fig. (41) : Positive recurrence</b>                                                                                |             |
| Stage IIIA carcinoma of the rectum                                                                                    | 145         |
| <b>Fig. (42) : Positive recurrence</b>                                                                                |             |
| Stage IIIB carcinoma of the rectum                                                                                    | 147         |
| <b>Fig. (43) : Positive recurrence</b>                                                                                |             |
| Stage IIIB carcinoma of the<br>transverse colon                                                                       | 149         |
| <b>Fig. (44 A, B, C) : Positive recurrence</b>                                                                        |             |
| Stage IV carcinoma of the rectum                                                                                      | 152         |
| <b>Fig. (45 A, B) : Positive recurrence</b>                                                                           |             |
| Stage IV carcinoma of the rectum                                                                                      | 154         |
| <b>Fig. (46) : Postoperative granulation followed<br/>    abdomino-perineal resection of<br/>    rectal carcinoma</b> | 156         |



## **List of Tables**

|                                                                                                 | <b><u>Page</u></b> |
|-------------------------------------------------------------------------------------------------|--------------------|
| <b>Table (1) : Duke's classification</b>                                                        | <b>39</b>          |
| <b>Table (2) : Modified Duke's classification</b>                                               | <b>39</b>          |
| <b>Table (3) : TNM staging system</b>                                                           | <b>42</b>          |
| <b>Table (4) : Sex and age of the cases</b>                                                     | <b>48</b>          |
| <b>Table (5) : Shows different types of clinical presentation</b>                               | <b>49</b>          |
| <b>Table (6) : Shows the sites of the tumors</b>                                                | <b>50</b>          |
| <b>Table (7) : Shows methods of initial diagnosis of the<br/>preoperative group of patients</b> | <b>50</b>          |
| <b>Table (8) : Shows CT staging of colorectal carcinoma<br/>according to (Thoeni, 1989)</b>     | <b>57</b>          |
| <b>Table (9) : Shows : Group I<br/>preoperative cases</b>                                       | <b>59</b>          |
| <b>Table (10) : Shows : Group II<br/>Postoperative cases</b>                                    | <b>75</b>          |



## **List of Abbreviation**

|                |                                                 |
|----------------|-------------------------------------------------|
| <b>CT</b>      | <b>Computed tomography.</b>                     |
| <b>H. unit</b> | <b>Housfield unit.</b>                          |
| <b>US</b>      | <b>Ultrasonography.</b>                         |
| <b>UICC</b>    | <b>Union International Centre of<br/>Cancer</b> |
| <b>TNM</b>     | <b>TNM classification</b>                       |
|                | <b>T: Primary tumor</b>                         |
|                | <b>N: Regional lymph node.</b>                  |
|                | <b>M: Distant metastases.</b>                   |



**Hegazy, Badr Mohamed, computed tomographic evaluation and staging of colorectal carcinoma.**

**Radiology Department, Faculty of Medicine, Ain Shams University.**

### **Abstract**

**This study includes 60 patients with colorectal carcinoma either preoperative or postoperative. CT abdomen and pelvis was done for all patients.**

**CT is an effective method for demonstration of the primary lesion, direct extension of the tumor and distant metastases . CT is advised for all patient presented with colorectal carcinoma for evaluation and staging of the disease.**



# ***INTRODUCTION & AIM OF WORK***

