

Dc

**Evaluation Of Myocardial Perfusion And Wall Motion
Using Dobutamine Stress Technetium-99m Sestamibi
Scintigraphy In Patients With Coronary Artery
Disease**

*Thesis Submitted For The Partial Fulfillment
Of The M.D. Degree In Cardiology*

By

AMR ADEL EL-SAYED
M.B., B.Ch., M.S. (Cardiology)



E16. 12307572
A. A

Supervised By

53443

PROFESSOR HASSAN EZZ EL-DIN ATTIA

Head of the Department of Cardiology
Ain Shams University

PROFESSOR LAILA FARIS

Head of The Department of Radiotherapy and Nuclear Medicine
Ain Shams University

PROFESSOR SAID ABD EL-HAFIZ KHALED

Professor of Cardiology
Ain Shams University

DR. SAMEH SHAHEEN

Lecturer of Cardiology
Ain Shams University

AIN SHAMS UNIVERSITY

1997





بِسْمِ اللَّهِ الرَّحْمَنِ الرَّحِيمِ

وَقَدْ رَبَّزِدَنِي عِلْمًا

صَدَقَ اللَّهُ الْعَظِيمِ

CONTENTS

Introduction	1
Aim of the work	3
Review of the literature	4
Patients and methods	64
Results	71
Discussion	95
Summary	103
Conclusion and recommendations	106
References	107
Arabic Summary	

List of Tables

Table (1) :	Incidence of symptoms during dobutamine thallium-201 scintigraphy in 144 patients, according to Hays et al .	12
Table (2) :	Detection of coronary artery disease by dobutamine stress myocardial perfusion imaging .	15
Table (3) :	Comparison of Tc-99m sestamibi and Tl-201 imaging .	27
Table (4) :	Sensitivity and Specificity of Tc-99m Sestamibi SPECT Imaging .	33
Table (5) :	Correlation of coronary angiography and ST-segment depression in response to dobutamine stress as regards detection of ischemic heart disease .	74
Table (6) :	Correlation of coronary angiography and occurrence of chest pain in response to dobutamine infusion .	74
Table (7) :	Side effects of dobutamine infusion .	76
Table (8) :	Correlation between reversibility as assessed by single study stress Tc-99m sestamibi gated SPECT (stress perfusion / rest wall motion) and reversibility assessed by stress / rest perfusion .	79

- Table (9) :** Correlation between coronary angiography and dobutamine stress gated SPECT as regards wall motion abnormalities . 79
- Table (10) :** Correlation between the results of coronary angiography and dobutamine stress Tc-99m sestamibi gated SPECT as regards detection of single, two, three vessel and left main disease . 81
- Table (11):** Correlation of coronary angiography and dobutamine stress Tc-99m sestamibi gated SPECT as regards detection of left anterior descending artery affection . 85
- Table (12) :** Correlation of coronary angiography and dobutamine stress gated SPECT as regards detection of left circumflex artery affection . 85
- Table (13) :** Correlation of coronary angiography and Tc-99m sestamibi gated SPECT as regards detection of right coronary artery affection . 87
- Table (14) :** Correlation of coronary angiography and dobutamine stress Tc-99m sestamibi gated SPECT as regards detection of multivessel disease (two, three vessel or left main disease) . 87

List of Figures

<i>Fig. (1)</i>	Left ventricular perfusion segments according to vascular territories .	70
<i>Fig. (2)</i>	Hemodynamic response to dobutamine infusion (Heart rate) .	72
<i>Fig. (3)</i>	Hemodynamic response to dobutamine infusion (Blood pressure) .	72
<i>Fig. (4)</i>	Reasons for termination .	77
<i>Fig. (5)</i>	Dose of dobutamine .	77
<i>Fig. (6)</i>	Sensitivity and specificity of dobutamine Tc-99m sestamibi gated SPECT .	83
<i>Fig. (7)</i>	Detection of single and multivessel disease .	83
<i>Fig. (8)</i>	Gated SPECT images in short axis (SA), vertical long axis (VLA) and horizontal long axis (HLA), showing normal perfusion .	88
<i>Fig. (9)</i>	Twelve -lead ECG of a 48-year old female patient at a dobutamine dose of 40 $\mu\text{g}/\text{Kg}/\text{min}$. She developed mild chest pain with no ECG changes .	89
<i>Fig. (10)</i>	Gated SPECT images of the same patient showing a reversible anterior defect .	90
<i>Fig. (11)</i>	Left coronary angiography of the same patient in right anterior oblique view showing a 70% lesion in the middle 1/3 of the LAD .	91

- Fig. (12)** Twelve-lead ECG of a 55-year old male patient at a dobutamine dose of 30 $\mu\text{g}/\text{Kg}/\text{min}$. He developed chest pain and non-sustained ventricular tachycardia . 92
- Fig. (13)** Gated SPECT scan of the same patient showing reversible anterior and inferior defects and a fixed lateral defect . 93
- Fig. (14)** Left coronary angiography of the same patient in right anterior oblique view showing a tight lesion of the proximal circumflex artery and another lesion in the LAD . 94

