

EFFECT OF CHANGING THE pH OF BURN WOUND ENVIRONMENT ON THE CONTROL OF BURN WOUND SEPSIS

Thesis
Submitted for partial fulfillment of
M.D. in Plastic and Reconstructive Surgery

By
Salah Nasser Mohammed Afifi
(M.B, B. Ch & M.Sc)

Supervised by
Prof.Dr. Ikram Ibrahim Safe
Prof of Plast. and Reconst.Surgery
Faculty of Medicine, Ain Shams University

Dr. Ayman Abo El-Makarem Shaker
Lecturer in Plast, and Reconst. Surgery
Faculty of Medicine, Ain Shams University

Dr. Omnia Abo El-Makarem Shaker
Lecturer in Clinical Pathology
Faculty of Medicine, Ain Shams University

Faculty of Medicine
Ain shams university,
1998





ACKNOWLEDGMENT

*I would like to express my deepest gratitude and appreciation to **Prof.Dr. Ikram Ibrahim Safe**, Prof of Plast. and Reconst. Surgery Faculty of Medicine, Ain Shams University for the time spent guiding me to come out with this work. We both had a hard time navigating through this difficult subject, but by his guidance and support, it was pleasure to face these difficulties.*

*My deep thanks and appreciation go to **Dr. Ayman Abo El-Makarem Shaker**, Lecturer in Plast, and Reconst. Surgery, Faculty of Medicine, Ain Shams University, for his kind supervision, meticulous revision and remarks which on many occasions put me in the right direction.*

*I am also very grateful to **Dr.Omnia Abo El-Makarem Shaker**, Lecturer in Clinical Pathology, Faculty of Medicine, Ain Shams University, for her kind supervision and for the help and observations she offered during the conduction of this work.*

*Special thanks to **Professor Dr. Hassan Badran**, Professor of Plastic and Reconstructive Surgery, Faculty of Medicine, Ain Shams University, as it was his idea to conduct this thesis and he guided me at its first steps.*

*I owe special gratitude to **Professor Dr. Moustafa Hemida**, Professor and Head of Department of Plastic and Reconstructive Surgery, Faculty of Medicine, Ain Shams University for his sound advice, moral support and continuous encouragement*

*I can not deny the help and effort of **all my professors and colleagues** in Plastic Surgery Department and the Burn Unit for their help and valuable advice*

*I wish also to extend my deep thanks to **Mr. Khalid Kotb**, Technician in the microbiology lab, of the Burn Unit, Faculty of Medicine, Ain Shams University hospitals for his efforts in preparing the work and processing the specimens*

CONTENTS

INTRODUCTION	(1)
AIM OF THE WORK	(6)
REVIEW OF LITERATURE:	
Burn injury	(7)
Pathology of the Burn wound	(15)
Burn wound Infection	(23)
Microbiology of the burn wounds	(29)
Effects of B.W.S. on the burned patient ..	(46)
Diagnosis of B.W.S.	(51)
General measures for control of B.W.S....	(65)
Treatment of invasive burn wound infection.	(79)
Effect of pH on bacterial survival and metabolism	(114)
Buffer solutions	(135)
 MATERIALS AND METHODS	 (143)
RESULTS	(163)
DISCUSSION	(206)
SUMMARY AND CONCLUSION	(217)
REFERENCES	(222)
ARABIC SUMMARY.	

LIST OF TABLES

Table (1):	(115)
pH scale	
Table (2):	(117)
pH range of different microorganisms	
Table (3):	(138)
The physiologically important buffers	
Table (4):	(139)
Some commonly used buffers	
Table (5):	(144)
Characteristics of the 20 patients undergoing burn wound cultures for the in vitro study.	
Table (6):	(144)
Patients data.	
Table (7):	(147)
Characteristics of the 50 patients undergoing in vivo study.	
Table (8):	(156)
Group one patients data.	

Table (9):		(157)
Characteristics of the 10 patients in group one		
Table(10):		(157)
Group two, patients data.		
Table(11):		(158)
Characteristics of the 15 patients in group two.		
Table(12):		(158)
Group three. Patients data		
Table(13):		(159)
Characteristics of the 15 patients in group three.		
Table(14):		(159)
Group four. Patients data.		
Table(15):		(160)
Characteristics of the 2 patients in group four.		
Table(16):		(161)
Nitrate-phosphate buffer		
Table(17):		(164)
Microorganisms isolated from the leg wounds by swab cultures		

List of Tables (B)

Table(18):		(164)
Staph.aureus:		
A) On adjusted pH media:		
B- On Unadjusted media "pH 7.4": Wells technique:		
Table(19):		(166)
2) Strept. Pyogenes:		
A- On adjusted pH media		
B- On unadjusted media "pH 7.4: Wells technique		
Table(20):		(167)
3) Pseudomonas aeruginosa:		
A- On adjusted media:		
B- On unadjusted media "pH 7.4": Wells technique		
Table(21):		(168)
4) Klebsiella		
A- On adjusted media		
B- On Unadjusted media "pH 7.4": Wells technique		
Table(22):		(169)
5) E.Coli:		
A- On adjusted media		
B-On Unadjusted Media "pH 7.4": Wells technique		
Table(23):		(170)
6) Candida albicans		
A- On adjusted media		
B- On Unadjusted media "pH 7.4": Wells technique		
Table(24):		(172)
Microorganisms isolated from burn		
wounds by swab cultures in group one		
in the first 4 days.		

List of Tables (C)

Table(25):	(174)
Group one. Results (Biopsy findings both qualitative and quantitative, wound pH and blood pH).	
Table(26):	(179)
Mean log counts of pseud. aeruginosa "group one" 8 patients	
Table(27):	(179)
Mean values of wound pH in patients with Pseud. aerug. wound infection.	
Table(28):	(179)
Mean values of blood pH in the same group.	
Table(29):	(180)
Mean log counts of other organisms (Staph. aureus., E.coli and klebsiella) in group one (7 patients).	
Table(30):	(181)
Mean values of wound pH in the same patients	
Table(31):	(181)
Mean values of blood pH in the same group.	
Table(32):	(183)
Micro-organisms isolated from burn wounds by swab cultures in group two in the first 4 days.	
Table(33):	(184)
Group two. Results (Biopsy findings both qualitative and quantitative, wound pH and blood pH).	

List of Tables (D)

Table(34):	(188)
Mean log counts of Pseud. aerug. group two "7 patients"	
Table(35):	(189)
Mean values of wound pH in patients with Pseud.aerug. wound infection	
Table(36):	(189)
Mean values of blood pH in the same group	
Table(37):	(190)
Mean log counts of other organisms (staph.aureus, E. coli and klebsiella) in group two. 8 patients	
Table(38):	(190)
Mean values of wound pH in the same patients.	
Table(39):	(191)
Mean values of blood pH in the same group	
Table(40):	(192)
Microorganisms isolated from burn wounds by swab cultures in group III patients in the first 4 days	
Table(41):	(194)
Group three. Results (biopsy findings both quantitative and quantitative, wound pH and blood pH)	

List of Tables (E)

.....
Table(42):(198)
Mean log counts of Pseud. aerug. in
group three "% patients"

Table(43):(199)
Mean values of wound pH in patients
with Pseud. aerug. wound infection in
group three.

Table(44):(199)
Mean values of blood pH in the same group
Mean log counts of other organisms
(Staph.aureus, E. coli, and
klebsiella) in group three "% patients".

Table(45):(200)
Mean log counts of other organisms

Table(46):(201)
Mean values of wound pH in the same
group

Table(47):(201)
Mean values of blood pH in the same
group

Table(48):(202)
The findings in the five patients of
group four (wound colony, wound pH,
blood pH)

.....
List of Tables (F)

Table(49):	(204)
Mean log counts of Pseud. aerug. in patients of group four (4 patients).	
Table(50):	(204)
Mean values of wound pH for the same patients	
Table(51):	(205)
Mean values of blood pH	

List of Tables (G)

Fig.(10):		(
	Weighing the specimens	
Fig.(11):		(
	Biopsy specimen after maceration	
Fig.(12):		
	Suspension and serial dilutions	
Fig.(13):		
	Measurement of wound pH by the narrow range indicator paper	
Fig.(14):		
	pH meter	
Fig.(15):		
	Growth of Staph aureus on adjusted media (acidic side)	
Fig.(16):		
	Growth of Staph aureus on unadjusted media (wells technique)	
Fig.(17):		
	Growth of Strept pyogenes on adjusted media (acidic side)	
Fig (18):		
	Growth of Strept pyogenes on unadjusted media (wells technique)	

List of figures (B)

Fig. (19):	(167)
Growth of <i>Pseudomonas aerug.</i> on adjusted media (acidic side)	
Fig. (20):	(167)
Growth of <i>Pseudomonas aerug.</i> on unadjusted media (wells technique)	
Fig. (21):	(168)
Growth of <i>Klebsiella</i> on adjusted media (acidic side)	
Fig. (22):	(168)
Growth of <i>Klebsiella</i> on unadjusted media (wells technique)	
Fig. (23):	(169)
Growth of <i>E.coli</i> on adjusted media (acidic side)	
Fig. (24):	(169)
Growth of <i>E.coli</i> on unadjusted media (wells technique)	
Fig. (25):	(178)
A) Deep dermal and full-thickness burn of both lower limbs at day 5 postburn	
Left lower limb dressed by the buffer solution (soaks/12 hours).	
Right lower limb dressed by the conventional technique.	
B) 12 days post-treatment.	

List of figures (C)