


بِسْمِ اللَّهِ الرَّحْمَنِ الرَّحِيمِ



**MAGNETIC RESONANCE IMAGING IN
INTRACEREBRAL HEMATOMAS**



THESIS

*Submitted For The Partial Fulfillment Of The
Master Degree In Radiodiagnosis
Faculty Of Medicine
Ain Shams University*

لجنة
الدراسة

By

د. خالد
تالاات
خايري

616.07548
H.A

HALA ABD EL MOHSEN SOLIMAN

(M.B.B.Ch)

52959

SUPERVISOR

Prof. Dr. KHALED TALAAT KHAIRY

Assistant Professor of Radiodiagnosis
Faculty Of Medicine - Ain Shams University



1996

CONTENTS

Acknowledgments	7
Introduction and aim of the work	11
Physical principles of magnetic resonance imaging.....	15
- Preparatory Alignment	16
- RF Excitation.....	20
- Relaxation and signal measurement.....	21
- Pulse Sequences and Image Contrast	25
- Rapid partial saturation	30
- Echo Planar Methods.....	31
- Spatial Localization.....	32
- Simultaneous Slice Acquisition.....	34
- Volume Acquisition	35
- Slice selection.....	35
- General principles of instrumentation.....	37
- Contrast agents	41
- Image artefacts	43
Anatomical aspects of MRI of the brain.....	49
- Anatomical aspects of the brain stem.....	50
- Anatomical aspects of the cranio-cervical junction	52
- Anatomical aspects of the white matter	53
- Anatomical aspects of the pituitary gland	55
Pathology of intracerebral hematomas	75
- Spontaneous intracerebral hematomas	75
- Primary intracerebral hematomas.....	83
- Spontaneous supratentorial hemorrhage	85
- Spontaneous infratentorial hemorrhage	92
- Secondary intracerebral hematomas.....	100
- Traumatic intracerebral hematomas.....	107
MRI manifestations of intracerebral hematomas.....	111
- Biochemical basis of the MR appearance of cerebral hemorrhage... 111	
- Clinical aspects of MR imaging of hemorrhage.....	121
Summary and Conclusion.....	145
References.....	153
Arabic Summary.....	162

INDEX OF FIGURES

Figure 2-1	A spinning nucleus having charge has a magnetic field around it in the form of a magnetic dipole moment.	17
Figure 2-2	If tissue is placed within a strong magnetic field each individual nucleus aligns itself either parallel (A) or antiparallel (B).	17
Figure 2-3	If the residual parallel aligned moments are decomposed into components along the transverse plane versus components in the longitudinal plane, it is found that all components subtract in the transverse plane while all the components add up in the longitudinal plane.	20
Figure 2-4	Illustration of 90° and 180° tipping of the bulk magnetization vector	21
Figure 2-5	Different views of the RF excitation process.	22
Figure 2-6	Longitudinal section through an MR magnet with the patient positioned in the body coil for scanning.	39
Figure 2-7	Three sets of gradient coils which after the main magnetic field shown in the three orthogonal planes	40
Figure 3-1	A-L Brain, Axial MR images.	56
Figure 3-2	A-D Brain. Sagittal MR images.	62
Figure 3-3	Midsagittal MR image.	64
Figure 3-4	Brain. Coronal MR images.	66
Figure 3-5	Pituitary fossa (sella tursica). Coronal MR images	70
Figure 4-1	Acute right putaminal hemorrhage in a patient with history of hypertension.	86
Figure 4-2	Acute hemorrhage in the head of the caudate nucleus, ant. limb of the internal capsule.	88
Figure 4-3	Thalamic hemorrhage from an occult vascular malformation	89
Figure 4-4	Left parietal lobar hemorrhage	90
Figure 4-5	Hemorrhage into chromophobe pituitary adenoma.	92
Figure 4-6	Cerebellar hemorrhage in the subacute stage..	94
Figure 4-7	Pontine hemorrhage.	97
Figure 4-8	Midbrain hemorrhage of unknown etiology.	99
Figure 4-9	Subacute hematoma in the right superior frontal gyrus and a second one in the left frontal operculum.	101
Figure 4-10	70 year old male with lung carcinoma with hemorrhage into a temporal lobe metastasis.	102
Figure 4-11	Right parietooccipital subcortical hemorrhage due to anticoagulants	104
Figure 4-12	Right occipital hemorrhagic infarction.	105
Figure 4-13	Occult vascular malformation in a 22 year old woman.	106
Figure 5-1	Proton relaxation in water..	112

Figure 5-2	A paramagnetic ion has been added to the aqueous solution. The large magnetic moment associated with the unpaired outer shell electrons of the paramagnetic ion results in enhanced T1 relaxation with respect to pure water	113
Figure 5-3	A macromolecule-metal complex has been added to the aqueous solution. This further slows the tumbling motion of the water molecule.	115
Figure 5-4	Illustration of the effects of intracellular compounds differing magnetic susceptibility on the magnetic field surrounding the cell.	116
Figure 5-5	Magnetic susceptibility of hemoglobin breakdown products	122
Figure 5-6	When water molecules diffuse through normal brain matter, they experience a homogenous magnetic field.	124
Figure 5-7	Graph showing evolution of signal intensity of cerebral hematoma on MRI.	126
Figure 5-8	Temporal changes of hemorrhage on MR images.	128
Figure 5-9	Illustration of signal loss due to intravoxel and intervoxel variations in magnetic susceptibility.	134
Figure 5-10	Axial FLASH MR image at 1.5 T.	135
Figure 5-11	Parasagittal T ₁ -weighted spin-echo MR image demonstrating a subarachnoid blood clot	136
Figure 5-12	Densely calcified fossa meningioma.	140
Figure 5-13	The axial T ₁ -weighted image demonstrates bright signal surrounding an area of signal loss bilaterally in the basal ganglia	141

INDEX OF TABLES

Table 2-1	Nuclei of interest in biological MRI	16
Table 4-1	Causes of intracerebral hemorrhage	76
Table 4-2	Distribution by site	77
Table 5-1	Summary of physical and magnetic properties of hemoglobin breakdown products	127
Table 5-2	Time course of hemorrhage signal intensity	127

.....

ACKNOWLEDGMENTS

ACKNOWLEDGMENTS

I would like to express my gratitude to Professor KHALED TALAAT KHAIRY for his helpful comments, suggestions and unlimited support. He has given individual assistance in numerous points of detail and continued to guide this work.

Last but not the least, my sincere thanks and gratitude are due to all members of my FAMILY who endured me during carrying out this work.

INTRODUCTION AND AIM OF THE WORK

