

Chemical Peeling in Egyptian Patients

Thesis
Submitted for partial fulfillment of
Master Degree (M.Sc.)
In Dermatology and Venereology

6165
y.A.

By
Yasser Abd El-Sattar Abo Mandour
"M.B.B.Ch."

65429

Supervised by
Prof. Dr. Mohamad Farid Abd El-Latif
Professor of Dermatology & Venereology
Faculty of Medicine, Ain Shams University

Prof. Dr. Mahira Hamdy El-Sayed
Assistant Professor of Dermatology & Venereology
Faculty of Medicine, Ain Shams University

Faculty of Medicine
Ain Shams University
1998

Acknowledgment

First and foremost, thanks is due to **ALLAH** the most merciful, the most compassionate.

I wish to express my sincere gratitude to my Prof. Dr. Mohamad Farid Abd El-Latif, Professor of Dermatology & Venereology, Faculty of Medicine, Ain Shams University, for his valuable help, whole hearted support and guidance that made this work possible.

I would like to express my deepest thanks and profound gratitude to Dr .Mahira Hamdy El-Sayed, Assesstant Professor of Dermatology & Venereology, Ain Shams University, for her continuous supervision, encouragement and guidance.

Finally, I would like to thank everyone who helped me throughout this work.



LIST OF CONTENTS

INTRODUCTION AND AIM OF THE WORK.....	1
REVIEW OF LITERATURE	3
CHAPTER I: History of Chemical Peeling	3
CHAPTER II: Clinical Strategies for Patient Selection....	6
CHAPTER III: Superficial Peeling	22
CHAPTER IV: Medium-Depth peeling	109
CHAPTER V: Deep Peeling	137
CHAPTER VI: Complications of Chemical Peeling.....	174
SUBJECTS AND METHODS.....	205
RESULTS.....	214
DESCUSSION.....	261
CONCLUSION.....	268
SUMMARY.....	269
REFERENCES.....	277
ARABIC SUMMARY.....	308

LIST OF TABLES

	Page
Table (1): History of peels.....	5
Table (2): Factors in patient evaluation for subsequent relative contraindications.....	6
Table (3): Fitzpatrick's classification of sun reactive skin types.....	7
Table (4): Features of actinically damaged skin.....	10
Table (5): Photoaging groups-Glogau's classification.....	12
Table (6): Some A.H.A.s, their chemical structure and the fruits they are present in.....	49
Table (7): List of A.H.A.s in relation to their antioxidant Properties.....	50
Table (8): pKa of AHA & non-AHA compounds at 25 C.....	51
Table (9): pH of Some A.H.As at different concentrations.....	53
Table (10) Glycolic Acid Peel Exposure Time.....	57
Table (11): Jessner's solution or Combes formula.....	69
Table (12): Letessier's modification of Unna's paste.....	71
Table (13): Salicylic Acid Ointment.....	80
Table (14):Stratum corneum and epidermal thickness following 36%T.C.A.and 70% glycolic acid peels.....	104
Table (15):Histological changes after combined-depth peeling with JS + TCA 35% and CO2 + TCA 35%.....	133-134
Table (16): Suggested sequence for full - face Baker's peel.....	153

Table (17): Complication of chemical peeling.....	175
Table (18): Possible complications of chemical peeling.....	176
Table (19): Pigmentary problems by percent.....	180
Table (20): Location of scarred areas by percent.....	184
Table (21): Cardiac complication by percent.....	198
Table (22): Relative contraindications.....	203
Table (23): Demographic and patient's characteristics.....	236
Table (24): The base data of patients under study.....	237
Table (25): Results of glycolic acid 50% peel after 6 weeks.....	238
Table (26): Results of glycolic acid 50% peel after 12 weeks.....	239
Table (27): Complications of chemical peeling.....	240
Table (28): Number and percentage of patients with different grades of improvement in comedones according to the total number of patients with comedones.....	241
Table (29): Number and percentage of patients with different grades of improvement in papules according to the total number of patients with papules.....	241
Table (30): Number and percentage of patients with different grades of improvement in pustules according to the total number of patients with pustules.....	241
Table (31): Number and percentage of patients with different grades of improvement in nodules according to the total number of patients with nodules.....	242

Table (32): Number and percentage of patients with different grades of improvement in cysts according to the total number of patients with cysts.....	242
Table (33): Number and percentage of patients with different grades of improvement in post inflammatory hyperpigmentation according to the total number of patients with post inflammatory hyperpigment- ation assessed by their number.....	242
Table (34): Number and percentage of patients with different grades of improvement in post inflammatory hyperpigmentation according to the total number of patients with post inflammatory hyperpigment- ation assessed by H.A.S.I. Score.....	243
Table (35): Number and percentage of patients with different grades of improvement in superficial acne scars according to the total number of patients with superficial acne scars.....	243
Table (36): Number and percentage of patients with different grades of improvement in deep acne scars according to the total number of patients with deep acne scars.....	243
Table (37): Improvement in comedones after 2 weeks from the start of the study.....	244
Table (38): Improvement in comedones after 6 weeks from the start of the study.....	244
Table (39): Improvement in comedones at the end of the study.....	245

Table (40): Improvement in papules after 2 weeks from the start of the study.....	245
Table (41): Improvement in papules after 6 weeks from the start of the study.....	246
Table (42): Improvement in papules at the end of the study.....	246
Table (43): Improvement in pustules after 2 weeks from the start of the study.....	247
Table (44): Improvement in pustules after 6 weeks from the start of the study.....	247
Table (45): Improvement in pustules at the end of the study.....	247
Table (46): Improvement in nodules through the study.....	248
Table (47): Improvement in cysts through the study.....	248
Table (48): Improvement in post inflammatory hyperpigmentation according to number after 2 weeks from start.....	249
Table (49): Improvement in post inflammatory- hyperpigmentation according to number after 6 weeks from start.....	249
Table (50): Improvement in post inflammatory- hyperpigmentation according to number at the end of the study.....	249
Table (51): Improvement in post inflammatory- hyperpigmentation according to H.A.S.I. after 2 weeks.....	250
Table (52): Improvement in post inflammatory- hyperpigmentation according to H.A.S.I. after 6 weeks.....	250
Table (53): Improvement in post inflammatory- hyperpigmentation according to H.A.S.I. at the end of the study.....	251

Table (54): Improvement in superficial acne scars after 2 weeks from the start of the study.....	251
Table (55): Improvement in superficial acne scars after 6 weeks from the start of the study.....	252
Table (56): Improvement in superficial acne scars at the end of the study.....	252
Table (57): Improvement in deep acne scars after 2 weeks from the start of the study.....	252
Table (58): Improvement in deep acne scars after 6 weeks from the start of the study.....	253
Table (59): Improvement in deep acne scars at the end of the study.....	253

LIST OF FIGURES

	Page
Fig.(1): Diagram of the skin showing the wounding depth of superficial, medium - depth, and deep peels.....	2
Fig (2): Nasolabial furrows, Perioral wrinkles and Lateral forehead wrinkles	11
Fig (3): Glogau's classification for photoaging: (A) Type I (B) Type II (C) Type III (D) Type IV	13
Fig.(4):A - Sixteen year old. Asian female with a 3 year history of recalcitrant papulo-pustular acne. B- Appearance after a series of six applications of 20-35% glycolic acid peels, septra D.S. and 15% glycolic acid gel.....	25
Fig.(5):A-Papulopustular acne with post-inflammatory hyperpigmentation in an African-American woman. B - Appearance after a series of six applications of 50 %glycolic acid peels while on oral erythromycin.....	27
Fig.(6):A-A twenty year old Caucasian female with inflammatory papulo -pustular acne with post- acne scars B- Appearance following five months of treatment including five glycolic acid peels 20 - 70 % and four months of oral erythromycin 250 mg three times daily	29
Fig. (7) : A - Melasma in a Hispanic woman B- Appearance 4 months later following a series of three 50 % glycolic acid peels and topical at home use of 2 % hydroquinone and 10% glycolic acid gel.....	33

Fig. (8): A - A 22 years old female with melasma B-Appearance 3 months later after a series of four 50% glycolic acid peels.....	35
Fig.(9):A - Acne with post inflammatory hyper- pigmentation in an African - American woman B- Three months later following at home topical application of 2 % hydroquinone with 10% glycolic acid gel	36
Fig. (10): A- Actinic keratoses in photo damaged skin of a Caucasian woman B-One year later following single 50%glycolic acid peel and at home twice daily use of 2% hydroquinone and 10% glycolic acid gel	37
Fig.(11):A- Lamellar Ichthyosis in a 14 years old girl B- Appearance after application of 4 times daily of 5% glycolic acid in hydrophilic ointment for one week and once daily for the next two weeks	41
Fig. (12):A- Lower leg of 63 years old woman with winter time dry skin B- Appearance after lactic acid 12% partially neutralized in oil in water cream twice daily for one week	43
Fig. (13): Tray set-up for glycolic acid peel.From left to right: cleansing lotion, glycolic acid, alcohol and acetone.....	58
Fig. (14). Application of glycolic acid to the face: A-Skin is cleansed first B- Glycolic acid is applied with a large-tip cotton applicator C- A fan is used to alleviate any stinging or itching sensation	59

Fig. (15): Application of glycolic acid to the body	60
Fig. (16): The parent structure of trichloroacetic acid, with chloride atoms substituting for the hydrogen atoms on the alpha carbon.....	65
Fig. (17): Chemical Peel Consent.....	68
Fig (18): The whitish film produced by repeated T.C.A. application	78
Fig.(19): A- Photoaging skin and fine wrinkles, sagging skin B- After topical tretinoin for 4 months there is a limited improvement in texture and fine wrinkles but no change in sagging skin.....	78
Fig. (20): Normal skin. (A,Hematoxylin and eosin;B. Trichrome; C, Verhoeff- van Gieson)	85
Fig. (21) 35% TCA-treated skin at 2 days (A,B, and C) (A hematoxylin and eosin; B, trichrome; C, Verhoeff-van Gieson).....	88
Fig. (22): 70% glycolic acid-treated skin at 2 days (A,B, and C) (A hematoxylin and eosin;B, trichrome; C , Verhoeff-van Gieson)	91
Fig. (23): 35% TCA-treated skin at 2 weeks (A and B) (A, Hematoxylin and eosin; B , Verhoeff-van Gieson).....	94
Fig. (24): 70% glycolic acid-treated skin at 2 weeks (A and B). (A, Hematoxylin and eosin; B , Verhoeff-van Gieson).....	96
Fig. (25): 35% TCA - treated skin at 2 months (A, B and C) (A, Hematoxylin and eosin; B, Trichrome, C, Verhoeff-van Gieson.).....	98
Fig. (26): 70% glycolic acid-treated skin at 2months (A, B, and C) (A, Hematoxylin and eosin; B, Trichrome, C, Verhoeff-van Gieson.)	101

Fig (27): Tretinoin treated forearm skin: A-Before therapy B-After 4 months of therapy	108
Fig. (28): Medium depth peeling for pigmentary problems and photoaging. A-Pre-operative appearance. B- Post-operative appearance after a month.....	111
Fig.(29):A-Chemical peeling for facial skin. Preoperative appearance of keratoses, lentigins and rhytides. B-Materials include Jessner's solution, 35% T.C.A., saline, cotton tips and gauze. C-Application of Jessner's solution with a cotton tip applicator. D-Full application of 35%T.C.A. with frosting. E-Four days postoperative postoperative. F-Six months.....	118
Fig. (30):A-Severe Photoaging skin with deep wrinkles and sagging of skin. B-50% T.C.A. full face peel applied. C-Appearance four days post peel with crusting. D-Before T.C.A. peel E-After T.C.A. peel.....	113
Fig. (31):Colloidal iron stain before (A)and after (B) deep peeling. A thick band of new collagen can be noticed	172
Fig. (32): Patient record for glycolic acid peel,	207
Fig. (33): Glycolic acid chemical peel consent.....	208
Fig. (34): Pre- and post-glycolic acid peel patient instructions.....	209
Fig (35):Materials used: Ethyl alcohol, acetone, 50 % glycolic acid, neutralising solution. sunscreen, antibiotic cream. shower cap, fitting goggles and brush.....	212

Fig. (36):Glycolic acid 50 % peeling procedure: A-Patient after preparation and degreasing. B-The glycolic acid is neutralized after proper exposure time. C-Thorough washing. D-Cold fomentation to alleviate any residual pain.....	213
Fig.(37):Comedones improvement(mean).....	217
Fig.(38): Papules improvement (mean).....	218
Fig.(39): Pustules improvement (mean).....	219
Fig.(40): Nodules improvement (mean).....	220
Fig.(41): Cysts improvement (mean).....	221
Fig.(42): P.I.H. number improvement (mean).....	222
Fig.(43): P.I.H. improvement assessed by H.A.S.I. (mean).....	223
Fig.(44): Superficial acne scars improvement (mean).....	224
Fig.(45): Deep acne scars improvement (mean).....	225
Fig.(46): Male to female ratio in the study group.....	226
Fig.(47): Percentage of patients with different results as regards (comedones).....	227
Fig.(48): Percentage of patients with different results as regards (papules).....	228
Fig.(49): Percentage of patients with different results as regards (pustules).....	229
Fig.(50): Percentage of patients with different results as regards (nodules).....	230
Fig.(51): Percentage of patients with different results as regards (cysts).....	231
Fig.(52): Percentage of patients with different results as regards(P.I.H.).....	232
Fig.(53): Percentage of patients with different results as regards(superficial acne scars).....	233