

STRUCTURAL AND FUNCTIONAL RESPONSE OF
FRESHWATER FISHES TO ENVIRONMENTAL
POLLUTION BY LEAD

THESIS

Submitted for the Degree of
DOCTOR OF PHILOSOPHY

in
ZOOLOGY

31307

597.0929
A. M.




BY

AMINA MOHAMED FARAG-ALLAH
(M.Sc.)

Assistant Lecturer in Zoology Department
University College for Women
Ain Shams University

January
1990




”لَا يُكَلِّفُ اللَّهُ نَفْسًا إِلَّا وُسْعَهَا لَهَا
مَا كَسَبَتْ وَعَلَيْهَا مَا اكْتَسَبَتْ رَبَّنَا
لَا تُؤَاخِذْنَا إِنْ نَسِينَا أَوْ أَخْطَأْنَا“


صَدَقَ اللهُ الْعَظِيمُ
(سورة البقرة ٢٨٦)




STRUCTURAL AND FUNCTIONAL
RESPONSE OF FRESHWATER
FISHES TO ENVIRONMENTAL
POLLUTION BY LEAD

BOARD OF SCIENTIFIC
SUPERVISION

Professor Fatma M. Mazhar, (Ph.D.), 
Professor of Comparative Anatomy,
Embryology and Marine Biology and
Ex-Dean of the University College for
Women, Ain Shams University.

Professor Madiha A.M. Ashry, (Ph.D.), 
Professor of Histochemistry and
Head of Department of Zoology,
University College for Women,
Ain Shams University.



ACKNOWLEDGEMENT

The author of this thesis would like to express her deep thanks, gratitude and sincere appreciation to Prof. Dr. Fatma M. Mazhar, ex-Dean and Professor of Comparative Anatomy, Embryology and Marine Biology, Ain Shams University College for Women for her instructive supervision, constructive criticism, and critical reading of the manuscript. To her I owe a great deal for her continuous encouragement and valuable advice. No words can be sufficient to express my gratitude and indebtedness.

I wish also to express my deep thanks and gratitude to Prof. Dr. Madiha A. Ashry, Head of Zoology Department and Professor of Histochemistry, Ain Shams University College for Women for her instructive supervision. To her I feel indebted for her valuable advice, immeasurable time and effort, continuous encouragement, and for critical reading of the manuscript.

Also, I am grateful to Prof. Dr. Monir A. El-Ganzuri, Professor of Cell Biology, Faculty of Science, Ain Shams University, for helping me to estimate the LC_{50} . Thanks are also due to him for his help throughout the whole work.

CONTENTS

LIST OF TABLES.....	
LIST OF HISTOGRAMS.....	
LIST OF FIGURES.....	
LIST OF TEXT-FIGURES.....	
CHAPTER I : INTRODUCTION.....	1
AIM OF THE WORK.....	11
CHAPTER II : REVIEW OF LITERATURE.....	13
- HAEMATOLOGY.....	13
- HISTOLOGY.....	18
CHAPTER III : MATERIAL AND METHODS.....	23
I- ESTIMATION OF LC ₅₀	24
II- HAEMATOLOGICAL STUDIES.....	25
- Blood Collection.....	25
i - Erythrocyte Count.....	25
ii - Haemoglobin Content.....	27
iii - Haematocrit Percentage.....	28
iv - Total Leucocyte Count.....	28
v - Coagulation Time.....	29
- Statistical Analyses.....	30
III- HISTOLOGICAL TECHNIQUES.....	31

CHAPTER IV	RESULTS.....	34
	I- SIGNS OF POISONING AND ESTIMATION	
	OF LC ₅₀	33
	- Signs of Poisoning	33
	- Estimation of LC ₅₀	34
	II- HAEMATOLOGICAL STUDIES.....	37
	i - Erythrocyte Count.....	37
	ii - Haemoglobin Content.....	41
	iii- Haematocrit Percentage.....	45
	iv - Total Leucocyte Count.....	48
	v - Coagulation Time.....	51
	III- HISTOLOGICAL STUDIES.....	55
	1 - Liver.....	55
	2 - Ileum.....	69
	3 - Kidneys.....	80
	4 - Testes.....	88
	5 - Ovaries.....	99
	6 - Body wall (Integument and	
	Hypaxial Skeletal Muscles)...	109
CHAPTER V	: DISCUSSION.....	120
CHAPTER VI	: SUMMARY.....	173
CHAPTER VII	: BIBLIOGRAPHY.....	183
CHAPTER VIII	: ARABIC SUMMARY.....	

LIST OF TABLES

- Table (1) : Principal uses of lead.
- Table (2) : The concentrations of lead nitrate given and the percentage of mortality with each concentration.
- Table (3) : Abridged tabulation of the results of table (2).
- Table (4) : Average of erythrocyte counts \pm S.D. in control and experimental groups exposed to graded lead nitrate concentrations.
- Table (5) : Average of haemoglobin content (grams)% \pm S.D. of control and experimental groups exposed to different lead nitrate concentrations.
- Table (6) : Average of haematocrit (P.C.V.) \pm S.D. in control and experimental groups exposed to varying lead nitrate concentrations.
- Table (7) : Average of leucocyte counts \pm S.D. in control and experimental groups exposed to graded lead nitrate concentrations.
- Table (8) : Average of coagulation time (seconds) \pm S.D. in control and experimental groups exposed to graded lead nitrate concentrations.

LIST OF HISTOGRAMS

- Histogram (1) : Erythrocyte counts of control and experimental groups of Clarias lazera (Bars indicate standard deviations).
- Histogram (2) : Haemoglobin content of control and experimental groups of Clarias lazera (Bars indicate standard deviations).
- Histogram (3) : Haematocrit percentage of control and experimental groups of Clarias lazera (Bars indicate standard deviations).
- Histogram (4) : Leucocyte counts of control and experimental groups of Clarias lazera (Bars indicate standard deviations).
- Histogram (5) : Coagulation time of control and experimental groups of Clarias lazera (Bars indicate standard deviations).

LIST OF FIGURES

- Fig.(1) : Liver section of control fish.(X - 660).
Fig.(2) : Liver section of control fish.(X - 1650).
Fig.(3) : Liver section of control fish.(X - 660).
Fig.(4) : Liver section, 2 days after exposure to $1/8$ LC_{50} of lead nitrate. (X - 660)
Fig.(5) : Liver section, 2 days after exposure to $1/8$ LC_{50} of lead nitrate. (X - 660).
Fig.(6) : Liver section, 4 days after exposure to $1/8$ LC_{50} of lead nitrate. (X - 1050).
Fig.(7) : Liver section, 6 days after exposure to $1/8$ LC_{50} of lead nitrate. (X - 660).
Fig.(8) : Liver section, 10 days after exposure to $1/8$ LC_{50} of lead nitrate. (X - 460).
Fig.(9) : Liver section, 2 days after exposure to $1/4$ LC_{50} of lead nitrate. (X- 660).
Fig.(10) : Liver section, 2days after exposure to $1/4$ LC_{50} of lead nitrate. (X - 460).
Fig.(11) : Liver section, 4 days after exposure to $1/4$ LC_{50} of lead nitrate. (X - 660).
Fig.(12) : Liver section, 4 days after exposure to $1/4$ LC_{50} of lead nitrate. (X - 1650).
Fig.(13) : Liver section, 4 days after exposure to $1/4$ LC_{50} of lead nitrate. (X - 165).
Fig.(14) : Liver section, 6 days after exposure to $1/4$ LC_{50} of lead nitrate. (X - 460).
Fig.(15) : Liver section, 6 days after exposure to $1/4$ LC_{50} of lead nitrate. (X - 1650).

- Fig.(16) : Liver section, 6 days after exposure to
1/4 LC₅₀ of lead nitrate. (X - 1650).
- Fig.(17) : Liver section, 6 days after exposure to
1/4 LC₅₀ of lead nitrate. (X - 460).
- Fig.(18) : Liver section, 8 days after exposure to
1/4 LC₅₀ of lead nitrate. (X - 660).
- Fig.(19) : Liver section, 8 days after exposure to
1/4 LC₅₀ of lead nitrate. (X - 660).
- Fig.(20) : Liver section, 10 days after exposure to
1/4 LC₅₀ of lead nitrate. (X - 1050).
- Fig.(21) : Liver section, 10 days after exposure to
1/4 LC₅₀ of lead nitrate. (X - 460).
- Fig.(22) : Liver section, 10 days after exposure to
1/4 LC₅₀ of lead nitrate. (X - 460).
- Fig.(23) : Liver section, 2 days after exposure to
1/2 LC₅₀ of lead nitrate. (X - 1650).
- Fig.(24) : Liver section, 4 days after exposure to
1/2 LC₅₀ of lead nitrate. (X - 1650).
- Fig.(25) : Liver section, 6 days after exposure to
1/2 LC₅₀ of lead nitrate. (X - 260).
- Fig.(26) : Liver section, 6 days after exposure to
1/2 LC₅₀ of lead nitrate. (X - 660).
- Fig.(27) : Liver section, 8 days after exposure to
1/2 LC₅₀ of lead nitrate. (X - 660).
- Fig.(28) : Liver section, 8 days after exposure to
1/2 LC₅₀ of lead nitrate. (X - 1650).
- Fig.(29) : Liver section, 10 days after exposure to
1/2 LC₅₀ of lead nitrate. (X - 660).
- Fig.(30) : Liver section, 10 days after exposure to
1/2 LC₅₀ of lead nitrate. (X - 660).

- Fig.(31) : Liver section, 10 days after exposure to $1/2$ LC_{50} of lead nitrate. (X - 165).
- Fig.(32) : Liver section, 2 days after exposure to LC_{50} of lead nitrate. (X - 660).
- Fig.(33) : Liver section, 4 days after exposure to LC_{50} of lead nitrate. (X - 660).
- Fig.(34) : Liver section, 4 days after exposure to LC_{50} lead nitrate. (X - 165).
- Fig.(35) : Liver section, 6 days after exposure to LC_{50} lead nitrate. (X - 165).
- Fig.(36) : Liver section, 6 days after exposure to LC_{50} lead nitrate. (X - 660).
- Fig.(37) : Liver section, 10 days after exposure to LC_{50} lead nitrate. (X - 1650).
- Fig.(38) : T.S. ileum of fish of the control group. (X - 660).
- Fig.(39) : T.S. ileum of fish of the control group. (X - 660).
- Fig.(40) : T.S. ileum of fish of the control group. (X - 660).
- Fig.(41) : T.S. ileum of fish of the control group. (X - 660).
- Fig.(42) : T.S. ileum of fish of the control group. (X - 165).
- Fig.(43) : T.S. ileum of fish of the control group. (X - 1050).
- Fig.(44) : T.S ileum, 2 days after exposure to $1/8$ LC_{50} of lead nitrate. (X - 660).
- Fig.(45) : T.S ileum, 2 days after exposure to $1/8$ LC_{50} of lead nitrate. (X - 1050).

- Fig.(46) : T.S ileum, 2 days after exposure to $1/8$ LC_{50} of lead nitrate. (X - 660).
- Fig.(47) ; T.S ileum, 4 days after exposure to $1/8$ LC_{50} of lead nitrate. (X - 660).
- Fig.(48) : T.S ileum, 6 days after exposure to $1/8$ LC_{50} of lead nitrate. (X - 1050).
- Fig.(49) ; T.S ileum, 8 days after exposure to $1/8$ LC_{50} of lead nitrate. (X - 460).
- Fig.(50) : T.S ileum, 8 days after exposure to $1/8$ LC_{50} of lead nitrate. (X - 660).
- Fig.(51) ; T.S ileum, 10 days after exposure to $1/8$ LC_{50} of lead nitrate. (X - 660).
- Fig.(52) : T.S ileum, 10 days after exposure to $1/8$ LC_{50} of lead nitrate. (X - 660).
- Fig.(53) : T.S ileum, 10 days after exposure to $1/8$ LC_{50} of lead nitrate. (X - 660).
- Fig.(54) : T.S ileum, 2 days after exposure to $1/4$ LC_{50} of lead nitrate. (X - 165).
- Fig.(55) : T.S ileum, 4 days after exposure to $1/4$ LC_{50} of lead nitrate. (X - 260).
- Fig.(56) : T.S ileum, 4 days after exposure to $1/4$ LC_{50} of lead nitrate. (X - 660).
- Fig.(57) : T.S ileum, 4 days after exposure to $1/4$ LC_{50} of lead nitrate. (X - 260).
- Fig.(58) : T.S ileum, 6 days after exposure to $1/4$ LC_{50} of lead nitrate. (X - 1050).
- Fig.(59) ; T.S ileum, 10 days after exposure to $1/4$ LC_{50} of lead nitrate. (X - 460).
- Fig.(60) : T.S ileum, 10 days after exposure to $1/4$ LC_{50} of lead nitrate. (X - 660).

- Fig.(61) : T.S ileum, 2 days after exposure to $1/2$ LC_{50} of lead nitrate. (X - 460).
- Fig.(62) : T.S ileum, 2 days after exposure to $1/2$ LC_{50} of lead nitrate. (X - 460).
- Fig.(63) : T.S ileum, 4 days after exposure to $1/2$ LC_{50} of lead nitrate. (X - 1050).
- Fig.(64) : T.S ileum, 4 days after exposure to $1/2$ LC_{50} of lead nitrate. (X - 1050).
- Fig.(65) : T.S ileum, 6 days after exposure to $1/2$ LC_{50} of lead nitrate. (X - 165).
- Fig.(66) : T.S ileum, 2 days after exposure to LC_{50} of lead nitrate. (X - 660).
- Fig.(67) : T.S ileum, 4 days after exposure to LC_{50} of lead nitrate. (X - 460).
- Fig.(68) : T.S ileum, 6 days after exposure to LC_{50} of lead nitrate. (X - 660).
- Fig.(69) : T.S ileum, 8 days after exposure to LC_{50} of lead nitrate. (X - 460).
- Fig.(70) : T.S ileum, 8 days after exposure to LC_{50} of lead nitrate. (X - 460).
- Fig.(71) : T.S ileum, 10 days after exposure to LC_{50} of lead nitrate. (X - 660).
- Fig.(72) : T.S. of the proper kidney of control fish. (X - 660).
- Fig.(73) : T.S. of the proper kidney of control fish. (X - 660).
- Fig.(74) : T.S. of the proper kidney of control fish. (X - 660).
- Fig.(75) : T.S. of the proper kidney of control fish. (X - 660).

- Fig.(76) : T.S. of the proper kidney of control fish. (X - 660).
- Fig.(77) : T.S. of the proper kidney of control fish. (X - 460).
- Fig.(78) : T.S. of the proper kidney of fish exposed to $1/8$ LC_{50} of lead nitrate for 6 days. (X - 660).
- Fig.(79) : T.S. of the proper kidney of fish exposed to $1/8$ LC_{50} of lead nitrate for 8 days. (X - 660).
- Fig.(80) : T.S. of the proper kidney of fish exposed to $1/8$ LC_{50} of lead nitrate for 10 days. (X - 660).
- Fig.(81) : T.S. of the proper kidney, 4 days after exposure to $1/4$ LC_{50} of lead nitrate. (X - 660).
- Fig.(82) : T.S. of the proper kidney, 8 days after exposure to $1/4$ LC_{50} of lead nitrate. (X - 660).
- Fig.(83) : T.S. of the proper kidney, 10 days after exposure to $1/4$ LC_{50} of lead nitrate. (X - 660).
- Fig.(84) : T.S. of the proper kidney, 2 days after exposure to $1/2$ LC_{50} of lead nitrate. (X - 660).
- Fig.(85) : T.S. of the proper kidney, 6 days after exposure to $1/2$ LC_{50} of lead nitrate. (X - 660).
- Fig.(86) : T.S. of the proper kidney, 10 days after exposure to $1/2$ LC_{50} of lead nitrate. (X - 660).
- Fig.(87) : T.S. of the proper kidney, 4 days after exposure to LC_{50} of lead nitrate. (X - 660).
- Fig.(88) : T.S. of the proper kidney, 6 days after exposure to LC_{50} of lead nitrate. (X - 660).
- Fig.(89) : T.S. of the proper kidney, 8 days after exposure to LC_{50} of lead nitrate. (X - 660).
- Fig.(90) : T.S. of the proper kidney, 8 days after exposure to LC_{50} of lead nitrate. (X - 660).