

**Early Morbidity and Mortality After Second
Step of Staged Management of Extensive
Aortic disease.**

Thesis

Submitted in Complete Fulfillment of
The MD Degree in
Cardiothoracic Surgery

By

AHMAD ABDELSATTAR ABDELAZIZ DAOUD
(M.Sc)

Under Supervision of

PROF. DR. SAMIR ABDALLAH HASSAN
*Professor of Cardiothoracic Surgery,
Faculty of Medicine, Cairo University*

PROF.DR. SAID ABDELAZIZ BADR
*Professor of Cardiothoracic Surgery
Faculty of Medicine, Cairo University*

ASS.PROF.DR. ALAA ELDIN FAROUK MOHAMED
*Assistant Professor of Cardiothoracic Surgery
Faculty of Medicine, Cairo University*

DR. AHMED ABDELAZIZ ELSHARKAWY
*Lecturer of Cardiothoracic Surgery
Faculty of Medicine, Cairo University*

**Faculty of Medicine
Cairo University
2016**

بِسْمِ اللَّهِ الرَّحْمَنِ الرَّحِيمِ

Acknowledgments

First and foremost, thanks are due to ALLAH the most gracious and merciful, to whom I attribute my success in achieving any work in my life.

Second, to the soul of my father, who spent his life pushing me forward.

I would like to express my sincere gratitude and deepest appreciation to my Professor Dr. Samir Abdallah Hassan, Professor of Cardiothoracic Surgery, Faculty of Medicine, Cairo University, for his encouragement and precious guidance throughout the preparation of this work.

I am deeply grateful to Professor Dr. Said Abdelaziz, Professor of Cardiothoracic Surgery, Faculty of Medicine, Cairo University, for his help and comments that enlightened my way throughout this work and throughout my whole clinical life.

I would like also to thank Professor Dr. Alaa Farouk, Assistant Professor of Cardiothoracic Surgery, Faculty of Medicine, Cairo University, for his patience, suggestions and cooperation to achieve this work.

I would like also to thank Dr. Ahmed Elsharkawy, lecturer of cardiothoracic surgery, Faculty of Medicine, Cairo University, for his help and patience all through this work.

Finally, I wish to express my appreciation to all who shared their thoughts and comments with me to achieve this work.

Contents

Contents	1
List of Tables.....	2
List of Figures	3
Abbreviations & Acronyms.....	Error! Bookmark not defined.
Introduction and Aim of the work	6
Review of Literature.....	9
<i>Anatomy of the aorta</i>	10
<i>Epidemiology</i>	13
<i>Classifications</i>	15
<i>Etiology and Pathophysiology</i>	20
<i>Management</i>	27
<i>Indications for Intervention</i>	30
<i>Approach to repair</i>	35
<i>Operative techniques</i>	39
Patients and Methods	63
Results.....	73
Discussion.....	93
Summary and Conclusion	100
References.....	103
المخلص العربى.....	120

List of Tables

Table 1; Normal Adult Thoracic Aortic Diameters	15
Table 2; Suggested Follow-up of Aortic Pathologies after Repair or Treatment ⁷	29
Table 3; pre-operative data.	75
Table 4; Maximum diameters of ascending and descending aortic lesions in MSCT in first stage.....	78
Table 5; Maximum diameter of the aorta in MSCT before second stage (without the cases of coarctation and hypoplastic arch).	82
Table 6; Duration between first and second intervention in surgical and endovascular patients.	84
Table 7; Hospital stay in surgical and endovascular groups of patients.	87
Table 8; ICU stay time in surgical and endovascular patients.....	87
Table 9; Mechanical ventilation time in surgical and endovascular patients.	88
Table 10; Morbidities and mortalities in surgical and endovascular group of patients...	91

List of Figures

Figure 1; Variations in the origin of the aortic arch branches.	10
Figure 2; Anatomy and branches of the abdominal aorta.	11
Figure 3; Crawford's classification of thoracoabdominal aortic aneurysms.	16
Figure 4; Stanford and DeBakey classifications of aortic dissection.	18
Figure 5; Illustration of typical extensive aortic aneurysm of the ascending/arch and thoraco-abdominal aorta repaired in two stages with the elephant trunk technique. ...	27
Figure 6; For endovascular repair descending (type B) aortic dissection, the stent-graft is positioned to cover the tear, while avoiding obstructing flow to the left subclavian ¹⁴² .40	
Figure 7; Four types of endoleaks are seen after an endovascular repair of an abdominal aortic aneurysm.	52
Figure 8; Positioning during thoracoabdominal aortic repair surgery	55
Figure 9; Left heart bypass.....	56
Figure 10; Clamping the aorta during thoracoabdominal aortic repair.	57
Figure 11; Over-sewing the intercostal arteries after aortic transection.....	58
Figure 12; Proximal anastomosis to the Dacron tube during thoracoabdominal aortic repair.	59
Figure 13; Distal anastomosis (open fashion) during thoracoabdominal aortic repair. ..	59
Figure 14; Removal of thrombus and atherosclerotic debris to identify the major branches during thoracoabdominal aortic repair.	60
Figure 15; Organ perfusion during thoracoabdominal aortic operations.	61
Figure 16; The aortic cross-clamp can be moved distally with each successive anastomosis during thoracoabdominal aortic repair.	62
Figure 17; The final anastomosis is completed just above the aortic bifurcation.	62
Figure 18; Large fusiform aneurysm of the descending thoracic aorta.	68
Figure 19; Exposure of the right common femoral artery.	70
Figure 20; Aortogram showing a deployed stent in the proximal descending thoracic aorta followed by a huge aneurysm.	71
Figure 21; Aortogram showing successful deployment of 2 overlapping stents in the descending thoracic aorta.....	72
Figure 22; E-vita thoracic stent graft with 22F delivery system.	72
Figure 23; Gender distribution in surgery and endovascular groups in the second stage.	74
Figure 24; Distribution of patients according to aortic pathology before first stage.	76
Figure 25; extent of the lesion in MSCT before first stage.....	77
Figure 26; interventions in the first stage.	78

Figure 27; aortic pathologies before the second stage.	80
Figure 28; indications of surgery in the second stage.	83
Figure 29; Surgeries in the second stage.	85
Figure 30; Endovascular stenting in the second stage.	86
Figure 31, Neurological complications after second stage.	89
Figure 32; Renal impairment in surgical and endovascular patients (percent to total number of patients).	90

***Introduction
and
Aim of the work***

Introduction

Extensive aortic disease stands for diffuse arteriosclerotic disease, coarctation, aneurysmal dilatation or dissection involving two or more segments of the aorta, which may include the ascending aorta, aortic arch, descending thoracic or abdominal aorta¹.

Patients with extensive aortic diseases are still considered to be a challenge for many cardiovascular surgeons. The relatively proximal segment of the descending aorta (the proximal 1/3 of the descending aorta) can be operated upon via the median sternotomy, but this one-stage approach is often difficult because of poor and limited operative access combined with an increased risk of bleeding from distant sites. Therefore, many surgeons prefer a two-stage approach to treat the combined lesions of the aortic arch and the descending aorta, especially when the extent of the aneurysms is beyond the proximal 1/3 of the descending aorta.²

The introduction of the elephant trunk technique by Borst et al. in 1983 has greatly facilitated surgery on this kind of pathology and this technique has been recognised as the standard modality for treatment of extensive aortic diseases. In a second-stage operation, the elephant trunk can be extended to any desired level through a lateral thoracotomy³.

Since the introduction of endovascular stent-graft technology as a second stage for management of extensive thoracic aortic diseases by Dake et al. in 1998, it has been considered as an alternative treatment modality that may be associated with reduced mortality and morbidity⁴.

Our study is a descriptive both retrospective and prospective non randomized study over 30 patients with extensive aortic disease underwent first stage repair of the aorta and to whom a second stage of repair is done in our centre using surgical approach or endovascular stenting (TEVAR/EVAR) of the diseased part of the descending thoracic, abdominal aorta or arch after de-branching in the first stage.

Aim of the Work

Aim of this study is to report early results of second step of staged repair of extensive aortic disease either by surgery or by endovascular repair of the aorta (TEVAR/ EVAR).

Review of Literature

Anatomy of the aorta

The aorta is the major arterial conduit conveying blood from the heart to the systemic circulation. It originates immediately beyond the aortic valve ascending initially. Then it curves forming the aortic arch, and descends caudally adjacent the spine. The ascending thoracic aorta gives off the coronary arteries. The aortic arch branches are typically the brachiocephalic trunk (branches to the right common carotid and right subclavian arteries), left common carotid and left subclavian arteries; however, aortic arch anatomy is variable (figure 1) ⁵.




Figure 1; Variations in the origin of the aortic arch branches.

(A) and (B) represent the majority of anomalies found in the general population.

(A) Common origin of the left common carotid artery and brachiocephalic artery (bovine arch). Represents 73 percent of all branch variations.

(B) Origin of the left common carotid from the mid- to upper brachiocephalic artery. Represents 22 percent of all branch variations.

(C) Common carotid trunk giving origin to the left subclavian artery.

(D) Common carotid trunk, independent from both subclavian arteries.

(E) Left and right brachiocephalic arteries.

(F) Single arch vessel (brachiocephalic artery) originates the left common carotid and left subclavian arteries.

The descending thoracic aorta provides paired thoracic arteries (T1-T12) and continues through the hiatus of the diaphragm to become the abdominal aorta which extends retroperitoneally to its bifurcation into the common iliac arteries at the level of the fourth lumbar vertebra (figure 2) 5. The abdominal aorta lies slightly left of the midline to accommodate the inferior vena cava which is in close apposition. The branches of the aorta (superior to inferior) include the left and right inferior phrenic arteries, left and right middle suprarenal arteries, the celiac axis, superior mesenteric artery, left and right renal arteries, left and right internal spermatic arteries, inferior mesenteric artery, left and right common iliac artery, middle sacral artery and the paired lumbar arteries (L1-L4).




Figure 2; Anatomy and branches of the abdominal aorta.

The common iliac artery bifurcates into the external iliac and internal iliac arteries at the pelvic inlet. The internal iliac artery gives off branches to the pelvic viscera and also supplies the musculature of the pelvis. The external iliac artery passes beneath the inguinal ligament to become the common femoral artery ⁶.