



1,3277

Comparative Study Between Bupivacaine Combined with Sufentanil, Fentanyl in Continuous Extradural Analgesia During Labor

Thesis Submitted for Partial Fulfillment of M.D. Degree in Anesthesiology

BY

Ashraf Mahmoud Hazem El-Hakim

M.B.B.Ch., M.Sc., Faculty of Medicine, Ain Shams University

Supervised BY

Prof. Dr. Salah El-Din Mostafa El-Halabi

Professor of Anesthesia and Intensive Care Faculty of Medicine - Ain Shams University

Prof. Dr. Bahira Mohamed Tawfik Halmy

Professor of Anesthesia and Intensive Care Faculty of Medicine - Ain Shams University

Prof. Dr. Galal Abou El-Soud Saleh

Assistant Professor of Anesthesia and Intensive Care Faculty of Medicine - Ain Shams University

Dr. Bassem Bolis Ghobrial


Lecturer of Anesthesia and Intensive Care Faculty of Medicine - Ain Shams University


Faculty of Medicine
Ain Shams University
*** 1999 ***





سورة الناء آية ١١٣





Acknowledgment

It was indeed an honor to have been supervised by **Prof. Dr. Salah El-Din Mostafa El-Halabi**, Professor of Anesthesia and Intensive Care, Faculty of Medicine, Ain Shams University. His incessant encouragement and fruitful remarks throughout the different phases of these work were of great value to me, along with the precious time he offered me inspite of his great responsibilities leave me really indebted.

I do feel greatly indebted and grateful to **Prof. Dr. Bahira Mohamed Tawfik**, Professor of Anesthesia and Intensive Care,
Faculty of Medicine, Ain Shams University, for her supervision,
encouragement, help, her valuable support, advice, and precious
guidance which were kindly given.

I would like to express my deepest gratitude to **Dr. Galal Abou El-Soud Saleh**, Assistant Professor of Anesthesia and Intensive Care, Faculty of Medicine, Ain Shams University, for his kind moral support, fruitful supervision, continuous guidance, and valuable advice that are responsible for the production of this work.

I would like to express my deep appreciation to **Dr. Bassem Bolis Ghobrial**, Lecturer of Anesthesia and Intensive Care, Faculty of Medicine, Ain Shams University, for the marvelous help he offered me in completing the clinical part of this work and this work owed a lot to his insight, initiative and support.

Ashraf Hazem

List of Contents

| | PAGE |
|------------------------------|-----------|
| Introduction | 1 |
| Aim of the Work | 2 |
| Review of Literature | 3 – 56 |
| Epidural Space in Pregnancy | 3 – 7 |
| Epidural Narcotics | 8 – 10 |
| Pharmacology of Opioids | 11 – 16 |
| Epidural Local Anesthetics | 17 – 26 |
| Pharmacology of Bupivacaine | 27 – 30 |
| Epidural Anesthesia | 31 – 46 |
| Epidural Analgesia for Labor | 47 – 50 |
| Physiology of Labor Pain | 51 - 56 |
| Patients and Methods | 57 - 62 |
| Results | 63 – 111 |
| Discussion | 112 – 122 |
| Conclusion | 123 |
| Summary | 124 – 125 |
| References | 126 – 137 |
| A rabic Summary | 1 – 2 |



List of Tables

| No. | Table | Page |
|-------------|---|------|
| Table (1): | Relationship between the plasma binding of amide local anesthetics and UV/M ratio. | 24 |
| Table (2): | Pharmacological parameters of anticoagulants and antiplatelet agents. | 42 |
| Table (3): | Pain in labor: Pathways and mechanisms. | 54 |
| Table (4): | Demographic and zero time state characteristics of patients. | 63 |
| Table (5): | The effect of epidural bupivacaine, bupivacaine-fentanyl, and bupivacaine-sufentanil mixture on the onset of analgesia. | 64 |
| Table (6): | The effects of study solutions on sensory level. | 64 |
| Table (7): | The effect of epidural bupivacaine on pain score during labor using a 5-point verbal scale. | 67 |
| Table (8): | The effect of epidural bupivacaine-fentanyl mixture on pain score during labor using a 5-point verbal scale. | 69 |
| Table (9): | The effect of epidural bupivacaine-sufentanil mixture on pain score during labor using a 5-point verbal scale. | 71 |
| Table (10): | The difference in pain sensation by using 5-point verbal scale between the three studied groups. | 73 |
| Table (11): | The association of motor block with the three groups. | 75 |
| Table (12): | The effect of epidural bupivacaine on heart rate (group I). | 78 |
| Table (13): | The effect of epidural bupivacaine and fentanyl on heart rate (group II). | 80 |
| Table (14): | The effect of epidural bupivacaine and sufentanil on heart rate (group III). | 82 |
| Table (15): | The difference in heart rate between the three studied groups. | 84 |
| Table (16): | The effect of epidural bupivacaine on mean arterial blood pressure in mmHg during labor (group I). | 87 |

| Table (17): | The effect of combined epidural bupivacaine and fentanyl on mean arterial blood pressure in mmHg during labor (group II). | 89 |
|-------------|--|------|
| Table (18): | The effect of combined epidural bupivacaine and sufentanil on mean arterial blood pressure in mmHg during labor (group III). | 91 |
| Table (19): | The difference in mean arterial blood pressure in mmHg between the three studied groups. | 93 |
| Table (20): | The difference in respiratory rate during labor between the three studied groups. | 95 |
| Table (21): | Effect of and difference between each of group I, group II and group III as regards maternal SaO ₂ . | 97 |
| Table (22): | Effect of and difference between each of group I, group II and group III as regards maternal tidal volume. | . 99 |
| Table (23): | The association of side effects with the three studied groups. | 102 |
| Table (24): | Difference in fetal heart rate between the three studied groups. | 103 |
| Table (25): | Effect of and difference between each of the three studied groups as regard Apgar score at one minute. | 105 |
| Table (26): | Effect of and difference between each of the three studied groups as regard Apgar score at five minute. | 105 |
| Table (27): | Effect of and difference between each of group I, group II and group III as regard the neurobehavioral response. | 106 |
| Table (28): | Effect of and difference between each of group I, group II and group III as regards PO ₂ , PCO ₂ , pH and base excess in umbilical venous blood. | 107 |
| Table (29): | The results of the post-delivery questionnaire. | 108 |

List of Figures

| No. | Figure | Page |
|--------------|--|------|
| Figure (1): | Sagittal section of lumbar vertebrae to show the principal ligaments. | 3 |
| Figure (2): | Transverse section to show the anatomy and relations to epidural space. | 4 |
| Figure (3): | Molecular shape of fentanyl. | 11 |
| Figure (4): | Molecular shape of sufentanil. | 13 |
| Figure (5): | Chemical structure of bupivacaine. | 18 |
| Figure (6): | Mode of action of local anesthetic drug. | 20 |
| Figure (7): | Schematic diagram illustrating time-dependent changes in sodium channel states and block of sodium channels associated with the cardiac action potential in the presence of a local anesthetic drug. | 21 |
| Figure (8): | Systemic absorption of bupivacaine and lidocaine after extradural injection. | 22 |
| Figure (9): | Schematic diagram of pain pathways during parturition. | 55 |
| Figure (10): | The effect of epidural bupivacaine, bupivacaine-fentanyl, and bupivacaine-sufentanil mixture on the onset of analgesia. | 65 |
| Figure (11): | The effect of epidural bupivacaine on pain score during labor using a 5-point verbal scale. | 68 |
| Figure (12): | The effect of epidural bupivacaine-fentanyl mixture on pain score during labor using a 5-point verbal scale. | 70 |
| Figure (13): | The effect of epidural bupivacaine-sufentanil mixture on pain score during labor using a 5-point verbal scale. | 72 |
| Figure (14): | The difference in pain sensation by using 5-point verbal scale between the three studied groups. | 74 |
| Figure (15): | The association of motor block with the three groups. | 76 |
| Figure (16): | The effect of epidural bupivacaine (group I) on maternal heart rate. | 79 |

| Figure (17): | The effect of epidural bupivacaine-fentanyl mixture (group | |
|--------------|--|-----|
| | II) on heart rate. | 81 |
| Figure (18): | The effect of epidural bupivacaine-sufentanil mixture (group III) on maternal heart rate. | 83 |
| Figure (19): | The difference in heart rate between the three studied groups. | 85 |
| Figure (20): | The effect of epidural bupivacaine (group I) on mean arterial blood pressure in mmHg. | 88 |
| Figure (21): | The effect of epidural bupivacaine-fentanyl mixture (group II) on mean arterial blood pressure in mmHg. | 90 |
| Figure (22): | The effect of epidural bupivacaine-sufentanil mixture (group III) on mean arterial blood pressure in mmHg. | 92 |
| Figure (23): | The difference in mean arterial blood pressure in mmHg between the three studied groups. | 94 |
| Figure (24): | The difference in respiratory rate during labor between the three studied groups. | 96 |
| Figure (25): | Effect of and difference between each of group I, group II and group III as regards maternal SaO ₂ . | 98 |
| Figure (26): | Effect and difference between each of group I, group II and group III as regards maternal tidal volume. | 100 |
| Figure (27): | The association of side effects with the three studied groups. | 102 |
| Figure (28): | Difference in fetal heart rate between the three studied groups. | 104 |
| Figure (29): | Effect of and difference between each of group I, group II and group III as regard the neurobehavioral response. | 106 |
| Figure (30): | The results of post-delivery questionnaire: Overall satisfaction with extradural infusion. | 109 |
| Figure (31): | The results of post-delivery questionnaire: Time to first micturition. | 110 |
| Figure (32): | The results of post-delivery questionnaire: Time to return to full sensation. | 111 |