
THE ROLE OF SOME EGYPTIAN WILD BIRDS AS ENVIRONMENTAL POLLUTERS

THESIS

Submitted in Partial Fulfilment ofthe Requirements for the Degree of Master of Science



By

SAID AHMED MAHMOUD MAZYAD

B.Sc. of Agricultural Sciences (Plant Protection)

Entomology (Ain Shams University)
Diploma in Medical Entomology (Faculty of Science)
Diploma in Environmental Sciences, Department of
Biological Science

To

Institute of Environmental Studies and Research Ain Shams University

16487

1993

THE ROLE OF SOME EGYPTIAN WILD BIRDS AS ENVIRONMENTAL POLLUTERS

. . . .

By

SAID AHMED MAHMOUD MAZYAD

B.Sc. of Agricultural Sciences
(Plant Protection)
Entomology (Ain Shams University)
Diploma in Medical Entomology (Faculty of Science)
Diploma in Environmental Science, Department of
Biological Science

Under the Supervision of

Prof. Dr. WALY A. ASHOUR

Prof. of Plant Pathology, Faculty of Agriculture, Ain Shams University.

Prof. Dr. AHMED SHOUKRY,

Dean of Faculty of Agriculture, Suez Canal University.

Dr. MAHMOUD E. THARWAT,

Associate Professor in Plant Protection Dept., Faculty of Agriculture, Ain Shams University

1993



APPROVAL SHEET

NAME: SAID AHMED MAHMOUD MAZYAD

Title : THE ROLE OF SOME EGYPTIAN WILD BIRDS

AS ENVIRONMENTAL POLLUTERS

This Thesis For The M.Sc. Degree in Environmental Sciences Has Been Approved.

BY

Prof. Dr. AHMED SHOUKRY ABD EL-AZIZ

Prof. of Plant Protection

Dean of Fac. of Agric.

Suez Canal Univ.

A. Stuly

Prof. Dr. AHMED ZAKY ALY

Prof. of Plant Pathology

Agric, Bot ., Dept.

Fac. of Agric .. Zagazig Univ.

3.2.214

Dr. SAWSAN EL-GHAZALI

Assoc. Prof. Community Medicine

Ain Shams Univ., Cairo, Egypt

Dr. MAHMOUD E. THARWAT

Assoc. Prof in Ecology
Plant Protection Dept
Fac of Agric Ain Shams Univ

(1993)

ACKNOWLEDGEMENT

The author is very grateful to Prof. Dr. WALY EL-DIN ASHOUR, Professor of Plant Pathology, Faculty of Agriculture, Ain Shams University and Prof. Dr. AHMED SHOUKRY, Dean of Faculty of Agriculture, Suez Canal University, for their valuable supervision and guidance.

Thanks are also due to Dr. MAHMOUD THARWAT, Associate Professor in Plant Protection Department, Faculty of Agriculture, Ain Shams University for his help and assistance during this work.

Many sincere thanks to Dr. MOHAMED EL ZAYAT., Head of Environmental Department, Institute of Environmental Studies and Research, Ain Shams University, for his continuous guidance, discussions and careful revision of all points investigated in this study.

Thanks are due to Dr. Amany Mostafa, Dr. Fatehia Mohamed and Dr. Zomored Anwar for their sincere help in assisting the author during this work. The writer would like to express his deep thanks to all members of Microbiology Department, Faculty of Medicine, Ain Shams University and the Institute of Animal Health, for their encouragement and supplying the facilities.

I would like to acknowledge all staff members of the research and Training Center on Vectors of Diseases for their continuous encouragement.

LIST OF TABLES

ţ

Table 1	Seasonal fluctuation of temperature and relative humidity in Qalubia and Giza Governorates from January to December, 1989.
Table 2:	Seasonal fluctuation of temperature and relative humidity in Qalubia and Giza governorates from January to December, 1990.
Table 3:	Air temperature in great Cairo (Cairo, Giza and Qalubia) during the period from January to December, 1989.
Table 4a	: Meteorological records of El-Arish City (North Sinai Governorate) during the study period from (January, 1989 to December, 1989).
Table 4b	Governorate) during the study period from January, 1990 to December, 1990.
Table 5:	Pathogenic and non pathogenic bacteria isolated from the excreta of wild birds at South Delta and North Sinai.
Table 6:	Biochemical differentiation between bacterial species recovered from the excreta of wild birds collected from the South Delta and North Sinai.
Table 7:	Internal pathogenic bacteria isolated from wild bird collected from El-Kanater El-Khairya, Qalubia, governorate during the study period (Jan., 1989 to Dec., 1989).

	Page
Table 8: Internal pathogenic bacteria isolated from wild birds collected from El-Kanater El-Khairya Qalubia	
Governorate during Jan., 1990 to Dec., 1990.	71
Table 9: Internal pathogenic bacteria isolated from wild birds collected from El-Marg area (Qalubia Governorate) during the study period (Jan., 1989 to Dec., 1989).	<i>7</i> 5
Table 10: Internal pathogenic bacteria isolted from wild birds collected from El-Marg area (Qalubia Governorate) during Jan., 1990 to Dec. ,1990.	78
Table 11: Internal pathogenic bacteria isolated from the excreta of wild birds living at El-Giza Governorate (South Delta area during the study period (January. 1989 to December, 1989).	82
Table 12: Internal pathogenic bacteria isolated from the excreta of wild birds living at El-Giza Governorate (South Delta area) during the study period (January, 1990 to December, 1990).	85
Table 13: Internal pathogenic bacteria isolated from wild bird as migratory birds collected from El-Arich City during the study period (January 1989 to December 1990).	
Table 14: Internal pathogenic bacteria isolated from wild birds, as migratory birds collected from El-Sheikh Zoued during the study period (January, 1989 to December, 1990).	93
Table 15: Internal pathogenic bacteria isolated from wild birds, as migratory birds, collected from El-Bardaweal during the study period (Jan. 1989 to Dec. 1990).	96

Table 1	6: Percentage of Salmonella detected in wild birds collected from El-Kanater El-Khairya during the studying period.
Table 17	Percentage of recovery rate of Salmonella detected in wild birds collected from El-Marg during the studying period.
Γable 18	Percentage of recovery rate of Salmonella detected in wild birds collected from El-Giza Governorate (South Delta area)
Table 19	serotypes of Salmonella recovered from 5 wild bird species execretion on various selective media using tetrathionate enrichement media.
Table 20	3: Salmonella serotypes recovered from 5 wild bird species excreta on various selective media using specific enrichment media.
Γable 21	: Type of pathogenic bacteria isolated from different water samples.
Γable 22	: Effect of pathogenic organisms infesting migratory birds in North Sinai
Γable 23	: Percentage of H.A. and H.I. tests carried out on egg fluids using ND antiserum as reference (virology

LIST OF FIGURES

Fig. 1:	Descriptive map of the first area (South Delta) showing the place of collection of the wild birds.
Fig. 2:	Temperature and relative humidity recorded at Qalubia and Giza Governorates from January, 1989 to December, 1989.
Fig. 3:	Temperature and relative humidity recorded at Qalubia and Giza Governorates from January, 1990 to December, 1990.
Fig. 4:	Air temperature in Great Cairo (Cairo, Giza and Qalubia) during the period from January to December, 1989,
Fig. 5:	Descriptive map of the second area (North Sinai) showing the place of collection of migratory birds.
Fig. 6:	Temperature and relative humidity recorded at El-Arish City (North Sinai Governorate) during the study period from January, 1989 to December, 1989.
Fig. 7:	Temperature and relative humidity recorded at El-Arish City (North Sinai Governorate) during the study period from January, 1990 to December, 1990.
Fig. 8:	Distribution of pathogenic bacteria isolated from excreta collected from wild birds at El-Kanater El-Khairya (samples were collected from January, 1989 to December, 1989).
Fig. 9:	Relative distribution of pathogenic bacteria isolated from excreta collected from El-Kanater El-Khairya during

Fig. 10:	Distribution of pathogenic bacteria isolated from excreta collected from wild birds living at El-Kanater El-Khairya (samples were collected from January, 1990 to December, 1990).	
Fig. 11:	Relative distribution of pathogenic bacteria isolated from excreta collected from January, 1990 to December 1990. (samples were collected from El-Kanater El-Khairya area).	
Fig. 12:	Distribution of pathogenic bacteric isolated from excreta collected from wild birds living at El-Marg (samples were collected from January, 1989 to December, 1989).	
Fig. 13:	Relative distribution of pathogenic bacteria isolated from excreta collected from El-Marg during 1989.	
Fig. 14:	Distribution of pathogenic bacteria isolated from excreta collected from wild birds living at El-Marg (samples were collected from January, 1990 to December, 1990).	
Fig. 15:	Relative distribution of pathogenic bacteria isolated from excreta collected from wild birds living at El Marg during 1990.	
Fig. 16:	Distribution of pathogenic bacteria isolated from excreta collected from wild birds living at El-Giza (samples were collected from January, 1989 to December, 1989).	
Fig. 17:	Relative distribution of pathogenic bacteria isolated from wild birds at El Giza area during the year of 1989.	

Fig. 18: Di	istribution of pathogenic bacteria isolated from excreta collected from wild birds living at El-Giza (samples were collected from January, 1990 to December, 1990).
Fig. 19: R	Relative distribution of pathogenic bacteria isolated from wild birds at El-Giza area during 1990 .
Fig. 20: D	istribution of bacteria isolated from excreta collected from wild birds living at El Arish (1989-1990)
Fig. 21:	Relative distribution of pathogenic bacteria isolated from excreta collected from wild migratory birds at El Arish City (Samples were collected from Jan., 1989 to Dec., 1990).
_	stribution of pathogenic bacteria isolated from excreta collected from wild birds living at Sheikh Zoued 1989-1990.
	elative distribution of pathogenic bacteria isolated from excreta collected from wild migratory birds at Sheikh Zoued (samples were collected from Jan., 1989 to Dec., 1990).
-	stribution of pathogenic bacteria isolated from excreta collected from wild birds living at El-Bardaweel (1989-1990)
	elative distribution of pthogenic bacteria isolated from excreta collected from wild birds migratory at El Bardaweel (samples were collected from Jan., 1989 to Dec., 1990).
	ercentage of Newcastle disease shows the tests of H.I. and H.A. (virology test) . (samples were collected from migratory birds at North Sinai).

TABLE OF CONTENTS

	Page
ACKNOWLEDGEMENT	i
List of Tables	ii
List of Figures	iv
INTRODUCTION	1
REVIEW OF LITERATURE	3
- Bacterial disease	4
- Viral disease	12
MATERIAL AND METHODS	24
I. The study areas	24
A. South of Delta (First area)	24
1- Description of the first area	24
2- The common birds in the first area	26
3- Climatic condition in the first area	28
B. North Sinai (Second area)	35
1- Description of the second area	35
2- Topography of Sinai peninsula	35
3- Climate of Sinai peninsula	37
4- Phytogeography	37
5- Human population	38
6- Water sources	38
7- Meteorological records of North Sinai during the study	
period (January, 1989 to December, 1990)	39
II. The birds under test	44
1. Cattle egret (Bubulcus ibis)	44
2. Quail (Coturnix coturnix)	44
3. Brown-necked raven (Corvus corax ruficollis)	44
4. House sparrow (Passer domesticus)	44

	Page
5. Starling (Sturnus vulgaris)	45
6. Turtle dove (Streptopelia turtur)	45
7. Coot (Fulica atra)	45
8. Egyptian goose (Alopochen agypticus)	45
9. Teal (Anas crecca)	45
10. Pintail (Anas acuta)	45
III. Sample collection	46
A. From birds	4 6
B. From sources of water	46
IV. Laboratory studies	47
A Bacteriological investigation	47
1- Preparation of samples	47
2- Isolation and identification of pathogenic and non	
pathogenic bacteria	48
i. Charbohydrate fermentation test.	48
ii. Isolation and purification of the colonies.	48
iii. Biochemical identification	49
1. Sugar fermentation reaction	49
2. Indole test	49
3. Methyl red reaction	50
4. Vogos-proskawer reaction	50
5. Citrate utilization	50
6. Gelatin liquification test	51
7. Urea test	51
iv. Salmonella isolation and identification	51
B. Examination of water samples	52
C. Plant inoculation with pathogenesis	53
D. Virology test	54

	Page
E. Description of the used media	56
i. The media of bacteria	56
ii. Virology media	61
- RESULTS	64
- DISCUSSION	112
- CONCLUSION AND RECOMMENDATION	118
- SUMMARY	120
- REFERENCES	123
- ARABIC SUMMARY	

INTRODUCTION AND AIM OF THE WORK