

SOME NEW APPROACHES TO PROTECT AGRICULTURAL ENVIRONMENT IN QENA REGION (UPPER EGYPT) AGAINST SOME INSECT PESTS

BY



72820

MOURTADA AHMED ALI AHMED EISSA

B.Sc. Agric. Sc. (Plant Protection), Assiut Univ., 1979

M. Sc. Agric. Sc. (Economic Entomology), Al-Azhar Univ., 1986

THESIS

**Submitted in Partial Fulfillment
of Requirement for the Degree of
DOCTOR OF PHILOSOPHY**

632-7
H. A

**In
Environmental Science
(Agricultural Science)**



**Faculty of Agriculture
Department of Agricultural Sciences
Institute of Environmental Sciences
Ain Shams University**

SOME NEW APPROACHES TO PROTECT AGRICULTURAL ENVIRONMENT IN QENA REGION (UPPER EGYPT) AGAINST SOME INSECT PESTS

BY

MOURTADA AHMED ALI AHMED EISSA
B.Sc. Agric. Sc. (Plant Protection), Assiut Univ., 1979
M. Sc. Agric. Sc. (Economic Entomology), Al-Azhar Univ., 1986

This Thesis for the Ph. D. degree has been approved by :

Prof. Dr. Galal Mahmoud Moawad
Head Researcher and Director, Plant Protec.
Res. Inst., Agric. Res. Centre, Min. of
Agriculture

(*Galal Moawad*)

Prof. Dr. Mohamed Ibrahim Abdel-Megied
Prof. of Pesticides, Fac. of Agric., Ain Shams
Univ.

(*M. Abdel Megied*)

Prof. Dr. Zidan Hendy Abdel-Hameid
Prof. of Pesticides and Vice Dean, Fac. of
Agric., Ain Shams Univ. (Supervisor)

(*Zidan*)

Date of Examination : 10 / 4 / 1997



SOME NEW APPROACHES TO PROTECT AGRICULTURAL ENVIRONMENT IN QENA REGION (UPPER EGYPT) AGAINST SOME INSECT PESTS

MOURTADA AHMED ALI AHMED EISSA
B.Sc. Agric. Sc. (Plant Protection), Assiut Univ., 1979
M. Sc. Agric. Sc. (Economic Entomology), Al-Azhar Univ., 1986

Under Supervision of

Prof. Dr. Zidan Hendy Abdel-Hameid

Prof. of Pesticides and Vice Dean, Fac. of Agric., Ain Shams Univ.

Prof. Dr. Ahmed Ismail Gadallah

Prof. of Economic Entomology and Head of Plant Protection Dept., Fac. of Agric., Al-Azhar Univ.

Prof. Dr. Ahmed Abdou Hamed Amin

Head Researcher and Head of Cutworms and Mole-cricket Dept., Plant Protec., Res., Inst., Agric. Res. Centre

ABSTRACT

The mainly goal of this study is the protection of agricultural environment in Qena Governorate using certain new approaches. Studies concern with the plant coverage and general survey of insects persiste in this area of Egypt using light traps throughout the period from April, 1992 to March, 1994. In this study, 41448 insect species were surveyed. Studies on the ecology and control of the most economic pests demonated in Qena were conducted. These Pests are *A. ipsilon*, *S. exigua* and *G. africana* which cause severe damage for field crops, vegetables and medical and ornamentals.

CONTENTS

	Page
ACKNOWLEDGEMENT	
INTRODUCTION	1
REVIEW OF LITERATURE	2
1. Studies on <i>Agrotis ipsilon</i>	2
1.1. Economic importance of <i>A. ipsilon</i> (Hufn.)	2
1.2. Population dynamics and seasonal abundance of the black cutworm correlated with thermal summations ...	4
1.3. Chemical control of the black cutworm <i>A. ipsilon</i>	10
2. The seasonal activity and abundance of <i>Spodoptera</i> <i>exigua</i> (Hb.)	17
3. Ecological studies on the mole-crickets	22
3.1 Economic importance and behaviour of mole-crickets ...	22
3.2. Mole-crickets' activity and population fluctuations	27
3.3. The Role of Mole-cricket Trapping As Means of Its Control	28
MATERIAL AND METHODS	32
1. Survey, seasonal abundance and population dynamics of certain pests in Qena Governorate (Upper Egypt)	32
2. Generation and prediction possibility of adult cutworm, <i>Agrotis ipsilon</i> (Hufn.) in relation to thermal unit accumulations	34
3. Relation between crop rotation regime and some insect pests infestation at Qena Governorate	36
4. Control of cutworm, <i>Agrotis ipsilon</i> (Hufn.)	37
4.1. Laboratory tests	37
4.2. Efficacy of certain insecticides on <i>A. ipsilon</i> under field conditions	40
5. Controlling Mole-crickets by using pitfall trap	42
	44
RESULTS AND DISCUSSION	
I: Survey of green vegetation and insect orders	44
I.1. Green cover	44

CONTENTS Cont.,

	Page
I.2. Insect survey	48
I.2.1 Order Lepidoptera	49
I.2.2. Order Orthoptera	52
I.2.3. Order Neuroptera	52
I.2.4. Order Odonata	55
I.2.5 Order Coleoptera	55
I.2.6. Order Hemiptera	55
I.2.7. Order Dictyoptera	57
II : Ecological studies on some key insect pests in Qena Governorate	59
II.1. Ecological studies on the black cutworm (BCW), <i>A. ipsilon</i>	59
II.1.1. Seasonal abundance and population dynamics	59
II.1.2. Population density and field generations	63
II.1.2.1 Broods occurrence	70
II.1.2.2. Generations times	73
II.1.3. Prediction of <i>A. ipsilon</i> moths emergence using the integration between Icob (1977) method and Luckman (1976) formula	80
II.1.4. The effects of some weather factors on the activity and abundance of the BCW at Qena Governorate	82
II.1.4.1 Effect of day maximum temperature (D.Mx.T.)	82
II.1.4.2. Effect of night minimum temperature (D.Mn.T.)	84
II.1.4.3. Effect of day mean relative humidity (R.H. %)	84
II.1.4.4. Effect of the day wind velocity	84
II.1.4.5. The combined effect of the four weather factors on the seasonal activity of <i>A. ipsilon</i>	85
II.1.5. Cutworm infestation and crop rotation	87
II.1.5.1. Sesame/bean/maize/wheat rotation	87
II.1.5.2. Okra/squash/tomato/pepper/onion rotation	90
II.2. Ecological studies on lesser cotton leafworm, <i>Spodoptera exigua</i> (Hb.)	94
II.2.1. The fluctuations in the seasonal abundance of <i>S. exigua</i> associated by light trap catches	94



CONTENTS Cont.,

	Page
II.2.2. Effect of mean weather factors on the activity and abundance of <i>Spodoptera exigua</i> (Hb.) in Qena Governorate	111
II.2.2.1 Effect of day maximum temperature	111
II.2.2.2. Effect of the night minimum temperature	113
II.2.2.3. Effect of day mean relative humidity	113
II.2.2.4. Effect of the day mean wind velocity	114
II.2.2.5. The combined effect of the four studied weather factors on the seasonal activity of <i>S. exigua</i>	114
II.2.3. The lesser cotton leafworm infestation and crops rotation	117
II.2.3.1. The first rotation (a field crops rotation system)	117
II.2.3.2. The second rotation (vegetables)	120
II.3. Ecological Studies on Mole Cricket, <i>Gryllotalpa africana</i>	124
II.3.1. Seasonal abundance of <i>Gryllotalpa africana</i> adults as indicated by light trap catches	124
II.3.1.1. Nage Hammadi district	130
II.3.1.2. Qena District	1130
II.3.2. Analysis of the effect of the considered main weather factors on the seasonal activity of <i>G. africana</i> adults ..	133
II.3.2.1 Effect of day maximum temperature	133
II.3.2.2. Effect of the night minimum temperature	133
II.3.2.3. Effect of day mean relative humidity	139
II.3.2.4. Effect of wind velocity	139
II.3.2.5. Combined effect of the four weather factors on the seasonal activity of <i>G. africana</i>	140
II.3.3. Mole-cricket infestation and crop rotation	142
II.3.3.1. The first regime (field crops)	142
II.3.3.2. The second regime (vegetable crops)	145
III : Control of Some Insect Pests	150
III.1. Effect of certain pesticides against <i>A. gossypii</i>	150
III.1.1. Acute toxicity of certain pesticides	150
III.1.2. Lethal effect of certain pesticides on <i>A. gossypii</i> 1 st instar larvae	154

CONTENTS Cont.,

	Page
III.1.3 Latent biological effect of pyriproxyfen Jovenoid on <i>A. ipsilon</i>	157
III.1.3.1. Larval and pupal weight	157
III.1.3.2. Pupation, adult emergence, sex ratio, egg deposition and hatchability	157
III.1.4. Latent biological effect of prempet on <i>A. ipsilon</i>	164
III.1.4.1. Larval and pupal weight	164
III.1.4.2. Pupation, adult emergence, sex ratio, egg deposition and hatchability.	164
III.1.5. Latent biological effect of fenvalerate on <i>A. ipsilon</i>	169
III.1.5.1. Larval and pupal weight	169
III.1.5.2. Pupation, adult emergence, sex ratio, egg deposition and hatchability	174
III.1.6. Latent biological effect of cyanophos on <i>A. ipsilon</i>	179
III.1.6.1. Larval and pupal weight.	179
III.1.6.2. Pupation, adult emergence, sex ratio, egg deposition and hatchability.	179
III.1.7. Field efficacy and performance of tested insecticides against <i>A. ipsilon</i> in Qena Governorate	189
III.2. Performance and efficacy of pitfall traps in reducing the population of mole-cricket in pepper field	194
 DISCUSSION	 202
SUMMARY	210
REFERENCES	214
 ARABIC SUMMARY	

LIST OF TABLES

Table No :	Page
1 Summary crops cultivated in the experimental area during 1992-1993 and 1993-1994, seasons, Qena Governorate	46
2 Wintery crops cultivated in the experimental area during 1992-1993 and 1993-1994, seasons, Qena Governorate	47
3 Fruit and wooden trees cultivated in the experimental area during 1992-1993 and 1993-1994, seasons, Qena Governorate	48
4 Total numbers of class insects captured adults as indicated by light traps in Qena Governorate, 1992/1993 and 1993/1994	50
5 Total numbers of Lepidopterous families and species as indicated by light traps in Qena Governorate, 1992/1993 and 1993/1994	53
6 Total numbers of Orthopterous sub-orders families and species as indicated by light traps in Qena Governorate, 1992/1993 and 1993/1994	54
7 Total numbers of Neuropterous families and species as indicated by light traps in Qena Governorate, 1992/1993 and 1993/1994	54
8 Total numbers of Odonatous families and species as indicated by light traps in Qena Governorate, 1992/1993 and 1993/1994	56

LIST OF TABLES Cont.,

Table No :	Page
9 Total numbers of Coleopterous families and species as indicated by light traps in Qena Governorate, 1992/1993 and 1993/1994	56
10 Total numbers of Hemipterous sub-orders families and species as indicated by light traps in Qena Governorate, 1992/1993 and 1993/1994	58
11 Total numbers of Dictyopterous families and species as indicated by light traps in Qena Governorate, 1992/1993 and 1993/1994	58
12 Fluctuation in the monthly of <i>A. ipsilon</i> moths as indicated by two light traps in Qena Governorate 1992/1993 and 1993/1994	60
13 Seasonal changes in <i>A. ipsilon</i> moths caught in light traps in Qena Governorate 1992/1993 and 1993/1994 ..	64
14 Weekly catches of black cutworm <i>A. ipsilon</i> in a light trap in Qena Governorate and the corresponding averages of 4 weather factors (1992/1993)	66
15 Weekly catches of black cutworm <i>A. ipsilon</i> in a light trap in Qena Governorate and the corresponding averages of 4 weather factors (1993/1994).....	68
16 Generation time of BCW adopted by Icob (1977), Qena Governorate, 1992 - 93 and 1993 - 94	76
17 Timing of peaks occurrence of BCW moths as indicated by heat units summation, Qena Governorate, 1992 - 93 and 1993 - 94	76

LIST OF TABLES Cont.,

Table No :	Page
18 Simple correlation (r), simple regression (b) and partial regression (Pb) values indicating the relation between and effect of 4 weather factors on the weekly catches of <i>A. ipsilon</i> in Qena Governorate during 92/93 & 93/94 .	83
19 Analysis of variance of the seasonal activity of <i>A. ipsilon</i> on four main weather factors along with percentage of explained variance, Qena Governorate (1992/93 & 1993/94)	86
20 Number of cutworm larvae/100 plants of different crops cultivated in a 2-year rotation system in Qena Governorate between April, 1992 and April, 1994	88
21 Number of cutworm larvae/100 plants of different crops (mainly vegetables) cultivated in Qena Governorate between April, 1992 and April, 1994	91
22 Fluctuation in the monthly numbers of <i>S. exigua</i> moths as indicated by two light traps in Qena Governorate 1992/1993 and 1993/1994	95
23 Weekly catches of lesser cotton leafworm, <i>S. exigua</i> in a light trap in Qena Governorate and the corresponding averages of 4 weather factors (1992/1993)	98
24 Weekly catches of lesser cotton leafworm, <i>S. exigua</i> in a light trap in Qena Governorate and the corresponding averages of 4 weather factors (1993/1994)	100
25 Seasonal changes in <i>S. exigua</i> moths caught in light traps in Qena Governorate 1992/1993 and 1993/1994 ..	104
26 Generation occurrence of <i>S. exigua</i> according to heat units summation in qena Governorate, 1992-1994	111

LIST OF TABLES Cont.,

Table No :	Page
27 Simple correlation (r), simple regression (b) and partial regression (Pb) values indicating the relation between and effect of 4 weather factors on the weekly catches of <i>S. exigua</i> Qena Governorate during 92/93 & 93/94 .	112
28 Analysis of variance of the seasonal activity of <i>S. exigua</i> on four main weather factors along with percentage of explained variance, Qena Governorate (1992/93 & 1993/94)	115
29 Number of lesser cotton leafworm larvae/100 plants of different crops cultivated in a 2-year rotation system in Qena Governorate between April, 1992 and April, 1994	118
30 Number of lesser cotton leafworm larvae/100 plants of different crops (mainly vegetables) cultivated in Qena Governorate between April, 1992 and April, 1994	121
31 Fluctuation in the monthly numbers of <i>G. africana</i> adults as indicated by two light traps in Qena Governorate during 1992/1993 and 1993/1994	125
32 Seasonal changes in <i>G. africana</i> adults caught in light traps in Qena Governorate 1992/1993 and 1993/1994 ..	128
33 Weekly catches of african mole-cricket, <i>G. africana</i> in a light trap in Qena Governorate and the corresponding averages of 4 weather factors (1992/1993)	134
34 Weekly catches of african mole-cricket, <i>G. africana</i> in a light trap in Qena Governorate and the corresponding averages of 4 weather factors (1993/1994)	136

LIST OF TABLES Cont.,

Table No :	Page
35 Simple correlation (r), simple regression (b) and partial regression (Pb) values indicating the relation between and effect of 4 weather factors on the weekly catches of <i>G. africana</i> Qena Governorate during 92/93 & 93/94 .	138
36 Analysis of variance of the seasonal activity of <i>G. africana</i> on four main weather factors along with percentage of explained variance, Qena Governorate (1992/93 & 1993/94)	141
37 Number of mole-cricket galleries/40 m ² in the soil of different crops cultivated in a 2-year rotation system in Qena Governorate between April, 1992 and April, 1994	143
38 Number of mole-cricket galleries/40 m ² in the soil of different crops (mainly vegetables) cultivated in Qena Governorate between April, 1992 and April, 1994	146
39a Estimation of Ld-P lines obtained by exposure of <i>A. ipsilon</i> 4 th instar larvae to certain pesticides	151
39b "t" values for differences in LC ₅₀ values between pyriproxyfen and other toxicants	151
40 Effect of Pyriproxyfen 10% (JHM) on the black cutworm, <i>A. ipsilon</i> 4 th instar larvae	158
41 Effect of Prempet (mixture of Pyriproxyfen 5% and fenpropathrin 15 %) on the black cutworm, <i>A. ipsilon</i> 4 th instar larvae	165
42 Effect of fenvalerate 10% on the black cutworm, <i>A. ipsilon</i> 4 th instar larvae	172

LIST OF TABLES Cont.,

Table No. :	Page
43 Effect of Cyanophos on the black cutworm, <i>A. ipsilon</i> 4th instar larvae	180
44 chronic toxicity of certain conventional insecticides against <i>A. ipsilon</i>	187
45 Effect of certain pesticides on the black cutworm, <i>A. ipsilon</i> in maize field, Qena Governorate, 1994	190
46 Daily number of trapped mole-cricket insects by pitfall traps in Pepper field, Qena Governorate, 1993	195
47 Number of mole-cricket insects captured in Pepper field by pitfall traps as a control method, Qena Governorate, 1993	198

LIST OF FIGURES

Figure No :	Page
1 Percentage of total catch for different surveyed insect orders during 1992/1993 and 1993/1994 in Qena Governorate	51
2 Percentage of <i>A. ipsilon</i> moths total catch during successive months of 1992/1993 in Qena Governorate	61
3 Percentage of <i>A. ipsilon</i> moths total catch during successive months of 1993/1994 in Qena Governorate	62
4 Seasonal abundance of cutworm, <i>A. ipsilon</i> based on percentages of annual total catch at Qena Governorate during the two successive years of 92/93 and 93/94	65
5 Fluctuation in the population density of BCW, <i>A. ipsilon</i> at Nage Hammadi and Qena districts, Qena Governorate, 1992/1993	71
6 Fluctuation in the population density of BCW, <i>A. ipsilon</i> at Nage Hammadi and Qena districts, Qena Governorate, 1993/1994	72
7 Number of BCW field generations, according to Audemard & Millair (1975) and Isob (1977) method at Nage-Hammadi district, Qena Governorate, 1992/1993	74
8 Number of BCW field generations, according to Audemard & Millair (1975) and Isob (1977) method at Nage-Hammadi district, Qena Governorate, 1993/1994	75
9 Number of BCW field generations, according to Audemard & Millair (1975) and Isob (1977) method at Qena district, Qena Governorate, 1992/1993	78