

D

HEPATIC INVOLVEMENT IN TYPHOID FEVER AMONG EGYPTIAN INFANTS AND CHILDREN

Thesis

Submitted in partial fulfillment of
the Master Degree in **Pediatrics**

By



Mahmoud Mohamed Mahmoud Madian

M.B., B.Ch. (Ain Shams)

Supervised By

Prof. Dr. Sanaa Youssef Shaaban

Prof. of Pediatric

Faculty of Medicine

Ain Shams University

Dr. Moustafa Abdel-Aziz Sayed El-Hodhod

Lecturer of Pediatrics

Faculty of Medicine

Ain Shams University

**Faculty of Medicine
Ain Shams University**

1997





Acknowledgement

First of all I would like to thank God.

I wish to express my most sincere and heartfelt gratitude to Dr. Sanaa Youssef Shaaban, Prof. of Pediatrics, Faculty of Medicine, Ain Shams University, not only because this work would have never seen the light without her generous support and guidance, but also because she taught me how to be serious, systematic and scientific.

I'm grateful to Dr. Moustafa Abdel-Aziz El Hodhod, Lecturer of Pediatrics, Ain Shams University, for his unfailing support and continuous encouragement for me throughout this study.

I would like to thank my colleagues at the Abbassia Fever Hospital who participated or helped me in one way or the other to complete this work.

Last but not the least, I would like to thank my patients and their families for accepting to undergo this study.

Mahmoud Madian

CONTENTS

	Page
List of Tables	ii
List of Figures	vi
List of Abbreviations	vii
Introduction and aim of the work	1
Review of Literature	
Definition	3
Epidemiology of typhoid fever	3
Pathology and pathogenesis	9
Immunity of typhoid fever	16
Prevention and control	19
Clinical picture of enteric fever	24
Typhoid in children	29
Complications of enteric fever	30
Mortality in typhoid fever	40
Diagnosis of typhoid fever	41
Treatment of typhoid fever	51
The liver in typhoid fever	64
Subjects and Methods	67
Results	77
Discussion	117
Summary, Conclusion and recommendations	133
References	136
Arabic Summary	

LIST OF TABLES

Tab. No.	Title	Page
1	Biochemical differences between <i>S. typhi</i> , <i>S. paratyphi A</i> and <i>S. schottmuelleri</i> .	4
2	Serologic classification using the kauman-white agglutination scheme of antigenic analysis is summarized.	5
3	Approximate mean liver span of infants and children in mid-clavicular line.	74
4	Comparison between patients at the end of 1st week and at the end of 2nd week of illness regarding the symptoms.	77
5	Comparison between patients at the end of 1st week and at the end of 2nd week of illness regarding the signs.	79
6	Comparison between patients at the end of 1st week of illness and controls regarding the TLC.	81
7	Comparison between patients at the end of 1st week of illness and controls regarding the ESR.	82
8	Comparison between patients at the end of 1st week of illness and controls regarding total serum bilirubin.	83
9	Comparison between patients at the end of 1st week of illness and controls regarding AST.	84
10	Comparison between patients at the end of 1st week of illness and controls regarding ALT.	85
11	Comparison between patients at the end of 1st week of illness and controls regarding ALP.	86
12	Comparison between patients at the end of 2nd week of illness and controls regarding TLC.	87

13	Comparison between patients at the end of 2nd week of illness and controls regarding ESR.	88
14	Comparison between patients at the end of 2nd week of illness and controls regarding total serum bilirubin.	89
15	Comparison between patients at the end of 2nd week of illness and controls regarding AST.	90
16	Comparison between patients at the end of 2nd week of illness and controls regarding ALT.	91
17	Comparison between patients at the end of 2nd week of illness and controls regarding ALP.	92
18	Comparison between patients at the end of 1st week and at the end of the 2nd week of illness regarding the pulse rate.	93
19	Comparison between patients at the end of 1st week and at the end of the 2nd week of illness regarding the temperature.	94
20	Comparison between patients at the end of 1st week and at the end of the 2nd week of illness regarding the Hb .	96
21	Comparison between patients at the end of 1st week and at the end of the 2nd week of illness regarding ESR.	97
22	Comparison between patients at the end of 1st week and at the end of the 2nd week of illness regarding TLC.	98
23	Comparison between patients at the end of 1st week and at the end of the 2nd week of illness regarding total serum bilirubin.	100
24	Comparison between patients at the end of 1st week and at the end of the 2nd week of illness regarding AST.	101

25	Comparison between patients at the end of 1st week and at the end of the 2nd week of illness regarding ALT.	102
26	Comparison between patients at the end of 1st week and at the end of the 2nd week of illness regarding ALP.	103
27	Comparison between patients at the end of 1st week and at the end of the 2nd week of illness using cut off values regarding liver function tests.	105
28	Relation between some clinical parameters and serum bilirubin level among cases with typhoid fever at the end of the first week of illness.	107
29	Relation between some clinical parameters and AST level among cases with typhoid fever at the end of the first week of illness.	108
30	Relation between some clinical parameters and ALT level among cases with typhoid fever at the end of the first week of illness.	109
31	Relation between some clinical parameters and alkaline phosphatase level among cases with typhoid fever at the end of the first week of illness.	110
32	Relation between some clinical parameters and serum bilirubin level among cases with typhoid fever at the end of the second week of illness.	111
33	Relation between some clinical parameters and AST level among cases with typhoid fever at the end of the second week of illness.	112
34	Relation between some clinical parameters and ALT level among cases with typhoid fever at the end of the second week of illness.	113

35	Relation between some clinical parameters and alkaline phosphatase level among cases with typhoid fever at the end of the second week of illness.	114
36	Correlation coefficients between different studied parameters among cases at the end of the 1st week of illness.	115
37	Correlation coefficients between different studied parameters at the end of the second week of illness.	116

LIST OF FIGURES

Fig. No.	Title	Page
1	Comparison between patients at the end of 1st week and at the end of 2nd week of illness regarding the symptoms.	78
2	Comparison between patients at the end of 1st week and at the end of the 2nd week of illness regarding the signs.	80
3	Comparison between typhoid patients at the end of the first week and at the end of the second week of illness regarding pulse rate and temperature according to the mean value.	95
4	Comparison between typhoid patients at the end of the first week and at the end of the second week of illness regarding mean of hematological values ESR, Hb and TLC.	99
5	Comparison between controls, typhoid patients at the end of the first week and at the end of the second week of illness regarding mean of liver function tests.	104
6	Comparison between patients at the end of the first week and patients at the end of the second week of illness using cut off values regarding percentage of cases with abnormal liver function tests.	106

LIST OF ABBREVIATIONS

ALP	Serum alkaline phosphatase
ALT	Serum alanine amino transferase
ASOT	Antistreptolysin O titre
AST	Serum aspartate amino transferase
BMAC	Bone marrow aspiration culture
°C	Celsius degree
CIE	Counter immuno electrophoresis
cm(s)	Centimeter(s)
CMI	Cell mediated immunity
CMIR	Cell mediated immune response
ELISA	Enzyme linked immuno-sorbent assay
ESR	Erythrocyte sedimentation rate
gm	Gram
H antigen	Flagellar antigen
Hb	Haemoglobin
IU	International unit
KAU	King Armestrong unit
Kg(s)	Kilogramm(s)
L(s)	Liter(s)
M	Mean
mg(s)	Milligram(s)
O antigen	Somatic antigen
r	Correlation coefficient
RBCs	Red blood cells
S. typhi	Salmonella typhi
SD	Standard deviation
SMZ	Sulfamethoxazole
T.F.	Typhoid fever
TLC	Total leucocytic count
TMP	Trimethoprim
Vi	Surface antigen

