BIOCHEMICAL ABERRATION INDUCED BY CERTAIN INSECT GROWTH REGULATORS (IGRs) IN HOUSE FLY, MUSCA DOMESTICA (MUSCIDAE - DIPTERA)



THESIS

SUBMITTED BY SAADYA MOHAMED ABD EL - NABI EL - BERMAWY

(Faculty of Education , Ain Shams Univ .)

For the Award of the Degree of

Doctor of Philosophy

In Entomology

593.77 5. M

from

Faculty of Science
Ain Shams University



1994.

THESIS	EXAMINATION	COMMITTEE
NAME	TITLE	SIGNATURE

SUPERVISORS:

PROF. DR. NAIMA A. ABD EL-RAZIK

ASSIS. PROF. DR. ABDEL-MOTY. H. M. HASSANEIN

DR. NABIL ABDUL KADER SALEH

ALL ()

HEAD OF DEPARTMENT

PROF . DR . BAHIRA M . EL- SAWAF



DEDICATION

I have a great honour to dedicate this dissertation and the including research to the spirit of Professor Dr. Awni Mohamed Guneidy. who learned me much in both moral and scientific aspects.

Saadya Mohamed El - Bermawy.

ACKNOWLEDGEMENT

The author is indebted with sincerest gratitude to prof Dr. Naima A. Abd El-Razik, Professor of Entomology, Entomology Department, Faculty of Science, Ain shams University for her encouragement and continuous advice throughout the study and for reading and correcting the manuscript.

The author wishes to express her sincere thanks and gratitude to Dr: Abd El-Moty H. M. Hassanein, Assistant professor of Entomology, Faculty of Education, Ain Shams University for his direct and greed supervision, valuable criticism and advice that made this study possible.

The author feel deeply grateful and indebted to Dr. Nabil Abdul Kader Saleh, Lecturer of Biochemistry, Department of Biochemistry, Ain Shams University. To him, I owe more than can be expressed for his generous help, constructive criticism, practious advice and valuable suggestions, stimulating and enlighting discussion during the experimental work and help in preparing the text.

Thanks are also due to Dr. Reda. F. Bakr, Assistant professor of Entomology, Department of Entomology, Ain Shams University for his consistent proposals, kind encouragement, and interest in progress of the study.

Grateful appreciation is also due to professor Dr. Fahmy.T. Ali, professor of Biochemistry, Department of Biochemistry, Ain shams University, for his valuable advices during the program of the study.

The author is also grateful to Dr. Ahmed. F. Afifi., Head of Biological Science and Geology Department, Faculty of Education, Ain Shams University for his encouragement and facilities offered during the present study.

Thanks are also due to all staff members and colleagues of both the Entomology, Biochemistry, Department, Faculty of Science and Biological Sciences and Geology Department, Faculty of Education, Ain Shams University for their ssincere help.

ABSTRACT

The present study investigated the possible toxicological and biochemical effects of three selected insect growth regulators (IKI 7899, BAY SIR 8514 and Sumilarv 71639) on different treated stages of the house fly M. domestica, using different techniques of application.

Most treatments led to various morphological aberrations which mainly observed in the pupal and adult stages . The pattern of protein contents and behavior of enzymatic activities of LDH, α - GPDH and AcP were also explored in all developmental stages after treatments .

The IKI was found to be the most effective inhibitor for all parameters studied regardless the technique applied followed by BAY SIR which shown equipotent.

Key words: Musca domestica - Insect growth regulators - Morphological aberrations - Proteins - AcP - LDH and α - GPDH.

Abbreviations

AcP Acid Phospatase

AkP Alkaline Phosphatase

BSA Bovine serum albumin

EMS ethyl methanesulfonate

lpha- GPDH , lpha - Glycerophosphate dehydrogenase

20- HE 20 - Hydroxy ecdysone

IGR Insect growth regulator

JH Juvenile hormone

JHA Juvenile hormone analogue

 JHa_{1} (Isopropyl - 3 , 7, 11 - trimethyl - 2,4 - dodeca dienoate) = ZR515

JHE Juvenile hormone esterase

JHM Juvenile hormone mimic

LDH lactate dehydrogenase

MDH Malate dehydrogenase

ME Malic enzyme

NAD⁺ *Nicotinamide adenine dinucleotide*

PNP Para - nitrophenol

Precocene I (7 - methoxy - 2 -2- dimethyl chromene)

PTU Phenylthiourea

RSA Relative specific activity

R-20458 [(1-4-ethylphenoxy)-6.7-epoxy-3.7-dimethyl-2-octene] = stauffer

SDH Succinate dehydrogenase

SJH Synthetic Juvenile hormone

List of Tables Included

		Page
Table (1)	Biological activities of IKI 7899 against one - day old larvae	
	of the house fly M. domestica fed at different times	(42)
Table (2)	Biological activities of BAY 8514 against one - day old larvae	
	of house fly M. domestica fed at different times	(43)
Table (3)	Biological activities of Sumilarv against one - day old larvae	
	of the house fly M. domestica fed at different times	(44)
Table (4)	Biological activities of IKI 7899 against M. domestica larvae	
	using dipping technique (50 and 100 seconds exposure)	(49)
Table (5)	Biological activities of BAY SIR 8514 against M. domestica	
	larvae, using dipping technique (50 and 100 seconds	
	exposure)	(50)
Table (6)	Biological activities of Sumilarv against M. domestica	
	larvae using dipping technique (50 and 100 seconds.	
	exposure)	(51)
Table (7)	Biological activities of IKI 7899 against M. domestica pupae	
	(dipping technique - 100 and 200 seconds exposure)	(57)
Table (8)	Biological activities of BAY SIR 8514 against M. domestica	
	pupae (dipping technique - 100 and 200 seconds exposure)	(58)
Table (9)	Biological activities of Sumilarv against M. domestica pupae	
	(dipping technique - 100 & 200 seconds exposure)	(59)
Table (10)	Biological activities of IKI, BAY SIR and Sumilarv against	
	M. domestica. Adult feeded for 24 hours	(62)

Table (HA)	Changes in total protein content of M. domestica larvae			
	after continuous feeding on treated media with different			
	concentrations of IGRs	(65)
Table (11B)	Changes in fraction protein content of M. domestica larvae			
	after continuous feeding on treated media with different			
	concentrations of IGRs	((56)
Table (12A)	Changes in total protein content of M. domestica adult after			
	feeding on treated milk with different concentrations of IGRs			
	for 24 hours	('	71)
Table (12B)	Changes in fraction protein content of M. domestica adult			
	after feeding on treated milk with different concentrations of			
	IGRs for 24 hours	(72)
Table (13A)	Changes in total protein content of M. domestica larvae after			
	dipping in different concentrations of IGRs for 50 seconds	(78)
Table (13B)	Changes in fraction protein content of M. domestica larvae			
	after dipping in different concentrations of IGRs for 50			
	seconds	(79)
Table (14A)	Changes in total protein content of M. domestica pupae after			
	dipping in different concentrations of IGRs for 100 seconds	(84)
Table (14B)	Changes in fraction protein content of M. domestica pupae			
	after dipping in different concentrations of IGRs for 100			
	seconds	(85)
Table (15A)	Changes in total AcP of M. domestica larvae after continuous			
	feeding on treated media with different concentrations of			
	IGRs	(89)

Table (15B)	Changes in fraction AcP of M. domestica larvae after			
	continuous feeding on treated media with different			
	concentrations of IGRs	(90)
Table (16A)	Changes in total AcP of M. domestica adult after feeding on			
	treated milk with different concentrations of IGR's for 24			
	hours	(95)
Table (16B)	Changes in fraction AcP of M. domestica adult after feeding			
	on treated milk with different concentrations of IGRs for 24			
	hours	(96)
Table (17A)	Changes in total AcP of M. domestica larvae after dipping in			
	different concentrations of IGR s for 50 seconds	(101)
Table (17B)	Changes in fraction AcP of M. domestica larvae after dipping			
	in different concentrations of IGR's for 50 seconds	(102)
Table (18A)	Changes in total AcP of M. domestica pupae after dipping in			
	different concentrations of IGRs for 100 seconds	(107)
Table (18B)	Changes in fraction AcP of M. domestica pupae after			
	dipping in different concentrations of IGRs for 100 seconds	(108)
Table (19)	Relative LDH activity during the ontogenesis of M .			
	domestica after continuous feeding larvae on treated media			
	by different concentrations of IGRs	(113)
Table (20)	Relative LDH activity during the ontogenesis of M .			
	domestica after feeding adult on treated milk by different			
	concentrations of IGRs for 24 hours	(116)
Table (21)	Relative LDH activity during the ontogenesis of M .			
	domestica after dipping larvae in different concentrations			
	IGRs for 50 seconds	(118)

Table (22)	Relative LDH activity during the ontogenesis of M .			
	domestica after dipping pupae in different concentrations of			
	IGRs for 100 seconds	(12	21)	
Table (23)	Relative α -GPDH activity during the ontogenesis of M .			
	domestica after continuous feeding larvae on treated media			
	with different concentrations of IGRs	(12	24)	
Table (24)	Relative α -GPDH activity during the ontogenesis of M .			
	domestica after feeding adults on treated milk by different			
	concentration of IGRs for 24 hours	(12	27)	
Table (25)	Relative α -GPDH activity during the ontogenesis of M .			
	domestica after dipping larvae in different concentrations of			
	IGRs for(50 sec .)	(1.	30)	į
Table (26)	Relative α -GPDH activity during the ontogenesis of M .			
	domestica after dipping pupae in different concentrations of			
	IGRs for (100 sec .)	(1	33))
Table (27)	Pattern of IKI 7899 effects after feeding and dipping of			
	M. domestica larvae	(1	35))
Table (28)	Pattern of IKI 7899 effects on different parameters after			
	Feeding adults and Dipping pupae of M. domestica	(1	3 6))
Table (29)	Pattern of BAY BAY 8514 effects feeding and dipping of			
	M. domestica larvae	(1	37)
Table (30)	Pattern of BAY SIR 8514 effects on different parameters			
	after Feeding adults and Dipping pupae of M. domestica	(1.	38)
Table (31)	Pattern of Sumilarv effects after feeding and dipping of			
	M damestica larvae	(1	30	١

Table (32)	Pattern of Sumilary effects on different parameters after			
	feeding adults and dipping pupae of M. domestica	(140)
Table (33)	Reliability of the application of the tested IGRs using			
	feeding and dipping techniques	(141)

List of Figures Included

		P	age	<u> </u>
Fig (1)	Biological activities of <i>IKI 7899</i> for one - day old larvae of M .			
	domestica (different times feeding)	(45)
Fig (2)	Biological activities of BAY SIR 8514 for one - day old larvae			
	of M. domestica (different times feeding)	(46)
Fig (3)	Biological activities of Sumilarv for one - day old larvae of			
	M. domestica (different times feeding)	(47)
Fig (4 A,B,C)	Biological activities of IGRs against M. domestica Larvae			
	using dipping technique (50& 100 Sec .)	(52)
Fig (5)	Morphogenic aberrations induced in the house fly larvae after			
	feeding larvae on treated media with IGRs	(53)
Fig (6)	Morphogenic aberrations induced in the house fly pupae after			
	feeding larvae on treated media with IGRs	(54)
Fig (7)	Various degrees of adult house fly eclosion induced by IGRs			
	after feeding larvae on treated media	(55)
Fig(8 A,B,C)	Effect of dipping pupae in different cancentrations of IGRs for			
	(100 & 200 Sec.) on hatching of egg laid	(60)
Fig (9)	Effect of feeding M. domestica adult on treated milk with			
	IGRs for 24 hours on hatching of egg laid	(63	;
Fig (10)	Effect of IKI 7899 on total proteins (A) and their fractions (B)			
	after continuous feeding of M. domestica larvae	(67	')
Fig(11)	Effect of BAY SIR 8514 on total proteins (A) and their			
	fractions (B) after continuous feeding of M, domestica larvae-	1	(68	3)
Fig (12)	Effect of Sumilary on total proteins (A) and their fractions			
	(B) after continuous feeding of M. domestica, larvae	1	(69	9)

		Pag	e
Fig (13)	Effect of IK1 7899 on total proteins (A) and their fractions (B)		
	after feeding of M. domestica adults for 24 hours	(73)
Fig (14)	Effect of BAY SIR 8514 on total proteins (A) and their		
	fractions (B) after feeding of M . $domestica$ adults for 24		
	hours	(74)
Fig (15)	Effect of Sumilary on total proteins (A) and their fractions (B)		
	after feeding of M. domestica adults for 24 hours	(75	;)
Fig (16)	Effect of IKI 7899 on total proteins (A) and their fractions (B)		
	after dipping of M. domestica larvae for (50 Sec.)	(80)
Fig (17)	Effect of BAY SIR 8514 on total proteins (A) and their		
	fractions (B) after dipping of M. domestica larvae for (50Sec)	(81)
Fig (18)	Effect of Sumilary on total proteins (A) and their fractions (B)		
	after dipping of M. domestica larvae for (50 Sec.)	(8	2)
Fig (19)	Effect of 1K1 7899 on total proteins (A) and their fractions		
	(B)after dipping of M. domestica pupae for (100 sec.)	(8	6)
Fig (20)	Effect of BAY SIR 8514 on total proteins (A) and their		
	fractions (B)after dipping of M. domestica pupae for (100		
	sec.)	(8'	7)
Fig (21)	Effect of Sumilary on total proteins (A) and their fractions		
	(B)after dipping of M. domestica pupae for (100 sec.)	(88	;)
Fig (22)	Percentage increase or decrease in Total AcP activity (A) and		
	their fractions (B) after continuous feeding of M. domestica		
	larvae on treated media with IKI 7899	(9	1)
Fig (23)	Percentage increase or decrease in Total AcP activity (A) and		
	their fractions (B) after continuous feeding of M. domestica		
	larvae on treated media with BAY SIR 8514.	(9	2)

		Page
Fig (24)	Percentage increase or decrease in Total AcP activity (A) and	
	their fractions (B) after continuous feeding of M. domestica	
	Iarvae on treated media with Sumilary	(93)
Fig(25)	Percentage increase or decrease in Total AcP activity (A) and	
	their fractions (B) after feeding of M. domestica adults on	
	treated milk by IKI 7899 for for 24 hours	(97)
Fig(26)	Percentage increase or decrease in Total AcP activity (A) and	
	their fractions (B) after feeding of M. domestica adults on	
	treated milk by BAY SIR 8514 for 24 hours	(98)
Fig(27)	Percentage increase or decrease in Total AcP activity (A) and	
	their fractions (B) after feeding of M. domestica adults on	I
	treated milk by Sumilarv for for 24 hours	(99)
Fig(28)	Percentage increase or decrease in Total AcP activity (A) and	
	their fractions (B) after dipping larvae of M. domestica in IKI	
	7899 for (50 Sec.)	(103)
Fig(29)	Percentage increase or decrease in Total AcP activity (A) and	
	their free (B) after dipping larvae of M. domestica in BAY	
	SIR 8514 for (50 Sec.)	(104)
Fig(30)	Percentage increase or decrease in Total AcP activity (A) and	
	their fractions (B) after dipping larvae of M. domestica in	
	Sumilary for (50 Sec.)	(105)
Fig(31)	Percentage increase or decrease in Total AcP activity (A) and	
	their fractions (B) after dipping pupae of M. domestica in IKI	
	7800 for (100 Sec.)	(100)