

**Evaluation of combination of certain
molluscicides against *Biomphalaria
alexandrina* and the free living
stages of *Schistosoma mansoni***

A Thesis submitted for
**THE DEGREE OF DOCTOR OF PHILOSOPHY
(IN ZOOLOGY)**

BY

FATHIA ALY MOHAMED GAWISH

B. Sc. Zoology & Chemistry Ain Shams University, 1977

M. Sc. Zoology El-Azhar University, 1985

Department of Zoology
Girls College for Arts, Science and Education
Ain-Shams University

1997



بِسْمِ اللَّهِ الرَّحْمَنِ الرَّحِيمِ

وَعَلَّمَكَ مَا لَمْ تَكُن تَعْلَمُ
وَكَانَ فَضْلُ اللَّهِ عَلَيْكَ عَظِيمًا

صَدَقَ اللَّهُ الْعَظِيمُ

سورة النساء - آية ١١٣

SUPERVISORS

Prof. Dr. ERIAN GEORGE KAMEL
Professor of Zoology and Head of Zoology
Department, Girls College for Arts,
Science and Education
Ain-Shams University

Prof. Dr. MOHAMED A. EL-EMAM
Professor of Medical Malacology,
Theodor Bilharz Research Institute
Embaba, Egypt.

Dr. M.A.El-Emam
17-2/97

❀ ACKNOWLEDGMENT ❀

❀ The author wishes to express her deep gratitude to **PROF. DR. ERIAN GEORGE KAMEL**, Professor of Zoology and Head of Zoology Department, Girls College for Arts, Science and Education Ain-Shams University, for supervising the work, reading the final manuscript and continuous encouragement.

❀ The author is greatly indebted to **PROF. DR. MOHAMED A. EL-EMAM** Professor and head of Medical Malacology laboratory, Theodor Bilharz Research Institute, for direct supervision, continuous guidance, valuable help in preparation the manuscript and encouragement.

❀ The author wishes to thank also **PROF. DR. MENRIET ZAKI ROUSHDY** Professor of Medical Malacology and Head of Environmental research and Medical Malacology Department, Theodor Bilharz Research Institute, for her help and kind encouragement.

CONTENTS

ACKNOWLEDGEMENT	Page
INTRODUCTION	1
REVIEW OF LITERATURE	3
MATERIALS AND METHODS	22
1- Breeding and maintenance of snails	22
2- Calculation of sublethal doses of molluscicides	23
3- Effect of sublethal concentrations of Bayluscide, copper sulphate and their mixtures on growth, survivorship and egg laying of <i>Biomphalaria alexandrina</i> snails and hatchability of their eggs in laboratory	25
3.1.The survival and growth of <i>Biomphalaria alexandrina</i> juvenile snails.	25
3.2.Survivorship and egg-laying capacity of adult <i>Biomphalaria alexandrina</i> snails.	26
3.3.Hatchability of <i>Biomphalaria alexandrina</i> eggs.	27
4- Effect of Bayluscide, copper sulphate and their mixtures on the free living stages of <i>Schistosoma mansoni</i> (miracidia & cercariae)	28
4.1.Miracidia	28
4.2.Cercariae	28
5- Effect of sublethal concentrations of Bayluscide, copper sulphate and their mixtures on the maturation of <i>Schistosoma mansoni</i> infection in <i>Biomphalaria alexandrina</i> snails and cercarial production.	29

	<i>Page</i>
5.1.Exposure of snails to molluscicides	29
5.2.Exposure of snails to miracidia	30
5.3.Examination of the snails for cercarial shedding	31
 6- Effect of sublethal concentrations of Bayluscide, copper sulphate and their mixtures on the infectivity of <i>Schistosoma mansoni</i> cercariae	 32
6.1.Experimental animal groups	32
6.2.Preparation and counting of cercarial suspension	32
6.3.Infection of mice	33
6.4.The Oogram method	34
6.5.Digestion of liver and intestine (ova count/gm tissue)	36
	37
RESULTS	
1- Susceptibility of adult <i>Biomphalaria alexandrina</i> to Bayluscide and copper sulphate	37
 2- Effect of sublethal concentrations of Bayluscide, copper sulphate and their mixtures on life span, survival and growth rates of <i>Biomphalaria</i> <i>alexandrina</i> juvenile snails	 40
2.A.Survival rate	40
2.B.Growth rate	44
2.C.Life span	49
 3- Effect of sublethal concentrations of Bayluscide, copper sulphate and their mixtures on life span, survival rate and fecundity of adult <i>Biomphalaria</i> <i>alexandrina</i> snails	 53
3.A. Survival rate	53
3.B.Life span	57
3.C.Snail fecundity	60
 4- Effect of Bayluscide, copper sulphate and their mixtures on hatchability of adult <i>Biomphalaria</i> <i>alexandrina</i> eggs	 64

	<i>Page</i>
5- Effect of Bayluscide, copper sulphate and their mixtures on the free living stages of <i>Schistosoma mansoni</i> (miracidia and cercariae)	70
5.A.Miracida	70
5.B.Cercariae	70
6- Effect of sublethal concentrations of Bayluscide, copper sulphate and their mixtures on infection of <i>Biomphalaria alexandrina</i> with <i>Schistosoma mansoni</i>	74
6.1.Survival rate of snails at first shedding	75
6.2.Infection rate of <i>Biomphalaria alexandrina</i> snails with <i>Schistosoma mansoni</i>	78
6.3.The prepatent period (cercarial incubation period).....	80
6.4.The life span of infected <i>Biomphalaria alexandrina</i> snails	81
6.5.Duration of cercarial shedding	84
6.6.Cercarial production	88
6.7.Pattern of <i>Schistosoma mansoni</i> cercarial production from infected <i>Biomphalaria alexandrina</i> snails post first shedding.	95
7- Effect of Bayluscide, copper sulphate and their mixtures on infectivity of <i>Schistosoma mansoni</i> cercariae	106
7.1.Worm load/mouse	106
7.2.. <i>Schistosoma mansoni</i> ova per gram tissue in liver and intestine of infected mice	111
7.3.Oogram (developmental stages of <i>Schistosoma mansoni</i> ova)	118
DISCUSSION	133
SUMMARY	157
REFERENCES	163
ARABIC SUMMARY	191

LIST OF TABLES

Table	Page
1- Comparative susceptibility of adult <i>Biomphalaria alexandrina</i> snails to Bayluscide and copper sulphate after 24 hours of exposure.	38
2- Survival rate (L_x) and mean shell diameter (mm) of juvenile <i>Biomphalaria alexandrina</i> snails continuously exposed to sublethal concentrations of Bayluscide, copper sulphate and their mixtures.	41
3- Growth rate (%) of <i>Biomphalaria alexandrina</i> throughout their life span under continuous exposure to sublethal concentrations of Bayluscide, copper sulphate and their mixtures.	45
4- Life span of <i>Biomphalaria alexandrina</i> juvenile snails exposed continuously to sublethal concentrations of Bayluscide, copper sulphate and their mixtures.	51
5- Survivorship (L_x) and fecundity (M_x) of adult <i>Biomphalaria alexandrina</i> snails continuously exposed to sublethal concentrations of Bayluscide, copper sulphate and their mixtures.	54
6- Life span of adult <i>Biomphalaria alexandrina</i> snails continuously exposed to sublethal concentrations of Bayluscide, copper sulphate and their mixtures.	58
7- Effect of Bayluscide, copper sulphate and their mixtures on hatchability of <i>Biomphalaria alexandrina</i> eggs.	65
8- Effect of Bayluscide, copper sulphate and their mixtures on <i>Schistosoma mansoni</i> miracidia.	71
9- Susceptibility of <i>Schistosoma mansoni</i> cercariae to Bayluscide, copper sulphate and their mixtures.	73

<i>Table</i>	<i>Page</i>
10- Effect of sublethal concentrations of Bayluscide, copper sulphate and their mixtures on the survival rate, infection rate, prepatent period and life span of <i>Biomphalaria alexandrina</i> exposed to <i>Schistosoma mansoni</i> miracidia.	76
11- Effect of sublethal concentrations of Bayluscide, copper sulphate and their mixtures periodic cercarial production and duration of shedding (days).	85
12- Periodic cercarial production from <i>Biomphalaria alexandrina</i> snails exposed to <i>Schistosoma mansoni</i> miracidia (mean number of cercariae/snail/3days).	96
13- Worm load of <i>Schistosoma mansoni</i> in albino mice infected with cercariae exposed to sublethal concentrations of Bayluscide, copper sulphate and their mixtures during mice infection.	107
14- Worm load of <i>Schistosoma mansoni</i> in albino mice infected with cercariae exposed to sublethal concentrations of Bayluscide, copper sulphate and their mixtures one hour pre-mice infection.	108
15- Worm load of <i>Schistosoma mansoni</i> in albino mice infected with cercariae exposed to sublethal concentrations of Bayluscide, copper sulphate and their mixtures 3 hours pre-mice infection.	109
16- Number of ova/gram tissue of albino mice infected with <i>Schistosoma mansoni</i> cercariae exposed to Bayluscide, copper sulphate and their mixtures during mice infection.	112
17- Number of ova/gram tissue of albino mice infected with <i>Schistosoma mansoni</i> cercariae exposed to Bayluscide, copper sulphate and their mixtures 1 hour pre-mice infection.	113