THE USE OF PERFLUOROCARBON LIQUIDS IN THE MANAGEMENT OF GIANT RETINAL BREAKS

ESSAY

Submitted for Partial Fulfilment of the Degree of M.Sc. in Ophthalmology

By

Mohamed Abdul-Rahman Mohamed Awadallah

M.B., B.CH.

525°8

Supervised By

Prof. Dr. Hussein Shaker EL-Markabi

Prof. of Ophthalmology Ain-Shams University

61773

Prof. Dr. Ahmed Abul-Naga

Prof. of Ophthalmology Ain-Shams University M.A

Dr. Khaleed AL-Tagory Lecturer of Ophthalmology

Ain-Shams University



Department of Ophthalmology Faculty of Medicine Ain-Shams University

1996







Ain-Shams University Faculty of Medicine Department of Ophthalmology

Name: Mohamed Abdul-Rahman Mohamed Awadallah

Title: THE USE OF PERFLUOROCARBON LIQUIDS IN THE MANAGEMENT OF GIANT RETINAL BREAKS

SUPERVISORS

APPROVED

Prof. Dr. Hussein Shaker EL-Markabi Prof. of Ophthalmology Ain-Shams University

Prof. Dr. Ahmed Abul-Naga Prof. of Ophthalmology Ain-Shams University

Dr. Khaleed AL-Tagory
Lecturer of Ophthalmology
Ain-Shams University

Head of Ophthalmology Department



ACKNOWLEDGMENT

First of all, cordial thanks to "ALLAH" who enabled me to overcome all problems that faced me throughout the work.

I would like to express my outmost gratitude and indebtedness to **Prof.Dr. Hussein Shaker EL-Markabi**, Prof. of Ophthalmology, Faculty of Medicine, Ain-Shams University, for suggesting the topic, their constructive guidance during the lay out of the work and illuminating criticism in reading the manuscript. and **Prof. Dr. Ahmed Abul-Naga**, Prof. of Ophthalmology, Faculty of Medicine, Ain-Shams University, for constructive guidance, illuminating criticism in reading the manuscript and advice.

My grateful thanks are due to **Dr. Khaleed AL-Tagory**, Lecturer of Ophthalmology, Faculty of medicine, Ain-Shams University, for his advice and encouragement.

Finally, my sincere thanks are due to the staff members of Opthalmology Department, Faculty of Medicine, Ain Shams University, for their help and cooperation.



CONTENTS

- Introduction and aim of the work	1
CHAPTER I	
- Anatomy	3
I - Anatomy of the retina	3
11-Anatomy of the vitreous	13
CHAPTER II	
- Definition of Giant Retinal Tear	17
CHAPTER III	
- Old Methods in the Management of Giant	
Retinal Tears	23
CHAPTER IV	
- Physical and Chemical Properties of	
Perfluorocarbon Liquids	33
CHAPTER V	
- Surgical Techniques	39
CHAPTER VI	
- Complications of Perfluorocarbon Liquids	67
CHAPTER VII	
- Advantages of Perfluorocarbon Liquids	80
- Summary and Conclusion	83
- References	89
- Arabic Summary.	



LIST OF TABLES

Table 1: Chemical formula and physical properties of perfluorocarbon liquids used in vitreo retinal	
surgery.	33
Table 2: Force of tamponade exerted by various substitut	es.
LIST OF FIGURES	34
LIST OF FIGURES	
Fig.1: Normal fundus.	10
Fig.2: Giant retinal tear.	17
Fig.3: Superior giant retinal tear.	22
Fig.4 :Preoperative ophthalmoscopic examination of pat	ient
with upper giant tear is ideally performed with	i
patient on tilting table.	25
Fig.5: Preoperative examination of superior giant retinal	tear
with an inverted retinal flap.	25

Fig.7: The basal vitreous gel is removed using deep scle	eral
depression.	43
Fig.8: A 20-gauge microsurgical instrument.	47
Fig.9: Injection of perfluorocarbon liquid 1mm above	the
optic disc.	49
Fig.10: Dissection within the vitreous base.	49
Fig.11: The perfluorocarbon liquid injected over the e	dge
of the retinal tear.	49
Fig.12: The flute needle with silicone tip is placed near	the
edge of the tear to aspirate saline and subretinal f	luid
completely before removing the perfluorocarbon	L
liquid.	52
Fig.13: A slow and deliberate fluid-air exchange is don	ie.

40

52

Fig.6: Circumfrential scleral buckle.

Fig.14:Endophotocoagulation can be applied	through the
perfluorocarbon liquid to all retina.	56

- Fig.15: Schematic drawing showing a procedure for complete removal of perfluorocarbon liquid.
- Fig.16: Removal of perfluorocarbon liquid by passive aspiration using flute needle.
- Fig. 17: Slipping of small perfluorocarbon bubbles under the retina.
- Fig.18: Fragmentation of the perfluorocarbon bubble by a forceful stream of balanced salt solution which disturbe the anterior eniscus of the perfluorocarbon bubble.
- Fig. 19: Fragmentation of the perfluorocarbon bubble caused by direct contact between the infusion cannula and the anterior meniscus of the perfluorocarbon bubble.

72

