POLYMORPHONUCLEAR ELASTASE

IN NEONATAL SEPSIS IN EGYPTIAN NEONATES

M.S.Thesis

Submitted for partial fulfillment of

requirements of Master Degree in Pediatrics

Bv

Essam Mahmoud Ebrahim (M.B.B.Ch)

Under Supervision of

Prof. Dr. Mohamed Mohamed Fa

Professor of Pediatrics Faculty of Medicine

Ain Shams University

Dr. Hesham Abd El-Samei Awad

Lecturer of Pediatrics

Faculty of Medicine Ain Shams University

Novema & Baden. Dr. Nashwa Ahmed Adel El-Badawi

Assistant Prof. of Clinical Pathology Faculty of Medicine

> Faculty of Medicine Ain Shams University

> Ain Shams University

1996



Acknowledgment

I would like to express my sincere gratitude to Prof. Dr. Mohamed Farid, Professor of Pediatrics, Ain Shams University for his meticulous supervision of this work.

I deeply thank Dr. Hesham Abd El-Samei Awad, Lecturer of Pediatrics, Ain Shams University for his patience and constructive criticism during the preparation of this thesis.

I wish to thank Dr. Nashwa Ahmed Adel El-Badawi, Assistant Prof. of Clinical Pathology, Ain Shams University for her valuable advice and keen supervision.

I also thank all the nurses and workers in the Neonatal Care Unit, Childrens' Hospital, Ain Shams University, for helping me in preparing this work.

Last but not least, I would like to thank my family for helping me to complete this work.



List of abbreviations

- α-1-AT: Alpha -1- antitrypsin.
- α-2-MG: Alpha -2- macroglobulin.
- α-1-Pl: Alpha -1- proteinase inhibitor.
- · Ante-part. hge: Ante-partum hemorrhage.
- APC: Antigen producing cells.
- ASD: Atrial septal defect.
- ATIII: Antithrombin III.
- B-N-Menigitidis: Group B- Neisseria meningitidis.
- C3: Complement 3.
- CIE: Countercurrent immunoelectrophoresis.
- CNS: Central nervous system.
- CRP: C reactive protein.
- · CSF: Cerebro-spinal fluid.
- · 1D: One day.
- · DIC: Disseminated intravascular coagulation .
- · DM: Diabetes mellitus.
- E.Coli.: Escherichia Coli.
- EDTA: Ethylene diamine tetra-acetic acid.
- ELP: Elastase like proteinase.
- · ESR: Erythrocyte sedimentation rate.
- ETI: Endo-tracheal intubation.
- ETT: Endo-tracheal tube.
- · F: Female.
- Fc: Crystallized fragment.
- · FSPs: Fibrin split products.
- · GBS: Group B-streptococci.
- Hb%: Haemoglobin percent.
- · HDN: Haemolytic diseases of the newborn.
- H-influenza: Hemophilus influenza.
- HIE: Hypoxic ischemic encephalopathy.

- · HNE: Human neutrophil elastase.
- · HSM: Hepato-splenomegaly.
- I.D.: In diameter.
- · IDM: Infant of diabetic mother.
- IG: Immunoglobulin.
- IMAC: Immunoactivation.
- IMV: Intermittent mandantory ventilation.
 I/T: Immature / total.
- IUGR: Intra-uterine growth retardation.
- KD: Kilo Dalton.
- ITI: Inter-α-trypsin inhibitor.
- I.V.: Intravenous.
- IVI: Intravenous immunoglobulin.
- LTB₄: Leukotreine B₄.
- · M: Male.
- · MAS: Meconium aspiration syndrome.
- MPI: Mucus proteinase inhibitor.
- · MPO: Myelo-peroxidase.
- MW: Molecular weight.
- NADPH oxidase: Nicotinamide adenosine dinucleotide phosphate oxidase.
- NEC: Necrotizing enterocolitis.
- · NICU: Neonatal intensive care unit.
- · NK: Natural killer.
- · NPV: Negative predictive value.
- NSP: Neutrophil storage pool.
- PA: Perinatal asphyxia.
- · PAF: Platelet activation factor.
- PC: Personal computer.
- · PDA: Patent ductus arteriosus.
- PEEP: Positive end expiratory pressure.
- PMNs: Polymorphonuclear leukocytes.

- PPV: Positive predictive value.
- PROM: Premature rupture of membranes.
- PT: Preterm.
- PT: Prothrombin time.
- PTT: Partial thromboplastin time.
- RBCs: Red blood cells.
- RD: Respiratory distress.
- SD: Standard deviation.
- TGVs: Transposition of great vessels.
- TLC: Total leukocytic count.
- TPN: Total parenteral nutrition.
- UTI: Urinary tract infection.
- VLBW: Very low birth weight.
- · WBCs: White blood cells.
- ZSR: Zeta sedimentation rate.

List of Tables

* Review of literature:

l	Defense mechanisms against infectious pathogens	4
2	Characteristics of neutrophil in newborn compared	
	with adult.	14
3	Oxidative metabolic response of PMNs.	15
4	The microbiology of neonatal sepsis between 1979	
	and 1988 at Yale New Haven Hospital.	18
5	The predominant causative organisms of neonatal	
	sepsis.	19
6	Distribution by age of 270 positive cultures at Yale	
	New Haven Hospital (1979 to 1988)	20
7	Clinical signs and symptoms of sepsis in neonate	26
8	The positive and negative predictive value of commonly 39	
	ordered laboratory tests.	36
9	The reference value for neutrophilic cells in healthy neonates	40
10	Deviation of hematologic scores according to hematologic scoring	
	system.	40
11	Sepsis score	47
12	The doses and frequency of administration of the various	
	chemotherapeutic agents used in treatment of neonatal sepsis	51
13	The commonly used inotropes, vasoconstrictors, and vasodilators used	
	in neonatal shock	52
14	Natural elastase inhibitors of human origin.	70

* RESULTS:

15	Comparison between the cases and control groups	
	considering the gestational age.	78
16	The sex distribution among our population	78
17	The birth weight of the cases and control groups.	79
18	The level of elastase measured in cases and control groups	80
19	The level of elastase in cases (survivors and non-survivors)	81
20	Comparison between cases with RD and cases without RD as regards	
	the level of elastas	82
21	Comparison between cases with perinatal asphyxia as regards the level	
	of elastase.	82
22	Comparison between cases with M.A.S. and those	
	without M.A.S. as regards the level of elastase.	83