Incidence of Injury of Recurrent Laryngeal Nerve During Thyroid Surgery

Thesis

Submitted for Partial Fulfilment of Requirements of Master Degree in General Surgery

The second secon

by

Mohamed Mohamed Tawfek M.B., B.Ch.

Supervised by

63425

Prof. Dr. Adel Abdel Kader

Professor of General Surgery, Faculty of Medicine, Ain Shams University

Dr. Gamal Abdel Rahman El Mowalled

Lecturer of General Surgery, Faculty of Medicine, Ain Shams University

> Faculty of Medicine Ain Shams University 1998



السالخ المرع

"قالوا سبحانك لا علم لنا إلا ما علمتنا، إنك أنت العليم الحكيم" بيرة -٢٢

Contents

Contents

	Page
Acknowledgment	iv
List of Figures	v
1- Introduction	1
2- Review of literature	4
Anatomy and Embryology of the Thyroid Gland	4
Anatomy of R.L.N. and Vocal Cords	19
• Physiology of R.L.N.	32
 Pathological Classification of R.L.N. Lesions 	43
Indications of Thyroidectomy	53
• Technique of Thyroidectomy	59
Complications of Thyroidectomy	77
 Management of R.L.N. Injury 	89
3- Patients and methods	104
4- Results	107
5- Discussion.	112

Incidence of Injury of R. L. N. During Thyroid Surgery

6- Conclusion	118
7- Summary	119
9- References	121
10- Arabic summary	





Acknowledgment

First thanks are all due to Allah for blessing work until it has reached its end, as part of his generous help throughout my life.

I would like to direct special thanks to Prof. Dr. Adel Abdel Kader, Professor of General Surgery, Faculty of Medicine, Ain Shams University, for the great support and encouragement he gave me throughout the whole work. It is a great honour to work under his guidance and supervision.

I am greatly honoured to express my utmost thanks to Dr. Gamal Abdel Rahman El Mowalled, Lecturer of General Surgery, Faculty of Medicine, Ain Shams University, from whom I received faithful supervision, valuable sugestions, and continuous guidance throughout this work.

Mohamed Mohamed Tawfek



List of Figures

Figure	Page
Figure (1): Thyroid gland, anterior view.	6
Figure(2): Thyroid gland, posterior view.	7
Figure (3): Arterial supply to the thyroid gland, The thyroid ima artery is only occasionally present.	9
Figure (4): Venous drainage of the thyroid gland.	11
Figure (5): The lymphatic drainage of the thyroid gland.	14
Figure (6): Commonly occurring anatomy of the recurrent laryngeal nerve which is ascending anterior to the tracheo-oesophageal groove, passing deep to the main branches of the inferior thyroid artery and dividing into two branches before passing deep to the ligament of Berry and entering the larynx beneath the inferior comu of the thyroid cartilage.	21
Figure (7): A. Recurrent laryngeal nerve running between the branches of the inferior thyroid artery. B. Recurrent laryngeal nerve anterior to inferior thyroid artery. C. Recurrent laryngeal nerve posterior to inferior thyroid artery.	22
Figure (8): The larynx.	23
Figure (9): The relation between the superior laryngeal nerve and the thyroid gland.	24
Figure (10): The relation between the recurrent larryngeal nerve and inferior thyroid artery.	25
Figure (11): A. Non-descended inferior or recurrent laryngeal nerve. B. Partially descended inferior or recurrent laryngeal nerve.	26
Figure (12).: A series of diagrams to show different positions of vocal cords and arytenoid cartilages.	29
A: Position of rest in quiet respiration. The intermembranous part of the rima dlottides is triangular and the intercartilagenous part is rectangular in shape	

- B: Forced inspiration. Both parts of rima glottides are triangular in shape.
- C: Abduction of the vocal cords. The arrows indicate the lines of pull of the posterior crico-arytenoid muscles. The abducted vocal cords and the abducted, retracted and internally rotated arytenoid cartilages are shown in dotted outline. The entire rima glottides is triangular.
- D: Adduction of the vocal folds. The arrows indicate the line of pull of the lateral crico-arytenoid muscles. The adducted vocal folds and the medially rotated arytenoid cartilages are shown in dotted outlines.
- E: Closure of the rima glottides. The arrows indicate the line of pull of the transverse arytenoid muscle. Both the arytenoids and the vocal cords are adducted, but there is no rotation of the former.
- F: Tension of the vocal folds, produced by the action of the crico-thyroid muscle which tilt the anterior part of the cricoid cartilage cranially and so carry the arytenoid cartilages dorsally.
- G: Relaxation of the vocal folds, produced by the action of thyro-arytenoid muscles, which draw the arytenoid cartilage ventrally.
- Figure (13): Skin incision. Division of cervical fascia in the mid-line after elevation of the flap.
- Figure (14): Division of the suspensory ligament. 63
- Figure (15): Opening of the cricothyroid space to expose the superior pole vessels to protect the external laryngeal nerve.
- Figure (16): Clamping of superior thyroid vessels with protection of external laryngeal nerve.
- Figure (17): Individual division of branches of the anterior thyroid artery.
- Figure (18): Subtotal thyroidectomy, transection of the capsule at the line of inferior thyroid artery branches.
- Figure (19): Subtotal thyroidectomy. 1gram of thyroid tissue is left bilaterally.

71

Figure (20): Diagrams illustrating six of the eight significant nerve- artery relationships. 72

Figure (21): a) Simultaneous stimulation of the recurrent nerve and palpation of impulse in the cricothyroid area are illustrated. b) Left lateral, course of recurrent laryngeal nerve as it ascends alongside tracheo-oesophageat groove. Right lateral, recurrent laryngeal nerve as it ascends obliquely toward larynx after winding around subclavian artery. Note prominent inferior comu of thyroid cartilage and its relation to recurrent laryngeal nerve.

73

Figure (22): Location of recurrent laryngeal nerve in triangle.

94

