

MODULATION OF TOXICITY OF SOME PHENYLALKYLAMINES ILLICIT "STREET" NARCOTICS BY XANTHINOL NICOTINATE

Thesis

Presented in Partial Fulfillment of the
Requirement for Master Degree In Clinical Toxicology
Department of Forensic medicine and Clinical Toxicology

By

Iman Saad Kamal Ahmed

M.B.B. Ch. Ain Shams University 1992

Assistant Researcher in Clinical Toxicology
National Research Centre, Dokki, Cairo, Egypt.

614.19

I.S

Supervisors

Prof. Dr.

M. SAFWAT SOLIMAN

Prof. of Forensic medicine &
Toxicology, Faculty of Medicine
Ain Shams University

Prof. Dr.

ALY MOHAMED DIAB

Prof. of Analytical Toxicology
Narcotic & Toxicology Lab.
National Research Center

Prof. Dr.

SANAA IBRAHIM BILAL

Assistant Prof. of Forensic medicine
& Toxicology
Faculty of Medicine
Ain Shams University

Submitted to the
Faculty of Medicine
Ain Shams University
1996



ACKNOWLEDGEMENT

I am deeply thankful to **ALLAH** by the grace of whom the present work was accomplished.

I wish to express my gratitude to Prof. Dr. **Mohamed Safwat Soliman**, Professor of Forensic medicine & Toxicology, Faculty of Medicine, Ain Shams University, who gave me every help possible, full support, with his vast experience and fathery encouragement to accomplish this work.

I would like to pay a sincere tribute to Prof. Dr. **Aly Mohamed Diab**, Professor of Analytical Toxicology, National Research Centre, Dokki, Egypt, for suggesting the point and planning the work. Actually his beneficial directions and supervision and constructive criticism were indispensable to me.

I wish to express my thanks to Dr. **Sanaa Ibrahim Bilal**, Assistant Professor of Forensic medicine & toxicology, Faculty of medicine, Ain Shams University, who gave me every encouragement, supervision and help possible.

My deep thanks to Dr. **Sahar A. Aggour**, Lecturer of Pathology, Faculty of Medicine, Cairo University & Dr. **Wafaa A. Shata**, Lecturer of pathology, National Research Centre, Dokki, for their help in examination of pathological specimens.

My deep thanks to every person assisted this research to appear in its final form, with special regards to General. Essam El Tersawy, General. Mostafa Taher, Anti Drug Egyptian - General organization, Ministry of Interior, Mr. Wael Aman, Vice-General Prosecutor of

North Cairo prosecution, Abassia, for their kind supply of seizures and valuable data.

I extend my deep thanks and profound gratitude to the National Research Centre, Dokki, Egypt, who provided all possible facilities to complete this work.

I wish to express my deep thanks and profound gratitude to my colleagues in the department of Forensic medicine & clinical toxicology, Faculty of Medicine, Ain Shams University, who gave all possible help and encouragement.

Last but not least, all my sincere and deep gratitude are due to my mother Mrs Kawsar El Shafei, Head Mistress of El Nasr School, Ministry of Education, Whose encouragement and support allowed me to finish this thesis appropriately and efficiently.

CONTENTS

	Page
Part I :	
1. Introduction	1
2. Aim of Work	4
3. Review of Literature	
a) Khat	6
b) Methcathinone	17
c) Xanthinol nicotinate	27
Part II :	
4. Experimental	
Material, reagents and equipments	30
Methodology	35
Part III :	
5. Results	
- Part I	41
- Part II	43
- Negative & positive control groups	43
- Khat treated group	44
- Khat treated followed by xanthinol nicotinate treated group	45
6. Discussion	
- Part I	84
- Part II	86
7. General summary & Conclusions	89
Conclusions	92
Recommendations	93
8. References	94
9. Arabic Summary	

LIST OF FIGURES

Fig. No.		Page
Fig. (1)	Structural Formulae of Cathine and Cathinone.	11
Fig. (2)	Structural formulae of Cathinone biodegradation.	11
Fig. (3)	Thin layer chromatography of the chloroform fraction of Khat extract.	39
Fig. (4)	Body weight changes in control and Khat treated rats.	56
Fig. (5)	The mean ALT levels in control and Khat treated rats.	57
Fig. (6)	The mean AST levels in control and Khat treated rats.	58
Fig. (7)	Rat's liver control group H & E stain X 200.	59
Fig. (8)	Rat's liver 0.8 ml group H & E stain X 200. Central vein is congested, hepatocytes are markedly vacuolated especially towards the periphery.	60
Fig. (9)	Rat's liver 0.8 ml group H & E stain X 200. Mononuclear cell infiltration of the portal tract, with marked vacuolation of the adjacent hepatocytes, Von-Kupffer cells are prominent.	61
Fig. (10)	Rat's liver 0.8 ml group H & E stain X 200. Central vein is congested prominent Von-Kupffer cells with mononuclear cell infiltration among the hepatocytes.	62
Fig. (11)	Rat's liver 0.2 ml H & E stain, prominent Von-Kupffer cells and portal area is markedly infiltrated with mononuclear cells and multinucleated giant cells.	63
Fig. (12)	Fig. (5) magnified X 400 portal area is infiltrated with mononuclear cells and giant cells.	64
Fig. (13)	Rat's liver 0.2 ml group. H & E stain X 400. Necrotic area infiltrated with mononuclear cells, other hepatocytes show abnormal vacuolation.	65

Fig. No.		Page
Fig. (14)	Rat's liver control group. Masson Trichome stain X 200 Minimal collagen around the central vein & in the portal tract.	66
Fig. (15)	Rat's liver 0.2 ml group. Masson trichome stain X 200, Collagen is deposited around a dilated central vein & in the portal tract. A markedly dilated and congested blood sinusoid has caused pressure atrophy of the surrounding hepatocytes.	67
Fig. (16)	Rat's liver 0.2 ml group. Masson trichome stain X 200, increased collagen deposition around a central vein, in the portal tract & in between the hepatocytes.	68
Fig. (17)	Rat's liver, control group, PAS stain X 200, Normal glycogen content (PAS positive).	69
Fig. (18)	Rat's liver 0.2 ml group, PAS stain X 200, Marked decrease of glycogen content especially towards the periphery of hepatic lobules.	70
Fig. (19)	Rat's liver 0.2 ml group, PAS stain X 200, Marked glycogen depletion, the portal vein is markedly dilated & congested. An area of haemorrhage & necrosis is also seen.	71
Fig. (20)	Body weight changes in the four studied group.	72
Fig. (21)	The mean ALT levels in the four studied groups.	73
Fig. (22)	The mean AST levels in the four studied groups.	74
Fig. (23)	The mean level of total protein concentrations in the four studied groups.	75
Fig. (24)	The mean level of serum albumin concentration in the four studied groups.	76
Fig. (25)	Albumin/Globulin ratio among the four studied groups.	77

Fig. No.		Page
Fig. (26)	Rat's liver control group. Hematoxylin & eosin stain X 150. Preserved hepatic architecture, hepatocytes are arranged in thin plates within lobules.	78
Fig. (27)	Rat's liver control group. High power X 300 H & E stain, showing preserved architecture.	79
Fig. (28)	Rat's liver of positive control group, H & E X 150 the central vein is markedly dilated (improved blood supply to the liver) without any pressure effect upon neighbouring hepatocytes.	80
Fig. (29)	Rat's liver of positive control group, H & E X 300, showing unremarkable pathological changes.	81
Fig. (30)	Rat's liver of Khat treated followed by xanthinol nicotinate group, H & E X 150, showing hydropic degeneration of the hepatocytes and narrow sinusoids.	82
Fig. (31)	Rat's liver of Khat treated followed by xanthinol nicotinate group, H & E X 300 showing diffuse hydropic degeneration of the hepatocytes, signs of regeneration i.e., appearance of binuclear cells, double and several rows of liver plates could be observed.	83

LIST OF TABLES

Table No.		Page
Table (1)	Synthetic phenylalkylamines.	3
Table (2)	Statistical analysis of body weight changes in Khat treated rats (g.).	47
Table (3)	Statistical analysis of ALT activity (UI/L) after treatment of albino rats with different dose levels of Khat extract (0.2 ml, 0.4 ml & 0.8 ml).	47
Table (4)	One way analysis of variance (ANOVA) of ALT activity between different groups (0.2, 0.4 & 0.8 ml).	47
Table (5)	Statistical analysis of AST activity (UI/L) after treatment of albino rats with different dose levels of Khat extract (0.2, 0.4 & 0.8 ml).	48
Table (6)	One way analysis of variance (ANOVA) of AST activity between different groups (0.2, 0.4 & 0.8 ml).	48
Table (7)	Statistical analysis of body weight changes (g.) in different groups of rats (negative control, positive control, Khat treated & khat treated followed by xanthinol nicotinate.	49
Table (8)	Statistical analysis of ALT activity (UI/L) in different groups of rats / negative control, positive control, Khat treated & Khat treated followed by xanthinol nicotinate).	50
Table (9)	One way analysis of variance (ANOVA) of ALT activity between different groups [negative control (1), positive control (2), Khat treated (3) & Khat treated followed by xanthinol nicotinate (4)].	50

Table No.	Page
Table (10) Statistical analysis of AST activity in different groups of rats (negative control, positive control, Khat treated & Khat treated followed by xanthinol nicotinate).	51
Table (11) One way analysis of variance (ANOVA) of AST activity between different groups [negative control (1), positive control (2), Khat treated (3) & Khat treated followed by xanthinol nicotinate (4)].	51
Table (12) Statistical analysis of total protein concentrations (g/dl) in different groups of rats (negative control, positive control, Khat treated and Khat treated followed by xanthinol nicotinate).	52
Table (13) One way analysis of variance (ANOVA) of total protein concentrations between different groups (negative control (1), positive control (2), Khat treated (3) & Khat treated followed by xanthinol nicotinate).	52
Table (14) Statistical analysis of albumin concentrations (g/dl) in different groups of rats (negative control, positive control, Khat treated & Khat treated followed by xanthinol nicotinate).	53
Table (15) One way analysis of variance (ANOVA) of albumin concentrations between different groups [negative control (1), positive control (2), Khat treated (3) & Khat treated followed by xanthinol nicotinate (4)].	53

Table No.		Page
Table (16)	Statistical analysis of globulins concentrations (g/dl) in different groups of rats (negative control, positive control, Khat treated & Khat treated followed by xanthinol nicotinate).	54
Table (17)	One way analysis of variance (ANOVA) of globulins concentrations between different groups [negative control (1), positive control (2), Khat treated (3) & Khat treated followed by xanthinol nicotinate (4)].	54
Table (18)	Statistical analysis of Albumin/Globulins ratio in different groups of rats (negative control, positive control, Khat treated & Khat treated followed by xanthinol nicotinate).	55
Table (19)	One way analysis of variance (ANOVA) of Albumin/Globulins ratio between different groups [negative control (1), positive control (2), Khat treated (3) & Khat treated followed by xanthinol nicotinate (4)].	55



Part I

THEORETICAL

