

**ECOLOGICAL AND MORPHOLOGICAL
STUDIES ON SYCAMORE FIG PSYLLID,
PAUROPSYLLA WILLCOCKSI DÇBSKI
(HOMOPTERA - PSYLLOIDEA).**

BY

FATMA ABD EL-HALIM MOHARUM MOHAMMAD

B. Sc. AGRICULTURE

AIN SHAMS UNIVERSITY (1983)

A THESIS

**SUBMITTED IN PARTIAL FULFILLMENT OF THE
REQUIREMENTS FOR THE MASTER DEGREE**

IN

ENVIRONMENTAL SCIENCES

(DEPARTMENT OF BIOLOGICAL AND PHYSICAL SCIENCE.)

INSTITUTE OF ENVIRONMENTAL STUDIES

AND RESEARCH. AIN SHAMS UNIVERSITY

UNDER THE SUPERVISION OF :

Dr. SAMY M. EL-SAYES

**ASSISTANT PROFESSOR OF
BIOLOGICAL AND PHYSICAL
SCIENCE DEPARTMENT.**

**INSTITUTE OF
ENVIRONMENTAL STUDIES
AND RESEARCH-AIN SHAMS
UNIVERSITY**

Dr. ZEINAT K. MOHAMMAD

**SENIOR RESEARCHER OF
ENTOMOLOGY, PLANT
PROTECTION RESEARCH
INSTITUTE AGRICULTURE
RESEARCH CENTER**

CAIRO

1997

the 1990s, the number of people with a mental health problem has increased by 50% (Mental Health Foundation 1999). The prevalence of mental health problems in the community has increased from 1.5% in 1979 to 2.5% in 1994 (Mental Health Foundation 1999).

There is a growing awareness of the need to address the needs of people with mental health problems in the community. The Department of Health (1999) has set out a strategy for mental health care, which includes a commitment to 'improve the lives of people with mental health problems, and to prevent and reduce the incidence of mental health problems'. The strategy also includes a commitment to 'improve the effectiveness of mental health services, and to ensure that they are accessible to all who need them'.

The strategy also includes a commitment to 'improve the effectiveness of mental health services, and to ensure that they are accessible to all who need them'. The strategy also includes a commitment to 'improve the effectiveness of mental health services, and to ensure that they are accessible to all who need them'. The strategy also includes a commitment to 'improve the effectiveness of mental health services, and to ensure that they are accessible to all who need them'.

The strategy also includes a commitment to 'improve the effectiveness of mental health services, and to ensure that they are accessible to all who need them'. The strategy also includes a commitment to 'improve the effectiveness of mental health services, and to ensure that they are accessible to all who need them'. The strategy also includes a commitment to 'improve the effectiveness of mental health services, and to ensure that they are accessible to all who need them'.

The strategy also includes a commitment to 'improve the effectiveness of mental health services, and to ensure that they are accessible to all who need them'. The strategy also includes a commitment to 'improve the effectiveness of mental health services, and to ensure that they are accessible to all who need them'. The strategy also includes a commitment to 'improve the effectiveness of mental health services, and to ensure that they are accessible to all who need them'.

The strategy also includes a commitment to 'improve the effectiveness of mental health services, and to ensure that they are accessible to all who need them'. The strategy also includes a commitment to 'improve the effectiveness of mental health services, and to ensure that they are accessible to all who need them'. The strategy also includes a commitment to 'improve the effectiveness of mental health services, and to ensure that they are accessible to all who need them'.

The strategy also includes a commitment to 'improve the effectiveness of mental health services, and to ensure that they are accessible to all who need them'. The strategy also includes a commitment to 'improve the effectiveness of mental health services, and to ensure that they are accessible to all who need them'. The strategy also includes a commitment to 'improve the effectiveness of mental health services, and to ensure that they are accessible to all who need them'.

The strategy also includes a commitment to 'improve the effectiveness of mental health services, and to ensure that they are accessible to all who need them'. The strategy also includes a commitment to 'improve the effectiveness of mental health services, and to ensure that they are accessible to all who need them'. The strategy also includes a commitment to 'improve the effectiveness of mental health services, and to ensure that they are accessible to all who need them'.

The strategy also includes a commitment to 'improve the effectiveness of mental health services, and to ensure that they are accessible to all who need them'. The strategy also includes a commitment to 'improve the effectiveness of mental health services, and to ensure that they are accessible to all who need them'. The strategy also includes a commitment to 'improve the effectiveness of mental health services, and to ensure that they are accessible to all who need them'.



بِسْمِ اللَّهِ الرَّحْمَنِ الرَّحِيمِ

وَبِهِ نَسْتَعِينُ



the 1990s, the incidence of *S. flexneri* has increased in the United Kingdom [10]. In the United States, *S. flexneri* has been reported to be the most common serotype of *Shigella* isolated from children with shigellosis [11].

There is a paucity of data on the epidemiology of *S. flexneri* in the United Kingdom. In the 1970s, *S. flexneri* was the most commonly isolated *Shigella* serotype from patients with shigellosis in the United Kingdom [12]. In the 1980s, *S. flexneri* was the most commonly isolated *Shigella* serotype from patients with shigellosis in the United Kingdom [13]. In the 1990s, *S. flexneri* was the most commonly isolated *Shigella* serotype from patients with shigellosis in the United Kingdom [14].

In the 1990s, *S. flexneri* was the most commonly isolated *Shigella* serotype from patients with shigellosis in the United Kingdom [14]. In the 1990s, *S. flexneri* was the most commonly isolated *Shigella* serotype from patients with shigellosis in the United Kingdom [14]. In the 1990s, *S. flexneri* was the most commonly isolated *Shigella* serotype from patients with shigellosis in the United Kingdom [14].

In the 1990s, *S. flexneri* was the most commonly isolated *Shigella* serotype from patients with shigellosis in the United Kingdom [14]. In the 1990s, *S. flexneri* was the most commonly isolated *Shigella* serotype from patients with shigellosis in the United Kingdom [14]. In the 1990s, *S. flexneri* was the most commonly isolated *Shigella* serotype from patients with shigellosis in the United Kingdom [14].

In the 1990s, *S. flexneri* was the most commonly isolated *Shigella* serotype from patients with shigellosis in the United Kingdom [14]. In the 1990s, *S. flexneri* was the most commonly isolated *Shigella* serotype from patients with shigellosis in the United Kingdom [14]. In the 1990s, *S. flexneri* was the most commonly isolated *Shigella* serotype from patients with shigellosis in the United Kingdom [14].

In the 1990s, *S. flexneri* was the most commonly isolated *Shigella* serotype from patients with shigellosis in the United Kingdom [14]. In the 1990s, *S. flexneri* was the most commonly isolated *Shigella* serotype from patients with shigellosis in the United Kingdom [14]. In the 1990s, *S. flexneri* was the most commonly isolated *Shigella* serotype from patients with shigellosis in the United Kingdom [14].

In the 1990s, *S. flexneri* was the most commonly isolated *Shigella* serotype from patients with shigellosis in the United Kingdom [14].

BIOGRAPHY

NAME IN FULL : FATMA ABD EL-HALIM MOHARUM
MOHAMMAD

DATE AND PLACE OF BIRTH : 13 APRIL, 1961, GIZA

DATE OF GRADUATION : MAY 1983

DEGREE AWARDED : B. SC. AGRICULTURE.
AIN SHAMS UNIVERSITY

GRADE : GOOD

OCCUPATION : AGRICULTURAL SPECIALIST.
PLANT PROTECTION RESEARCH
INSTITUTE DOKKI CAIRO, EGYPT.

DATE OF APPIONTMENT : 1983

DATE OF REGISTRATION : 1997

APPROVAL SHEET

ECOLOGICAL AND MORPHOLOGICAL
STUDIES ON SYCAMORE FIG PSYLLID
PAUROPSYLLA WILLCOCKSI DÇBSKI
(HOMOPTERA- PSYLLOIDEA).

BY

FATMA ABD EL-HALIM MOHARUM MOHAMMAD

B. SC. AGRICULTURE

AIN SHAMS UNIVERSITY (1983)

THIS THESIS FOR M. SC. OF SCIENCE DEGREE HAS
BEEN APPROVED BY :

f. Dr. A. H. KASCHEF, Professor of Entomology Faculty of
Science, Ain Shams University

A. Karschef

f. Dr. F. M. MARIY, Professor of Economic Entomology, Faculty
of Agriculture, Ain Shams University.

F. Mariy

SAMY, M. EL-SAYES, Assistant Professor of Biological And
Physical Science Department. Institute of
Environmental Studies And Research-
Ain Shams University.

Samy

ZEINAT, K. MOHAMMAD, Senior Researcher of Entomology,
Plant Protection Research Institute
Agriculture Research Center.

Zeinatt

COMMITTEE IN CHARGE

DATE / /1997

The first part of the paper discusses the importance of the study of the history of the English language. It is a branch of linguistics which deals with the changes in the language over time. The second part of the paper discusses the importance of the study of the history of the English language. It is a branch of linguistics which deals with the changes in the language over time. The third part of the paper discusses the importance of the study of the history of the English language. It is a branch of linguistics which deals with the changes in the language over time. The fourth part of the paper discusses the importance of the study of the history of the English language. It is a branch of linguistics which deals with the changes in the language over time. The fifth part of the paper discusses the importance of the study of the history of the English language. It is a branch of linguistics which deals with the changes in the language over time. The sixth part of the paper discusses the importance of the study of the history of the English language. It is a branch of linguistics which deals with the changes in the language over time. The seventh part of the paper discusses the importance of the study of the history of the English language. It is a branch of linguistics which deals with the changes in the language over time. The eighth part of the paper discusses the importance of the study of the history of the English language. It is a branch of linguistics which deals with the changes in the language over time. The ninth part of the paper discusses the importance of the study of the history of the English language. It is a branch of linguistics which deals with the changes in the language over time. The tenth part of the paper discusses the importance of the study of the history of the English language. It is a branch of linguistics which deals with the changes in the language over time.

ACKNOWLEDGMENTS

Firstly, ultimate thanks to God

The authoress wishes to express her deepest thanks and gratitude to Dr. **Samy, M. Ali El-Sayes**, Assistant professor of biological and physical sciences Department, Institute of Environmental Studies and Research, Ain Shams University for his valuable supervision, guidance and revision of the manuscript.

My deepest thanks are also due to Dr. **Zeinat, K. Mohammad**, Senior researcher of Entomology, Plant Protection Research Institute for her kindness, unfailing help, valuable advice, suggesting the problem and encouragement as well as continuous guidance.

Thanks due to all the staff members of scale insects and mealy bugs research Department, Plant Protection Research Institute.

My deep thanks extend to all my family, for their kind help and providing all facilities throughout the work.

CONTENTS

SUBJECT	PAGE
INTRODUCTION.	2
LITERATURE REVIEW.	5
1) MORPHOLOGY.	7
2) ECOLOGICAL STUDIES.	11
MATERIALS AND METHODS.	14
1) MORPHOLOGY.	16
2) ECOLOGICAL STUDIES.	18
RESULTS. AND DISCUSSION	21
1) SYSTEMATIC POSITION.	23
2) MORPHOLOGICAL CHARACTERISTICS.	23
1- DESCRIPTION OF THE EGG	23
2- DESCRIPTION OF THE FIRST NYMPHAL INSTAR	26
3- DESCRIPTION OF THE SECOND NYMPHAL INSTAR	31
4- DESCRIPTION OF THE THIRD NYMPHAL INSTAR	36
5- DESCRIPTION OF THE FOURTH NYMPHAL INSTAR	41
6- DESCRIPTION OF THE FIFTH NYMPHAL INSTAR	46
3) KEY TO DIFFERENTIATE THE NYMPHAL INSTARS.	51
4) POPULATION DENSITY AND DURATION OF <u>PAUROPSYLLA</u> <u>WILLCOCKIS</u> DÇBSKI	

GENERATIONS UNDER FIELD CONDITIONS	52
THE ORIENTATION OF <u>PAUOPSYLLA WILLCOCKSI</u> DÇBSKI INFESTATION OF THE TWO LEAF SURFACES OF <u>FICUS SYCOMORUS</u> L. TREE TAKEN FROM FOUR CARDINAL DIRECTION AND THREE LEVELS.	57
1- THE ABUNDANCE OF <u>PAUOPSYLLA WILLCOCKSI</u> DÇBSKI NYMPHS IN THE FOUR CARDINAL DIRECTIONS OF THE FIG TREE <u>FICUS SYCOMORUS</u> L..	57
2-THE ABUNDANCE OF <u>PAUOPSYLLA WILLCOCKSI</u> DÇBSKI NYMPHS IN THREE HORIZONTAL LEVELS OF THE FIG TREE <u>FICUS SYCOMORUS</u> L..	62
3- DISTRIBUTION OF <u>PAUOPSYLLA WILLCOCKSI</u> DÇBSKI ON EITHER SIDES OF OF THE <u>FICUS SYCOMORUS</u> LEAF SURFACES :	65
EFFECT OF SOME ECOLOGICAL FACTORS ON THE SEASONAL ABUNDANCE OF <u>PAUOPSYLLA WILLCOCKSI</u> DÇBSKI EGGS:	68
1- TEMPERATURE.	70
2- RELATIVE HUMIDITY.	70
3- PHOTO-PERIOD.	72
4- DEW POINT.	72
5- WIND VELOCITY.	72
6-COMBINED EFFECT OF THE ECOLOGICAL FACTORS ON THE SEASONAL ABUNDANCE OF <u>PAUOPSYLLA WILLCOCKSI</u> DÇBSKI EGGS	74
EFFECT OF SOME ECOLOGICAL FACTORS ON THE SEASONAL ABUNDANCE OF <u>PAUOPSYLLA WILLCOCKSI</u> DÇBSKI NYMPHS.	77

