### A COMPARATIVE STUDY BETWEEN THE RESULTS OF PERCUTANEOUS NEPHROLITHOTOMY AND EXTRACORPOREAL SHOCK WAVE LITHOTRIPSY IN THE TREATMENT OF LOWER CALYCEAL STONES

#### THESIS

Submitted for partial fulfillment of M.Sc Degree in:

UROLOGY

Presented by

## Mohamed Hamdi Ibrahim

M.B., B.Ch

Under the supervision of

Prof. Dr. Mohamed Tarek M. Fathy Zaher

Professor of Urology

Faculty of Medicine

Ain Shams University

Assis. Prof. Dr. Wael Aly Maged

Assis. Prof. of Urology Faculty of Medicine Ain Shams University

Dr. Mohamed Shukry Shoeb

Lecturer in Urology Faculty of Medicine Ain Shams University

FACULTY OF MEDICINE AIN SHAMS UNIVERSITY 1999







~ (CP

June !



#### ACKNOWLEDGEMENT

To **ALLAH**, goes all my deepest gratitude and thanks for achieving any work in my life.

I wish to express my cordial appreciation and utmost gratitude to *Prof. Dr. Tarek Zaher*, Professor of Urology, Faculty of Medicine, Ain Shams University, for his sincere guidance, supervision and kind encouragement throughout the whole thesis.

My sincere thanks to Assis. Prof. Dr. Wael Maged, Assistant Professor of Urology, Faculty of Medicine, Ain Shams University, for initiation the idea for this thesis, as well as providing an invaluable criticism and direction.

I am indeed most grateful to *Dr. Mohamed Shoeb*, Lecturer in Urology, Faculty of Medicine, Ain Shams University, for his relentless support, splendid efforts and creative thoughts.

Last but not least, my deep appreciation is expressed to all the staff of Urology department, Ain Shams University, and to all members of my family for their co-operation and encouragement

Mohamed Hamdi



TO MY FAMILY ..



## INDEX

|      |  | Page |
|------|--|------|
| I    | Introduction and Aim of The Work                 | 1    |
| 11   | Review of Literature                             |      |
|      | Historical aspects of urinary stone disease      | 3    |
|      | Epidemiological aspects of urinary stone disease | 5    |
|      | Theories of stone formation                      | 9    |
|      | Types of urinary stones                          | 15   |
|      | Methods of diagnosis                             | 23   |
|      | Pathological consequences                        | 29   |
|      | Treatment modalities                             | 32   |
|      | Extracorporeal shock wave lithotripsy (ESWL)     | 35   |
|      | Percutaneous nephrolithotomy                     | 52   |
| 11   | Patients and Methods                             | 62   |
| Ш    | Results  | 70   |
| IV   | Cases  | 79   |
| V    | Discussion                                       | 90   |
| VI   | Summary and Conclusion                           | 98   |
| VII  | References                                       | 101  |
| VIII | Arabic summary                                   |      |



## LIST OF TABLES AND FIGURES

|    | Tables  | Page |
|----|---|------|
| 1  | Pertinent features of the three forms of Hypercalciuria       | 17   |
| 2  | Radiographic densities of stones in relation to bone          | 26   |
| 3  | Advantages and disadvantages of shock wave generators         | 41   |
| 4  | Comparison of imaging modalities                              | 42   |
| 5  | Ease of fragmentation of various calculi                      | 43   |
| 6  | Incidence of complications in our patients treated with ESWL  | 75   |
|    | Figures   |      |
| 1  | Saturation and supersaturation of solutes in urine            | 10   |
| 2  | Mechanism of stone fragmentation                              | 37   |
| 3  | Electrohydrolic lithotriptor shock wave generation            | 38   |
| 4  | Electromagnetic lithotriptor shock wave generation            | 39   |
| 5  | Piezoelectric shock wave generation                           | 40   |
| 6  | Siemens lithostar plus lithotriptor                           | 66   |
| 7  | Percutaneous neprholithotomy set                              | 69   |
| 8  | Electromagnetic lithoclast                                    | 69   |
| 9  | Incidence of acute and obtuse infundibulopelvic angle in ESWL | 71   |
| 10 | Incidence of narrow and wide calyceal neck in ESWL            | 72   |
| 11 | Percentage of stone-free patients in ESWL                     | 73   |
| 12 | Incidence of stone disintegration in PCNL                     | 76   |
| 13 | Success rate of PCNL  | 77   |

# INTRODUCTION AND AIM OF THE WORK