

بِسْمِ اللَّهِ الرَّحْمَنِ الرَّحِيمِ

”قَالُوا سُبْحَانَكَ لَا عِلْمَ لَنَا إِلَّا مَا عَلَّمْتَنَا  
إِنَّكَ أَنْتَ الْعَلِيمُ الْحَكِيمُ”

(صدق الله العظيم)

(سورة البقرة ٣٢)

# **A New Pulp Capping Material Developed From Portland Cement**

*Thesis*

Submitted in partial fulfillment for Doctor's Degree in  
Endodontics

By

**Ahmed Maged Negm**

(B.D.S) October 6 University 2007

(M.Sc) in Endodontics Cairo University 2012

Faculty of Dentistry  
Ain-Shams University  
2015

# **Supervisors**

## **Professor Dr Ihab El Sayed Hassanein**

Professor of Endodontics, Faculty of Dentistry Ain  
Shams University

## **Professor Dr Ashraf Mohamed Abdel-Rahman Abu- Seida**

Professor of Surgery, Anesthesiology and Radiology,  
Faculty of Veterinary  
Medicine, Cairo University

## **Dr Mohamed Mokhtar Nagy**

Lecturer in Endodontics, Faculty of Dentistry Ain Shams  
University

# Acknowledgments

I would like to express my sincere gratitude and appreciation to **Prof Dr Ihab El Sayed Hassanein** Professor of Endodontics, Faculty of Dentistry, Ain Shams University, for his sincerity, unsurpassed kindness, thoughtful guidance, extraordinary decency, unlimited help, care and support.

Countless thanks to **Professor Dr Ashraf Mohamed Abdel-Rahman Abu-Seida** Professor of Surgery, Anesthesiology and Radiology, Faculty of Veterinary Medicine, Cairo University, for his unlimited support, valuable scientific opinions and effort.

My deep gratitude and respect to **Dr Mohamed Mokhtar Nagy** Lecturer in Endodontics, Faculty of Dentistry Ain Shams University, for his sincere help, support and encouragement during this study.

I also would like to thank **Professor Dr Heba Farag**

Professor of Oral Pathology and Vice Dean of Faculty of Oral and Dental Medicine, Cairo University; for her great efforts and guidance during the histopathological phases of this research..

## **Dedication**

This work is dedicated to my parents for their endless love, support and encouragement.

**"Thank you very much, without you this work would not have been possible"**

# Contents

<b>Introduction.....</b>	<b>1</b>
<b>Review of Literature.....</b>	<b>2</b>
<b>Aim of the study.....</b>	<b>36</b>
<b>Materials and methods.....</b>	<b>37</b>
<b>Results.....</b>	<b>64</b>
<b>Discussion.....</b>	<b>117</b>
<b>Summary and Conclusion.....</b>	<b>129</b>
<b>Bibliography.....</b>	<b>134</b>

## List of Tables

<b>Table</b>	<b>Title</b>	<b>Page</b>
Table (1)	Criteria for classification of inflammatory cell response	61
Table (2)	Scoring system for dentin bridge formation	61
Table (3)	Criteria for classification of pulp calcification	63
Table (4)	Means and standard deviations (SD) of initial setting time (Min) for the different materials	64
Table (5)	Means and standard deviations (SD) of final setting time (Min) for the different materials	66
Table (6)	Means and standard deviations (SD) of compressive strength (MPa) for the different materials after 24 hrs.	68
Table (7)	Means and standard deviations (SD) of compressive strength (MPa) for the different materials after 21 days.	70
Table (8)	Means and standard deviations (SD) of diametral tensile strength (MPa) for the different materials after 24 hrs.	72
Table (9)	Means and standard deviations (SD) of diametral tensile strength (MPa) for the different materials after 21 days.	74
Table (10)	Means and standard deviations (SD) of the push out bond strength (MPa) for the different	76



	materials.	
Table (11)	pH values and their average at different times.	78
Table (12)	Differences in mean inflammatory cell count between different groups after 3 weeks.	101
Table (13)	Differences in mean inflammatory cell count between different groups after 3 months	103
Table (14)	The frequencies and percentages of inflammatory scores for different groups.	107
Table (15)	The frequencies and percentages of dentin bridge scores for different groups.	110
Table (16)	The frequencies and percentages of pulp stone scores for different groups.	115

## List of Figures

<b>Figure No.</b>	<b>Title</b>	<b>Page no.</b>
Figure (1)	Gilmore needle used for testing setting time.	39
Figure (2)	pH meter.	40
Figure (3)	Customized Teflon mould for formation of the specimens with the required size for compressive strength testing.	41
Figure (4)	Customized Teflon mould for formation of the samples with the required size for diametral tensile strength testing.	41
Figure (5)	Computer controlled materials testing machine (Model LRX-plus; Lloyd Instruments Ltd., Fareham, UK).	43
Figure (6)	Testing compressive strength of a vertically mounted specimen with computer controlled testing machine.	44
Figure (7)	Testing diametral tensile strength of a horizontally mounted specimen using computer controlled testing machine.	46
Figure (8)	A sample placed in a dentin disc for push out bond strength testing using computer controlled testing machine.	49
Figure (9)	A schematic diagram of the present study	51
Figure (10)	Intravenous administration of general anesthesia.	54

Figure (11)	Class V cavities with pulp exposure in a dog.	56
Figure (12)	Placement of capping material on the exposure sites.	57
Figure (13)	Placement of final restoration over the capping materials.	58
Figure (14)	Light microscope used for histologic examination.	60
Figure (15)	Image analyzer system (Leica Queen 500) connected to Leica light microscopy Leica Microsystems) used for measuring dentin bridge thickness.	62
Figure (16)	Bar chart showing the mean initial setting time (Min) for the different materials.	65
Figure (17)	Bar chart showing the mean final setting time (Min) for the different materials.	67
Figure (18)	Bar chart showing the mean compressive strength (MPa) for the different materials after 24 hrs.	69
Figure (19)	Bar chart showing the mean compressive strength (MPa) for the different materials after 21 days.	71
Figure (20)	Bar chart showing the mean diametral tensile strength (MPa) for the different materials after 24 hrs.	73
Figure (21)	Bar chart showing the mean diametral tensile strength (MPa) for the different materials after 21 days.	75
Figure (22)	Bar chart showing the mean push out bond strength (MPa) for the different materials.	

		77
Figure (23)	Diagrammatic representation of pH values for the different materials.	78
Figure (24)	Histogram showing the pH values of different materials at different times.	79
Figure (25)	Photomicrograph showing normal pulp. (H&E X100)	80
Figure (26)	Photomicrograph of MTA after 3weeks showing vasodilatation and a big area of necrosis with no dentin bridge formation. (H&E X40).	82
Figure (27)	Photomicrograph of MTA after 3weeks showing vasodilatation and continuous odontoblastic layer. (H&E X100).	82
Figure (28)	Photomicrograph of MTA 3weeks showing pulp with vasodilatation, and areas of necrosis. (H&E X200)	83
Figure (29)	Photomicrograph of MTA after 3months showing complete dentin bridge formation with normal pulp. (H&E X 100).	83
Figure (30)	Photomicrograph of MTA after 3 months showing complete dentin bridge formation with normal pulp. (H&E x 100).	84
Figure (31)	Photomicrograph of MTA after 3 months showing normal pulp with continuous odontoblastic layer. (H&E x 100).	84
Figure (32)	Photomicrograph of MTA after 3 months	

	showing dilated blood vessels and mild inflammation. (H&E x 200).	85
Figure (33)	Photomicrograph of PortCal I after 3 weeks showing the exposure site without dentin bridge formation. (H&E X100).	87
Figure (34)	Photomicrograph of PortCal I after 3 weeks showing the exposure site without dentin bridge formation and normal pulp. (H&E X100).	87
Figure (35)	Photomicrograph of PortCal I after 3 weeks showing vasodilatation and areas of necrosis. (H&E X100).	88
Figure (36)	Photomicrograph of PortCal I after 3 weeks showing continuous odontoblastic layer. (H&E X100).	88
Figure (37)	Photomicrograph of Portcal I after 3 months showing the exposure site with partial dentin bridge formation. (H&E X100).	89
Figure (38)	Photomicrograph of PortCal I after 3 months showing the exposure site with partial dentin bridge formation. (H&E X100).	89
Figure (39)	Photomicrograph of PortCal I after 3 months showing inflamed pulp under complete dentin bridge. (H&E X100).	90
Figure (40)	Photomicrograph of PortCal I after 3 months showing normal pulp under complete dentin bridge. (H&E X100).	90
Figure (41)	Photomicrograph of PortCal I after 3 months	91

	showing normal pulp. (H&E X100).	
Figure (42)	Photomicrograph of PortCal I after 3 months showing vasodilatation and areas of necrosis. (H&E X100).	91
Figure (43)	Photomicrograph of Portland cement with bismuth oxide after 3 weeks showing no dentin bridge formation. (H&E X100).	93
Figure (44)	Photomicrograph of Portland cement with bismuth oxide after 3 weeks showing vasodilatation and continuous odontoblastic layer. (H&E X100).	93
Figure (45)	Photomicrograph of Portland cement with bismuth oxide after 3 weeks showing vasodilatation and necrosis. (H&E X100).	94
Figure (46)	Photomicrograph of Portland cement with bismuth oxide after 3 weeks showing severe inflammation. (H&E X100).	94
Figure (47)	Photomicrograph of Portland cement with bismuth oxide after 3 months showing a failed attempt to form dentin bridge over a necrotic pulp. (H&E X100).	95
Figure (48)	Photomicrograph of Portland cement with bismuth oxide after 3 months showing the exposure site with partial dentin bridge formation. (H&E X100).	95
Figure (49)	Photomicrograph of Portland cement with bismuth oxide after 3 months showing the exposure site with complete dentin bridge	96

	formation over necrotic pulp. (H&E X100).	
Figure (50)	Photomicrograph of Portland cement with bismuth oxide after 3 months showing almost normal pulp. (H&E X100).	96
Figure (51)	Plate of photomicrographs of Portland cement with bismuth oxide after 3 months showing vasodilatation. (H&E X100).	97
Figure (52)	Plate of photomicrographs of Portland cement with bismuth oxide after 3 months showing vasodilatation with fibrosis. (H&E X100).	97
Figure (53)	Photomicrograph of Portland cement with bismuth oxide after 3 months showing pulp with pulp stone. (H&E X100).	98
Figure (54)	Photomicrograph of Portland cement with bismuth oxide after 3 months showing pulp with large detached pulp stone. (H&E X100).	98
Figure (55)	Photomicrograph of Portland cement with bismuth oxide after 3 months showing fatty degeneration. (H&E X100).	99
Figure (56)	Photomicrograph of Portland cement with bismuth oxide after 3 months showing pulp inflammation and necrosis. (H&E X100).	99
Figure (57)	Graph I: Floating error bar represents mean inflammatory cell count between different groups after 3 weeks.	102
Figure (58)	Graph II: Floating error bar represents mean inflammatory cell count between different	104