



شبكة المعلومات الجامعية

بِسْمِ اللَّهِ الرَّحْمَنِ الرَّحِيمِ





شبكة المعلومات الجامعية



شبكة المعلومات الجامعية

التوثيق الالكتروني والميكرو فيلم




شبكة المعلومات الجامعية

جامعة عين شمس

التوثيق الالكتروني والميكرو فيلم



نقسم بالله العظيم أن المادة التي تم توثيقها وتسجيلها
على هذه الأفلام قد أعدت دون أية تغيرات



يجب أن

تحفظ هذه الأفلام بعيداً عن الغبار

في درجة حرارة من 15 – 20 مئوية ورطوبة نسبية من 20-40 %

To be kept away from dust in dry cool place of
15 – 25c and relative humidity 20-40 %



شبكة المعلومات الجامعية



بعض الوثائق الأصلية تالفة



شبكة المعلومات الجامعية



بالرسالة صفحات

لم ترد بالأصل

The Significance of Some ANCAs in Rheumatoid Arthritis

*Thesis submitted for partial fulfillment of the
MD Degree in Internal Medicine*

Presented by

Heba Ahmed Hamdi Zaki

M.B., B.Ph., M.Sc. Ain Shams University

Under the supervision of

Prof. Dr. **Mohamed Fathy Tamara**

*Professor of Internal Medicine and Rheumatology
Faculty of Medicine-Ain Shams University*

Prof. Dr. **Mohamed A.M. Makhoul** Prof. Dr. **Malak Hassan Bahgat**

*Professor of Internal Medicine Faculty of Medicine-
Ain Shams University* *Professor of Internal Medicine
Faculty of Medicine-
Ain Shams University*

Dr. **Nihad Shoukry Shoueb**

*Lecturer of Internal Medicine
Faculty of Medicine-
Ain Shams University*

Dr. **Tarek Mahmoud El-Beguermy**

*Lecturer of Radiology
Faculty of Medicine-
Ain Shams University*

Faculty of Medicine
Ain Shams University
2000

B
7/1/2

بسم الله الرحمن الرحيم
قالوا سبحانك لا علم لنا إلا

ما علمنا

إنك أنت العليم الحكيم

صدق الله العظيم
سورة البقرة الآية ٣٢

Acknowledgment

First and foremost, thanks to God for giving me the power and strength to carry out this work.

*It is a great honor to express my sincere and deepest gratitude to Prof. Dr. **Mohamed Fathy Tamara**, Professor of Internal Medicine and Rheumatology, Ain Shams University, for his kind, fatherly attitude, guidance, close supervision and expert advice which made me able to complete this work.*

*Also I would like to express my extreme thanks to Prof. Dr., **Mohamed Ali Marei Makhlouf**, Professor of Internal Medicine, Ain Shams University, for his great support, patience and careful supervision.*

*Thanks are extended to Prof. Dr., **Malak Hassan Bahgat**, Professor of Internal Medicine, Ain Shams University for her kind supervision and support.*

*I feel deeply obliged to Dr. **Nihad Shoukry Shoueb**, Lecturer of Internal Medicine, Ain Shams University, for the great effort and time she devoted in guiding me throughout the work.*

*I am also deeply grateful to Dr. **Tarek Mahmoud El-Beguermy**, Lecturer of Radiology, Ain Shams University, for his guidance and valuable assistance.*

*I would also like to express my appreciation to my colleagues in the Rheumatology Unit and Laboratory especially Dr. **Hanan Ewais**, Dr. **Elham Barakat** and to the biochemist Dr. **Manal Hasanain** for their cooperation and help.*

Heba Hamdi

List of Contents

	Page No.
<i>Introduction and Aim of the Work</i>	1
<i>Review of Literature</i>	
Chapter I: Rheumatoid Arthritis.....	3
Chapter II: Rheumatoid Vasculitis.....	11
Chapter III: Anti-neutrophil cytoplasmic antibodies (ANCA's).....	38
Chapter IV: Capillary Microscopy.....	90
<i>Patients and Methods</i>	114
<i>Results</i>	128
<i>Discussion</i>	165
<i>Summary and Conclusion</i>	176
<i>References</i>	180
<i>Arabic Summary</i>	

List of Tables

	<i>Page No.</i>
<i>Table (1): Neutrophil granule populations and their protein content.</i>	39
<i>Table (2): Pathologic findings in renal tissue from 100 ANCA positive patients with glomerulonephritis.</i>	84
<i>Table (3): Major capillaroscopic changes in connective tissue disorders.</i>	113
<i>Table (4): American College of Rheumatology criteria for the diagnosis of rheumatoid arthritis.</i>	125
<i>Table (5): American College of Rheumatology revised criteria for classification of functional status in RA.</i>	125
<i>Table (6): Activity assessment score (Duke et al, 1983).</i>	126
<i>Table (7): Showing the clinical data of group I (lone RA).</i>	137
<i>Table (8): Showing the clinical data of group II (RV).</i>	138
<i>Table (9): Showing the age, sex and ESR 1st hour of the control group.</i>	139
<i>Table (10): Showing some laboratory collagen markers in the group I (RA).</i>	140
<i>Table (11): Showing some laboratory collagen markers in the group II (RV).</i>	141
<i>Table (12): Showing the titers of ANCAs as detected by ELISA in group I (lone RA).</i>	142
<i>Table (13): Showing the titers of ANCAs as detected by ELISA in group II (RV).</i>	143
<i>Table (14): Showing the titers of ANCAs as detected by ELISA in control</i>	144

<i>Table (15): Showing the capillaroscopic findings of group I (lone RA).</i>	<i>145</i>
<i>Table (16): Showing the capillaroscopic findings of group II (Rh. Vasculitis).</i>	<i>146</i>
<i>Table (17): Showing the capillaroscopic findings of control group.</i>	<i>147</i>
<i>Table (18): Shows the mean, standard deviation, minimum and maximum values for each variable in group I (lone RA).</i>	<i>148</i>
<i>Table (19): Shows the mean, standard deviation, minimum and maximum values for each variable in group II (Rheumatoid vasculitis).</i>	<i>148</i>
<i>Table (20): Shows the mean, standard deviation, minimum and maximum values for each variable in control group.</i>	<i>149</i>
<i>Table (21): Showing a comparative study for age, ESR, capillary number, length and width between group I (lone RA) and control group.</i>	<i>149</i>
<i>Table (22): Showing a comparative study for age, ESR, capillary number, length and width between group II (rheumatoid vasculitis) and control group.</i>	<i>149</i>
<i>Table (23): Showing a comparative study for age, disease duration (DD), ESR, capillary number, length and width between group I (lone RA) and group II (rheumatoid vasculitis).</i>	<i>150</i>
<i>Table (24): Showing a comparative study for clinical parameters between group I (lone RA) and group II (rheumatoid vasculitis).</i>	<i>150</i>
<i>Table (25): Showing a comparative study for functional class between group I (lone RA) and group II (RV).</i>	<i>150</i>
<i>Table (26): Showing a comparative study for rheumatoid factor between group I (lone RA) and group II (RV).</i>	<i>151</i>

<i>Table (27): Showing a comparative study for RW results between group I (lone RA) and group II (RV).</i>	151
<i>Table (28): Showing a comparative study for ANCAs between group I (RA) and control group.</i>	151
<i>Table (29): Showing a comparative study for ANCAs between group II (RV) and control group.</i>	152
<i>Table (30): Showing a comparative study between group I (lone RA) and group II (RV) regarding the number of +ve results of MPO and cANCA.</i>	152
<i>Table (31): Shows the differences between -ve and +ve MPO patients among group II (rheumatoid vasculitis).</i>	152
<i>Table (32): Showing a comparative study for RW results between +ve MPO and -ve MPO patients among group II (RV).</i>	153
<i>Table (33): Showing a comparative study for clinical parameters between -ve and +ve MPO patients among group II (RV).</i>	153
<i>Table (34): Comparative study as regards capillaroscopic findings between group I (lone RA) and group II (RV) by NCM.</i>	153
<i>Table (35): Comparative study as regards Doppler U/S wave findings between group I (lone RA) and group II (RV).</i>	154