

New trends in management Of Acute Aortic Dissection

An essay

Submitted for partial fulfillment of the master degree in
General surgery

By

Nader Nabil Naguib Moawad

MB.B.Ch.
Cairo University

Supervised by

PROF. DR. TAREK MOHAMED FARID ELBAHAAR

Professor of general and vascular surgery
Faculty of medicine, Ain Shams University

DR. TAREK YOUSSEF AHMED YOUSSEF

Lecturer of general and vascular surgery
Faculty of medicine, Ain Shams University

**Faculty of Medicine
Ain Shams University**

Acknowledgement

I would like to thank all people who helped me in this research for their dedication and effort. As without their conscientious and diligent work, the production of this research would not have been possible.

I would like to thank **Professor Dr. Tarek Mohamed Farid Elbahaar** for his continuing support and great guidance.

Also I would like to thank **Dr. Tarek Youssef Ahmed Youssef** for his useful advice and suggestions. I have really benefited from his experience and valuable recommendations.

I would like to express my gratitude for the limitless encouragement I received while developing this research from my family; my father, my mother and my brother, they were next to me during each step in this research.

Special thanks to my consultants and colleagues in the department of cardiac surgery in **National Heart Institute** for their great help and support.

Thanks go also to my friends for their helping me and enhancing my motivation to produce this achievement.

CONTENTS

• List of figures.....	III
• List of tables	VII
• List of algorithms	VIII
• List of abbreviations.....	IX
1) Introduction	1
2) Aim of the work	4
3) History.....	5
4) Anatomy of Aorta	7
5) Histology	19
6) Pathology.....	21
▪ Etiology	23
▪ Pathogenesis	31
▪ Staging and classification	38
▪ Natural history	45
7) Clinical presentation	48
8) Investigations	58
9) Treatment	104
▪ Introduction	104
▪ Immediate medical treatment	104
▪ Surgical treatment.....	108
▪ Interventional therapy.....	129
▪ Management of dissection related complications.....	138
▪ Long term follow up an treatment.....	142
10) References.....	148
11) Arabic summary	

List of figures

No.	Figure shows:	Page No.
1	Normal aortic anatomy	7
2	Superior view of the aortic valve	8
3	Ascending aorta and aortic arch	9
4	The aortic arch	12
5	The descending thoracic aorta	14
6	Descending thoracic aorta	15
7	Abdominal aorta	16
8	Branches of Abdominal Aorta	18
9	Intimal Tear	34

10	Classification of aortic dissection	39
11	Classification of aortic dissection	40
12	Plain chest x-ray exhibiting many features of acute type A dissection	60
13	Visualization of the aorta on transthoracic echocardiography (TTE)	67
14	TEE image of dissection of the ascending aorta	69
15	TEE image of dissection of the descending aorta.	70
16	TEE image of dissection of the aortic arch	71
17	TEE image of dissection of the descending aorta complicated by pleural effusion (PL)	72
18	Intraoperative TEE reveals prolapse of the dissection flap through the aortic valve in the mid esophageal long axis view of the ascending aorta	72
19	Aortic dissection: Linear artifacts into ascending aorta	74
20	Axial image of CT arteriogram showing an intramural haematoma of the descending thoracic aorta	77

21	Coronal reformatted image and oblique reformatted image from contrast-enhanced, cardiac-gated CT in a patient with type A aortic dissection	80
22	Coronal reformatted image and an axial image from contrast enhanced, cardiac-gated CT in a patient with type B aortic dissection.	80
23	Axial contrast-enhanced CT images of left brachiocephalic vein simulating dissection of the brachiocephalic artery	83
24	Axial contrast-enhanced image of the aortic arch shows low attenuation of the IMH, compared to the contrast filled true lumen of the aorta	85
25	Axial CT images of calcified intraluminal thrombus simulating detached intima calcification	86
26	Multiphase reformatted contrast-enhanced CT images after replacement of the ascending thoracic aorta and aortic arch with a graft that extends into the descending thoracic aorta	87
27	Sagittal contrast-enhanced magnetic resonance image of type B aortic dissection	91
28	Aortogram of acute type B dissection	96
29	The false lumen of the distal aorta is closed and the aortic wall is reconstructed with inside and outside felt strips	118
30	The distal dissected aortic wall is reconstructed with inside and outside felt strips	118
31	Everting 2-0 pledgeted mattress sutures are placed shoulder to shoulder around the aortic annulus	120

32	The coronary ostia are attached to the graft by the button technique	121
33	Total aortic arch replacement	122
34	The brachiocephalic vessels are separated from the true lumen by the dissected false lumen	123
35	Percutaneous balloon fenestration	133
36	Completion aortogram after successful endovascular aortic repair	135
37	Malperfusion of the distal aorta by occlusive type B dissection	141

List of tables

No.	Table shows:	Page No.
1	Risk conditions for aortic dissection	30
2	Sensitivity and Specificity of Various Imaging Modalities Useful for the Diagnosis of Thoracic Aortic Dissection	101
3	Effect of temperature on cerebral metabolic rate	114
4	Considerations for surgical, medical, and interventional therapy in aortic dissection	137
5	Frequency and Location of Malperfusion in Acute Type A and B Thoracic Aortic Dissection	139

List of algorithms

No.	Algorithm shows:	Page No.
1	Acute Aortic Dissection evaluation pathway	103
2	Surgical management pathway for Acute Aortic Dissection	128
3	Acute Aortic Dissection management pathway	147

List of abbreviations

AAD	Acute Aortic Dissection
ACS	Acute Coronary Syndrome
AMI	Acute Myocardial Infarction
BAV	Bicuspid Aortic Valve
CHF	Congestive Heart Failure
CRP	C – reactive protein
CT	Computed Tomography
DAS	Data Acquisition System
DHCA	Deep Hypothermic Circulatory Arrest
EDS	Ehlers Danlos Syndrome
ELISA	Enzyme Linked Immunosorbant Assay
FBN 1	Fibrillin-1 protein
ICU	Intensive Care Unit
IMH	Intramural Haematoma
IRAD	International Registry of Acute Dissection
IVUS	Intravascular Ultrasound
MDCT	Multi-detector Computed Tomography

MDCTA	Multi-detector Computed Tomography Angiography
MFS	Marfan Syndrome
MPR	Multi-planar Reformatted
MRI	Magnetic Resonance Imaging
ROI	Region Of Interest
SD	Standard Deviation
SVC	Superior Vena Cava
TAV	Tricuspid Aortic Valve
TEE	Transoesophageal echocardiography
TTE	Transthoracic echocardiography

Introduction

INTRODUCTION

In 1760, Dr Nicholls, physician to King George II, first described on necropsy an acute aortic dissection. Over the course of more than 200 years, we have developed a growing awareness and understanding of acute and chronic aortic syndromes. Recent advances in imaging and therapeutic techniques have further emphasized the importance of early diagnosis of acute aortic syndromes, which continues to be crucial to survival. However, we continue to be uncomfortable in the diagnosis and management of this highly lethal disorder.

(Thomas et al., 2005)

Aortic dissection develops from a tear within the intima of the aortic wall, blood flows across this entry point into a weakened media splitting the medial layer along the direction of flow creating a new (false) channel within the aortic media. This new channel progresses downstream, and significant pressure/mechanical stress is exerted by the advancing column of blood on the aortic branches encountered in its path.

(Alberto and Joseph, 2007)

The true prevalence of aortic dissection is difficult to estimate. The incidence of aortic dissection is estimated to be 5-30 cases per 1 million people per year. Aortic dissection occurs once per 10,000 patients admitted to the hospital.

(Gale, 2008)

There are many risk conditions for aortic dissection; the most common predisposing factor in IRAD (International Registry of Acute Aortic Dissection) was hypertension (72%), a history of atherosclerosis was present in 31% of patients and a history of cardiac surgery in 18%.

(Thomas et al., 2005)

The DeBakey classification comprises 3 types of aortic dissection (I, II, and III); the Stanford classification, 2 types (A and B). Within the DeBakey system, type I denotes a dissection that involves the full span of the aorta; type II, a dissection from the aortic root through the level of the origin of the innominate artery; and type III, a dissection from the descending aorta (distal to the origin of the left subclavian artery) to the aortic bifurcation. Dissections in the Stanford classification are categorized based on the involvement of the ascending aorta (type A) or the lack of such involvement (type B).

(Vijay et al., 2009)

The challenge in managing acute aortic dissection is appropriate clinical suspicion and action in pursuing the diagnosis and therapy.

(Nesser et al., 2002)

Initial medical management has been the consensus for the treatment of acute type B aortic dissection unless associated with life-threatening complications.

(Anthony et al., 2006)

The primary objective in management is to normalize blood pressure and initiate anti-impulse medication thus reducing the force of left ventricular ejection (dP/dt), which are the main determinants of extension and rupture of the false lumen. Acute ascending aortic dissections (Stanford type A or De Bakey type I or II) should be treated as a surgical emergency because these patients are at high risk of life-threatening complications such as aortic rupture, stroke, visceral ischemia, cardiac tamponade, and circulatory failure. In the current era, endovascular interventions for acute descending (type B) aortic dissection is the primary approach in the setting of complications of acute distal dissection because surgical repair has no proven superiority over medical or interventional treatment in this scenario. Medical treatment focuses on invasive hemodynamic monitoring, β -blockade, and vasodilators.

(Hussein and Christoph, 2007)